

7401-216

7401-216

ACCEPTED
 AUG 7 1973
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR PESTICIDE PHOENIX REGISTERED
 ED UNID. R. NO. SUBJECT
 TO ATTACHED COMMENTS.

VOLUNTARY PURCHASING GROUP, Inc.
 Box 460
 Bonham, Texas 75418

PM 03

7401-216

8-7-73

182



ferti·lome

DIAZINON[®]
INSECT SPRAY

WARNING KEEP OUT OF REACH OF CHILDREN.
 See Back Panel for Additional Warnings

CROP	INSECTS	APPLY UP TO DAYS OF HARVEST	REMARKS
Peaches, Nectarines, Apricots	Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites,	20	Apply as full cover spray. Apply when aphids, mites and crawlers first appear. For borers apply to peaches, in May and June. For leathoppers (peaches only) apply as infestations develop. For apricot mealybug apply as a cover spray from petal fall to June.
	Peach Twig Borers, Scale	10	
	Crawlers, San Jose Scale Crawlers, Leafhoppers (peaches only), Apricot Mealybugs.	10	
Apples	Leafhoppers, Eye-Spotted Bud Moths, Fruit Tree Leaf Rollers, Mealybugs.	14	Apply as full cover spray. For leafhoppers apply as infestations develop. For Eye-Spotted Bud Moth, Fruit Tree Leaf Rollers and Mealybugs make 3-8 cover spray applications beginning in pink stage. Slight russetting may occur on some variety of apples, such as Golden Delicious.
Pears	Pear Psylla, Tentiform Leaf Miners, Mealybugs, San Jose Scale Crawlers, Forbs Scale Crawlers.	14	Apply as full cover spray. For Pear Psylla mix 3 tablespoons in 3 gallons of water, spray at pre-pink stage. Make 1-2 additional applications during the first and second cover sprays if necessary. For Tentiform Leaf Miners and Mealybugs apply in 3 cover sprays, beginning in April.
Grapefruit Lemons Limes Oranges Tangerines Langerines, Citrus Citron Kumquats and hybrids of these	Citrus Aphids, Fruit Tree Leaf Rollers, Cottony-Cushion Scale Crawlers, Soft Scale Crawlers, Citrus Thrips, Citrus Snow Scales.	21	For citrus aphids, Cottony-Cushion Scale Crawlers and soft scale crawlers and thrips, apply thorough cover spray when insects first appear. For Leaf Rollers apply when 50% of eggs in egg masses have hatched in spring. For citrus snow scale, apply as a postbloom spray and again as a summer spray. For heavy snow scale infestations apply 4 tablespoons in 3 gallons of water. Thorough coverage of tree trunk, limbs and twigs is essential. Do not apply during bloom.
Pecans	Aphids, Mites, Pecan Nut Casebearers, Walnut Caterpillars.		Apply as a thorough cover spray. Do not apply after husks crack.

VEGETABLES: Mix 2 tablespoons in 3 gallons of water unless otherwise specified, apply to 1000 sq. ft. Assure thorough coverage of upper and lower leaf surfaces. Repeat as necessary to maintain control.

THE FERTILIZER COMPANY, Inc.
Box 400
Kalamazoo, Michigan 49001

1. WHAT IS THIS PACKAGE FOR?

This insecticide controls major insects fast and for extended periods of time. It will greatly aid in your Home Garden insect problems.

2. HOW DO YOU USE IT?

Mix 2 teaspoons in 1 gallon of water unless otherwise specified. Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Penetrate dense foliage.

ROSES AND FLOWERS: ROSES, AZALEAS, CAMELLIAS, RHODODENDRON, CHRYSANTHEMUMS, CARNATIONS, LILAC. To kill Aphids; Cutworms; Evonymus Scale Crawlers; Webworms; Flea Beetles; Lace Bugs; Dipterous Leaf Miners; Leafhoppers; Mealybugs; Two-spotted Spider, European Red, Cyclamen, Carnation Bud, Carnation Shoot Mites; Thrips and Whiteflies.

SHRUBS AND TREES: AZALEA, CAMELLIA, HOLLY, ARBORVITAE, BIRCH, BOXWOOD, JUNIPER, RHODODENDRON, DOUGLAS FIR, ELM, LILAC, MIMOSA, OAK, PINE, POPLAR, SPRUCE and WILLOW. To kill Aphids, European Red, Two-spotted Spider, Clover, Cyclamen and Privet Mites; Juniper and Mimosa Webworms; Cottony-cushion, Lecanium, Pine Needle, Oak Pit Scale, San Jose and Soft Scale Crawlers; Whiteflies; Arborvitae, Azalea, Birch, Boxwood, Dipterous, Holly and Oak Leaf Miners; Leafhoppers; Mealybugs; Obscure Root Weevils and Omnivorous Leaf Tiers. For Mimosa Webworms apply a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals. NOTE: Do not use on ferns, poinsettias, hibiscus, pteris, gardenias and orchids since plant injury may occur.

LAWNS: ST. AUGUSTINE, BERMUDA, BLUEGRASS, DICHONDRA, BENT, FESCUS, and MARION. To kill Sod Webworms, Chinch Bugs, Cutworms, Armyworms, Lawn Moths, Ants, Chiggers, Fleas, Earwigs, Pillbugs, Crickets, Mites (both Bermuda and Clover), Springtails, Ticks, Bermuda Grass Thrips, Frit Flies, Lawn Billbugs and Leafhoppers, water grass thoroughly a few hours before applying. Mix 2 tablespoons in 3 gallons of water and apply to 125 sq. ft. or 1 cup in 24 gallons of water to 1000 sq. ft. NOTE: For Lawn Billbugs, water lawn thoroughly immediately after application. In the north, Chinch Bugs will be killed using 1 tablespoon in 3 gallons of water per 125 sq. ft.

FRUIT, CITRUS AND PECANS: Mix 2 tablespoons in 3 gallons of water. Apply as a full cover spray unless otherwise specified. Assure thorough coverage. Repeat as necessary to maintain control.