

CROP	INSECTS	APPLY UP TO DAYS OF HARVEST	REMARKS
Peaches Nectarines Apricots	Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Peach Twig Borers, Scale Crawlers, San Jose Scale Crawlers, Leafhoppers (peaches only), Apricot Mealybugs.	20 10 10	Apply as full cover spray. Apply when aphids, mites and crawlers first appear. For borers apply to peaches, in May and June. For leafhoppers (peaches only) apply as infestations develop. For apricot mealybug apply as a cover spray from petal fall to June.
Apples	Leafhoppers, Eye-Spotted Bud Moths, Fruit Tree Leaf Rollers, Mealybugs.	14	Apply as full cover spray. For Leafhoppers apply as infestations develop. For Eye-Spotted Bud Moth, Fruit Tree Leaf Rollers and Mealybugs make 3-8 cover spray applications beginning in pink stage. Slight russeting may occur on some variety of apples, such as Golden Delicious.
Pears	Pear Psylla, Teniform Leaf Miners, Mealybugs, San Jose Scale Crawlers, Forbs Scale Crawlers.	14	Apply as full cover spray. For Pear Psylla mix 3 tablespoons in 3 gallons of water, spray at pre-pink stage. Make 1-2 additional applications during the first and second cover sprays if necessary. For Teniform Leaf Miners and Mealybugs apply in 3 cover sprays, beginning in April.
Grapefruit Lemons Limes Oranges Tangelos Tangerines, Citrus Citron Kumquats and hybrids of these	Citrus Aphids, Fruit Tree Leaf Rollers, Cottony-Cushion Scale Crawlers, Soft Scale Crawlers, Citrus Thrips, Citrus Snow Scales.	21	For citrus aphids, Cottony-Cushion Scale Crawlers and soft scale crawlers and thrips, apply thorough cover spray when insects first appear. For Leaf Rollers apply when 50% of eggs in egg masses have hatched in spring. For citrus snow scale, apply as a postbloom spray and again as a summer spray. For heavy snow scale infestations apply 4 tablespoons in 3 gallons of water. Thorough coverage of tree trunk, limbs and twigs is essential. Do not apply during bloom.
Pecans	Aphids, Mites, Pecan Nut Casebearers, Walnut Caterpillars.		Apply as a thorough cover spray. Do not apply after husks crack.
Pole Beans Snap Beans	Black Bean Aphids, Mexican Bean Beetles, Leaf Miners.		Do not apply within 7 days of harvest of beans. Bean and pea forage or bean and pea hay may be fed to dairy and beef cattle and sheep.
Lima Beans	Dipterous Leaf Miners, Leafhoppers, Mites		Wait 4 days following treatment if forage is cut for hay. Peas may be
Peas	Aphids, Leaf Miners		picked immediately.
Cucumbers	Aphids, Sepentine Leaf Miners, Thrips		Do not apply within 7 days of harvest.
Tomatoes	Aphids, Dipterous Leaf Miners, Banded Cucumber Beetles, Fall and Southern Armyworms, Vinegar Flies (Drosophila spp.)		Do not apply within 1 day of harvest. For aphids and dipterous leaf miners mix 1 tablespoon in 3 gallons of water. For Aphid, Dipterous Leaf Miners and Vinegar Flies weekly applications are usually required.

2

Continued on page 3



Diazinon[®] \$12⁹⁸ Insect Spray

- PROVIDES POSITIVE CONTROL OF LISTED INSECTS ON ROSES, SHRUBS, ORNAMENTALS, SHADE TREES, LAWNS, FLOWERS, FRUIT AND NUT TREES AND VEGETABLES
- QUICK-KILLING, LONG-LASTING INSECTICIDE-MITICIDE. SCALCIDE CONTROLS MANY SAP SUCKING AND LEAF CHEWING INSECTS

2 TABLESPOONS MAKES 1 GALLON OF SPRAY

Active Ingredient:

O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	25%
Inert Ingredients	75%
Total	100%

NET CONTENTS ONE QUART

**KEEP OUT OF REACH OF CHILDREN
WARNING**

See Back Panel for Additional Warnings

ACCEPTED
NOV 20 1981
under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 7401-216

PM 03

7401-216

11/20/81

192

3. HERE ARE THE RESULTS YOU MAY EXPECT!

This new product, used as directed, will control listed insects in your lawn and garden.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be harmful or fatal if swallowed or absorbed through the skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating and smoking. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or household pets to go onto sprayed grass or come in contact with foliage until spray has completely dried. Food utensils such as a teaspoon or measuring cup should not be used for food purposes after use with insecticides.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed— Call a physician or Poison Control Center immediately. If possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled— Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin— Remove contaminated clothing and wash affected areas with soap and water.

If in eyes— Flush eyes with plenty of water. Call a physician immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated areas. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treated areas. Applications should be timed to provide maximum possible interval between treatment and the next period of bee activity.

STORAGE AND DISPOSAL

Do not reuse empty container bottle. Wrap container and put in trash collection.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

*Diazinon Reg. Trademark CIBA—GEIGY CORP.
VOLUNTARY PURCHASING GROUPS, INC.

EPA Est. 7401-TX-1

EPA Reg. No. 7401-216
BONHAM, TEXAS

14M-8-81

3

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

1. WHAT IS THIS PACKAGE FOR?

This insecticide controls major insects fast and for extended periods of time. It will greatly aid in your Home Garden insect problems.

2. HOW DO YOU USE IT?

Mix 2 teaspoons in 1 gallon of water unless otherwise specified. Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Penetrate dense foliage.

ROSES AND FLOWERS: ROSES, AZALEAS, CAMELLIAS, RHODODENDRON, CHRYSANTHEMUMS, CARNATIONS, LILAC. To kill Aphids, Cutworms, Euonymus Scale Crawlers, Webworms, Flea Beetles, Lace Bugs, Dipterous Leaf Miners, Leafhoppers, Mealybugs, Two-spotted Spider, European Red, Cyclamen, Carnation Bud, Carnation Shoot Mites, Thrips and Whiteflies.

SHRUBS AND TREES: AZALEA, CAMELLIA, HOLLY, ARBORVITAE, BIRCH, BOXWOOD, JUNIPER, RHODODENDRON, DOUGLAS FIR, ELM, LILAC, MIMOSA, OAK, PINE, POPLAR, SPRUCE and WILLOW. To kill Aphids, European Red, Two-spotted Spider, Clover, Cyclamen and Privet Mites, Juniper and Mimosa Webworms, Cottony-cushion, Lecanium, Pine Needle, Oak Pit Scale, San Jose and Soft Scale Crawlers, Whiteflies, Arborvitae, Azalea, Birch, Boxwood, Dipterous, Holly and Oak Leaf Miners, Leafhoppers, Mealybugs, Obscure Root Weevils and Omnivorous Leaf Tiers. For Mimosa Webworms apply a thorough foliar spray when adults first appear. Make two additional applications at 4—5 week intervals. NOTE: Do not use on ferns, poinsettias, hibiscus, pileas, gardenias and orchids since plant injury may occur.

LAWNS: ST. AUGUSTINE, BERMUDA, BLUEGRASS, DICHONDRA, BENT, FESCUS, and MARION: To kill White Grubs of (Japanese Beetles, European Chafer, Southern Chafer), Sod Webworms, Chinch Bugs, Cutworms, Armyworms, Lawn Moths, Ants, Chiggers, Fleas, Earwigs, Pillbugs, Crickets, Mites (both Bermuda and Clover), Springtails, Ticks, Bermuda Grass Thrips, Frit Flies, Lawn Billbugs and Leafhoppers, water grass thoroughly a few hours before applying. Mix 2 tablespoons in 3 gallons of water and apply to 125 sq. ft. or 1 cup in 24 gallons of water to 1000 sq. ft. NOTE: For Lawn Billbugs, water lawn thoroughly immediately after application. In the north, Chinch Bugs will be killed using 1 tablespoon in 3 gallons of water per 125 sq. ft.

FRUIT, CITRUS AND PECANS: Mix 2 tablespoons in 3 gallons of water. Apply as a full cover spray unless otherwise specified. Assure thorough coverage. Repeat as necessary to maintain control.

VEGETABLES: Mix 2 tablespoons in 3 gallons of water unless otherwise specified, apply to 1000 sq. ft. Assure thorough coverage of upper and lower leaf surfaces. Repeat as necessary to maintain control.

IMPORTED FIRE ANT: To aid in control of Imported Fire Ants in lawn and other recreational areas, apply 1 fl. oz. (2 tbsp.) in a minimum of 1 gallon of water. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Apply gently at a rate of 1 gallon per mound. High pressure sprays may disturb ants and cause migration, reducing produce effectiveness.

1

DIRECTIONS LIFT HERE

282