

1. WHAT IS THIS PACKAGE FOR?

Unique formulation that controls many fungal diseases found in home gardens

2. HOW DO YOU USE IT?

SHAKE WELL BEFORE USE OR SAMPLING: Unless otherwise specified mix 2 tablespoons in 1 gallon of water. Using hose on sprayer apply in such a manner to cover both leaf surfaces. (See special directions for citrus). For best results carefully follow spray schedules under individual headings. NOTE: Do not combine with other insecticide sprays.

LAWNS: (Nonpastured areas only) For Brown Patch, Brown Spot (St. Augustine only), Copper Spot, Leaf Spot, Melting Out, Seedling Blights mix 2 tablespoons in 1 gallon of water and apply to 100 sq. ft. of lawn area. Begin when growth starts in spring repeating at 7-10 day intervals throughout season. Do not graze treated areas. Do not feed clippings to livestock.

FRUIT TREES:

APPLES: For Bitter Rot, Black Rot, Frogeye Leaf Spot, Black Pox, White Rot apply in late cover and preharvest periods. For Botrytis Rot, Bullseye Rot apply in first and second cover sprays and in 1 or 2 pre-harvest sprays or as needed at 5-7 day intervals from prebloom through early cover periods. For Brooks Fruit Spot & Flyspeck apply in cover periods. Note: Red Delicious and other sensitive varieties may be injured by early season applications.

PEACHES: For Brown Rot, Rhizopus, Scab apply in pink, full bloom, shuck shed, cover and preharvest periods. For Brown Rot (blossom blight) apply at 3-4 day intervals during bloom. Do not apply within 1 day of harvest.

PEARS: For Fruit Spot, Scab (primary infection) apply in pink, petal fall and first two cover periods. For Scab (secondary infection) apply in cover periods. Notes: Do not use on D'Anjou pears. Russetting may be produced on Bosc pears.

CHERRIES: For Botrytis Rot apply at petal fall, shuck shed and in cover sprays. For Brown Rot (blossom blight) apply at 3-4 day intervals during bloom.

APRICOTS: For Brown Rot Twig Blight, Jacket Rot apply in red bud Bloom, 75% petal fall and cover periods.

CITRUS: For Brown Rot on grapefruit, lemons, limes, oranges, tangerines apply to skirts of trees before winter rains to a height of 3-4 feet. Repeat 10-12 weeks later.

VEGETABLES:

TOMATOES: For Anthracnose, Early and Late Blights, Gray and Septoria Leaf Spot mix 4 tablespoons in 1 gallon of water. Apply at first sign of diseases, repeating at 5-7 day intervals.

BEANS: For Anthracnose, Downy Mildew, Rust mix 1 tablespoon in 1 gallon of water. Apply before diseases appear, repeating at 4-10 intervals as necessary.

BEETS: For Alternaria, Cercospora, Septoria Leaf Spots, apply before diseases appear repeating at 7-10 day intervals.

POTATOES: For Early and Late Blights mix 4 tablespoons in 1 gallon of water. Apply when plants are 2-4 inches high, repeating at 5-7 day intervals throughout season.

ONIONS: For Downy Mildew, Purple Blotch apply at first sign of disease, Repeat at 7-10 day intervals.

CANTALOUPE, CUCUMBERS, HONEYDEW MELONS, PUMPKINS, SUMMER AND WINTER SQUASH, WATERMELONS: For Angular Leaf Spot, Anthracnose, Downy Mildew mix 3 tablespoons in 1 gallon of water, apply at first sign of disease, repeating at 5-7 day intervals.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

Control of listea fungal disease is easily obtained with use of this product.

CAUTION: Keep out of reach of children. Do not take internally. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Do not contaminate feed or foodstuffs.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Food utensils such as tablespoons or measuring cups should not be used for food purposes after use with pesticides.

Do not reuse container for any purpose. Destroy when empty. Product freezing temperature 32° F. Satisfactory utility is restored after thawing and mixing of product. Repeated 0-10° freezing may impair utility.

Buyer assumes all risk of use, storage or handling of this material in strict accordance with directions given herewith.

28M 1 93

EPA Reg. No. 7401-1-1

VOLUNTARY PURCHASING GROUPS, INC.
BONHAM, TEXAS



ferti·lome



LAWN & FRUIT TREE DISEASE CONTROL \$2.98

- FOR CONTROL OF DISEASES ON LAWNS, FRUIT & CITRUS TREES, VEGETABLES AND MELONS.
- CONTROL OF BROWN PATCH, BROWN SPOT, MELTING OUT (ST. AUGUSTINE ONLY), LEAF SPOTS, MIDGE ROT, BLIGHTS, MELT OUT, AND UNCERTAIN OTHER FUNGI.
- DIRECTIONS MAKE 1 GALLON OF SPRAY.
- Contains:
 - Control 13%
 - inert ingredient 87%
- N (Trichloromethylthio-4-cyclohexene-1,2-dicarboximide)
- NET CONTENTS 16 FL. OZS (ONE PINT)

CAUTION. KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

ACCEPTED UNDER THE FEDERAL FUNGICIDE AND ECONOMIC INSECTICIDE ACT

Oct 18, 1973 7901-232

VOLUNTARY PURCHASING GROUPS, INC. BONHAM, TEXAS

