

ferti-lome

Lawn & Fruit Tree

Disease Control

ACCEPTED
Per self Cert
OCT 13 1987
The Federal Insecticide
Fungicide and Rodenticide
Act, as amended, under
EPA Reg. No. 7401-232

For control of disease on lawns, fruit trees, vegetables and melons.

Controls brown patch, brown spot (on St. Augustine Only), leaf spots, mildews, rots, blights, melting out, and certain other fungi


2 tablespoons makes 1 gallon of spray

Active Ingredients:

- *Captan..... 12.71%
- Related Derivatives..... 0.29%
- Inert Ingredients:..... 87.00%
- *N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

Keep Out Of Reach Of Children
DANGER

See back panel for additional precautionary statements



(Back Panel)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. WHAT IS THIS PACKAGE FOR?

Unique formulation that controls many fungal diseases found in home gardens.

2. HOW DO YOU USE IT?

SHAKE WELL BEFORE USE OR SAMPLING: Unless otherwise specified mix 2 tablespoons in 1 gallon of water. Using a hose-on sprayer apply in such a manner to cover both leaf surfaces. For best results carefully follow spray schedules under individual headings. NOTE: Do not combine with other insecticide sprays.

LAWNS (Nonpastured areas only). For Brown Patch, Brown Spot (St. Augustine Only), Copper Spot, Leaf Spot, Melting Out, Seedling Blights mix 2 tablespoons in 1 gallon of water and apply to 100 sq. ft. of lawn area. Begin when growth starts in spring repeating at 7-10 day intervals throughout season. Do not graze treated areas. Do not feed clippings to livestock.

FRUIT TREES:

APPLES... For Bitter Rot, Black Rot, Frogeye Leaf Spot, Black Pox, White Rot apply in late cover and preharvest periods. For Botrytis Rot, Bulls-eye Rot, apply in first and second cover sprays and in 1 or 2 preharvest sprays or as needed at 5-7 day intervals from prebloom through early cover periods. For Brooks Fruit Spot & Flyspeck apply in cover periods. NOTE: Red Delicious and other sensitive varieties may be injured by early season applications.

PEACHES... For Brown Rot, Rhizopus, Scab apply in pink, full bloom, shuck shed, cover and preharvest periods. For Brown Rot (blossom blight) apply at 3-4 day intervals during bloom. Do not apply within 1 day of harvest.

PEARS... For Fruit Spot, Scab (primary infection) apply in pink, petal fall and first two cover periods. For Scab (secondary infection) apply in cover periods. NOTES: Do not use on D'Anjou pears. Russetting may be produced on Bosc pears.

CERRIES... For Botrytis Rot apply at petal fall, shuck shed and in cover sprays. For Brown Rot (blossom blight) apply at 3-4 day intervals during bloom.

APRICOTS... For Brown Rot, Twig Blight, Jacket Rot apply in red bud, bloom, 75% petal fall and cover periods.


VEGETABLES:

TOMATOES... For Anthracnose, Early and Late Blights, Gray and Septoria Leaf Spot mix 4 tablespoons in 1 gallon of water. Apply at first sign of disease, repeating at 5-7 day intervals.

CANTALOUPEs, CUCUMBERS... For Angular Leaf Spot, Anthracnose, Downy Mildew mix 3 tablespoons in 1 gallon of water, apply at first sign of disease, repeating at 5-7 day intervals.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

Control of listed fungal diseases is easily obtained with use of this product:



STORAGE: Store in original container and place in a locked storage area.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Discard container in trash.

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse. Avoid contact with skin or clothing. Wear chemical resistant gloves when using indoors or outdoors. When using wear long pants and long-sleeved shirt and apply with the wind to your back. Wash non-disposable gloves thoroughly with soap and water before removing.

STATEMENT OF PRACTICAL TREATMENT

If In Eyes - Flush eyes with plenty of water. Call a Physician.
If Swallowed-Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.
If On Skin - Remove contaminated clothing. Wash affected areas with soap and water.
If Inhaled - Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this material to conform to the chemical description on the label and is suited for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the user or buyer.

Voluntary Purchasing Groups, Inc.

Bonham, Texas 75418

EPA Est. No. 7401-TX-1

EPA Reg. No. 7401-232

NET CONTENTS _____ Fl. Ozs.

