



PM23 7401-234 ~~7401-234~~ 1015

7401-234

Makers of fine lawn & garden and farm chemicals / Natural Guard Brand / Farmers Cooperative Brand / ferti-lome Brand / AMERICAN Brand / Hi-Yield Brand

Feb. 22, 1993

Joanne I. Miller, Product Manager 23  
U.S. EPA  
OPP  
Registration Division (H7505C)  
401 M Street SW  
Washington, D.C. 20460

I, Being an authorized representative of Voluntary Purchasing Groups, INC. certify that all containers of ferti-lome Systemic Weed & Feed 11-3-6, produced by June 15, 1994 will bear revised labeling in accordance with the revised labeling required for Task Force Technical and Manufacturing-use products. I further certify that all containers of said products sold or distributed by this company by Jan. 1, 1995 will bear revised labeling in accordance with the revised labeling for Task Force Technical and Manufacturing-use products.

Sincerely,

Raymond H. Smith  
Product Manager

Voluntary Purchasing Groups, Inc. / Bonham, Texas 75418 / (903) 583-5501

2015

(Front Panel)

Controls Weeds.....and Feeds your Lawns  
Coverage  
3,000 Sq.Ft.

- \* Systemic Action
- \* Attacks Through Leaves
- \* Attacks Through Roots

ferti-lome(R)  
Systemic  
Weed & Feed

\*Kills Most Broadleaf and Vining Type Weeds in St.  
Augustine, Common Bermuda, Zoysia, Rye, Bluegrass,  
Bents, Centipede, Bahia, and other Lawn Grasses.

Active Ingredients:

- \* Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid.....0.047%
- \*\* Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid.....0.165%
- \*\*\* Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)..... .020%
- Inert Ingredients:.....99.768%

- \*0.78 lb. or 0.039% 2,4-Dichlorophenoxyacetic acid per ton or 0.226 lb. per acre.
- \*\*2.72 lbs. or 0.136% 2-(2 methyl-4-chlorophenoxy) propionic acid per ton or 0.789 lb. per acre.
- \*\*\*0.32 lb. or 0.016% 3,6-dichloro-o-anisic acid per ton or 0.093 lb. per acre.
- \*\* Isomer Specific by AOAC Method 6.D01-5

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
See Back Panel for Additional Precautionary  
Statements

3075

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** May be harmful or fatal if swallowed. Avoid contact with skin, eyes and clothing. When using this product, wear long-sleeved shirt, long pants, socks, shoes and chemical resistant gloves. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled.

**STATEMENT OF PRACTICAL TREATMENT**

- IF SWALLOWED-** Call a physician or Poison Control Center immediately. if possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce Vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF INHALED** - Remove victim to fresh air. Apply artificial respiration if indicated.
- IF ON SKIN** - Remove contaminated clothing and wash affected areas with soap and water.
- IF IN EYES** - Flush eyes with plenty of water. Call a physician immediately.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drifts or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters. Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

**VOLUNTARY PURCHASING GROUPS, INC.**

**BONHAM, TEXAS 75418**

NET WEIGHT \_\_\_\_\_ LBS.

**BEST AVAILABLE COPY**

4075

(Back Panel)

ferti-lome(R)  
Systemic  
Weed & Feed  
11-5-6

**BEST AVAILABLE COPY**

**GUARANTEED ANALYSIS**

Total Nitrogen (N)..... 11%  
11.00% Ammoniacal Nitrogen  
Available Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>)..... 5%  
Soluble Potash (K<sub>2</sub>O)..... 6%  
Derived from-Primary Plant Foods: Sulphate  
of Ammonia, Ammoniated Superphosphate, Muriate  
of Potash.

Voluntary Purchasing Groups, Inc.  
Bonham, Texas 75418

~~PRECAUTIONARY STATEMENTS~~

~~HAZARDS TO HUMANS & DOMESTIC ANIMALS~~

~~CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes, or clothing. Wash thoroughly after handling. Keep children and pets off treated areas until this material is washed into the soil.~~

~~STATEMENT OF PRACTICAL TREATMENT~~

~~IF ON SKIN - Remove contaminated clothing and wash affected areas with soap and water. IF IN EYES - Flush eyes with plenty of water. Get medical attention immediately.~~

~~ENVIRONMENTAL HAZARDS~~

~~Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.~~

EPA Est. No. 7401-TX-1

EPA Reg. No. 7401-234

NET WT. 40 LBS.

Controls Such Weeds As:

Dandelion	Oxalis	Black Medic	Purslane
Chickweed	Wild Carrot	Burdock	Ragweed
Buckhorn	Wild Onion	Chickory	Sheep Sorrel
Plantains	Wild Garlic	Dock	Shepherd's Purse
Clovers	Wild Lettuce	Heal-All	Speedwell
Poison Ivy	Morning Glory	Knotweed	Thistle
Poison Oak	Buttercup	Lambsquarter	Varrow
Henbit	Pigweed	Lespedeza	English Daisy
Spurges	Bedstraw	Mallow	Milkweed
Pennywort	Ground Ivy	Peppergrass	Puncturevine

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**1. WHAT IS THIS PACKAGE FOR?**

ferti-lome Systemic Weed & Feed kills most broadleaf and vining type weeds found in lawn grasses, and at the same time supplies plant food for vigorous growth.

**2. HOW DO YOU USE IT?**

Use Fertilizer spreader setting from chart below. Apply anytime broadleaf weeds are present, except avoiding St. Augustine grass in summer months (June through September) when it is growing rapidly, because temporary foliage discoloration may occur. Best results are obtained when weeds are young and growing vigorously.

5075

(Back Panel Cont'd)

If weeds are growing slowly, control may be slow and more than one application may be needed. Do not water for 24 hours after application.

ONE BAG COVERS

3,000 Sq. Ft.

SPREADER

SPREADER SETTING\*

ferti-lome Cyclone

4½

ferti-lome Deluxe

8

Central (Sears, Wards, J.C. Penny)

8

Scotts

6½

To build good turf feed with ferti-lome monthly during growing season.

\* These settings can be used only as a guide; they are not absolute. The speed at which the applicator walks, the condition of the spreader, and the texture of the turf will alter the rate of fertilizer application.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

The use of this product aids in controlling weeds and obtaining deep green lawns.

NOTE: Avoid application to garden plants, flowers, shrubs and trees. Do not use on Centipede, Bent Grasses, Lippia or similar ground cover. Do not use on Dichondra.

Sincerely,  
the ferti-lome people

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.