

HOW TO USE... DIRECTIONS... TIMES WHEN...

2. HOW TO USE:
SHAKE WELL BEFORE USE OR SAMPLING. Mix in 100 gallons of water...
 Repeat in 10-14 days if necessary... NOTE: 1 lb. = 16 teaspoons, 1/2 lb. = 8 teaspoons.

The following application schedule for the primary pests (codling moth, apple maggot, oriental fruit moth and plum curculio) must be followed. Apply at petal fall and every 10-14 days thereafter until control is achieved.

FRUIT	PEST	CONCENTRATION	APPLICATION	REMARKS
APPLE	Apple Maggot	1/2 lb. per 100 gal.	Apply at petal fall and every 10-14 days thereafter.	Apply to all fruit trees.
	Codling Moth	1/2 lb. per 100 gal.	Apply at petal fall and every 10-14 days thereafter.	Apply to all fruit trees.
PLUM	Plum Curculio	1/2 lb. per 100 gal.	Apply at petal fall and every 10-14 days thereafter.	Apply to all fruit trees.
	Other pests	1/2 lb. per 100 gal.	Apply at petal fall and every 10-14 days thereafter.	Apply to all fruit trees.

FRUIT	PEST	CONCENTRATION	APPLICATION	REMARKS
APPLE	Apple Maggot, Aphid, Japanese Beetle, Leaf Hopper, European Red and Two Spotted Mites	5 lbs. per gal. or 1 qt. in 40 gal.	Apply at petal fall and every 10-14 days thereafter.	Apply to all fruit trees.
PLUM	Plum Curculio	1/2 lb. per 100 gal.	Apply at petal fall and every 10-14 days thereafter.	Apply to all fruit trees.
APPLE	Brown Rat, Blossom Borer and Berries, Rot	1/2 lb. per 100 gal.	Apply at petal fall and every 10-14 days thereafter.	Apply to all fruit trees.
APPLE	Codling Moth, Aphid, Cherry Fruit Worm, Cherry Fruit Fly, Japanese Beetle and Leafhopper	1/2 lb. per 100 gal.	Apply at petal fall and every 10-14 days thereafter.	Apply to all fruit trees.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

Outstanding, easy and economical insect and disease control may be had on your valuable fruit trees with use of ferti-lome Improved FRUIT TREE SPRAY.

CAUTION: May be harmful if swallowed, inhaled or absorbed through the skin. Do not take internally. Do not breathe fumes or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.


Food utensils such as a teaspoon or measuring cup should not be used for food purpose after use with insecticides.

Do not reuse container for any purpose. Destroy empty container.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



ferti-lome



ACCEPTED

VOLUNTARY PURCHASING GROUPS, INC. BONHAM, TEXAS