

(FRONT PANEL)

7401-237

PM 12 1/3

Fruit Tree

Fruit Tree


Spray

BEST AVAILABLE COPY

Contains insecticide-miticicide-scalicide-fungicide for insect and disease control on apple, peach, plum, and cherry fruit trees.

Active Ingredients:

- *Methoxychlor, Technical... 9.60%
- **Malathion... 8.07%
- ***Captan... 6.26%
- Related Derivatives... 0.14%
- ****INERT INGREDIENTS... 75.93%
- ****(Contains 48.95% Xylene Range Aromatic Solvent)
- *Equivalent to 2.45% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 2.15% other isomers and related products
- **0,0-dimethyl dithiophosphate of diethyl mercaptan
- ***N-Trichloroethylthio-4-cyclohexene-1,2-dicarboximide



Keep out of Reach of Children
DANGER

See back panel for additional precautionary statements

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

(Back Panel)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. WHAT IS THIS PACKAGE FOR?

Fast, sure insect and disease control is provided for your fruit trees when using ferti-lome Improved Fruit Tree Spray.

SHAKE WELL BEFORE USE OR SAMPLING: Mix required dosage in water as listed in table below. Apply (using any convenient sprayer) as a thorough cover spray to all tree surfaces and fruit. Repeat in 10-14 days if necessary to maintain control. Use only as specified on this label. NOTE: This product is not to be used on citrus trees.

For controlling apple diseases require for the Oriental Fruit Moth, Apply Maggot, Oriental Fruit Moth and Plum Curculio must be followed: Apply at petal fall and every 10-14 days thereafter until control is achieved.

CROP	INSECTS	DISEASES	RATES	REMARKS
APPLE	Red Banded Leaf Roller; Codling Moth; Plum Curculio.	Bitter, Black White, Botrytis and Bullseye Rots, Leaf spots, Brooks Fruit Spot and Flyspeck,	4½ tbs. per gal. or 1 qt. in 14 gals.	May injure McIntyre, Cortland and Red Delicious Apples. Do not apply within 7 days of harvest. On apples for Bitter, Black, White Rots and Leaf Spot apply in late cover and preharvest periods. For Botrytis and Bullseye Rots apply in first and second cover sprays and 1 or 2 preharvest sprays or as needed at 5-7 day intervals from preblossoms through early cover periods. For Brooks Fruit Spot and Flyspeck apply in cover periods.
	Tent Caterpillar and bud Moth (Apple only) and Mealy bug, Japanese Beetle, leaf hopper, European Pear Sawfly		2½ tbs. per gal. or 1 qt. in 14 gals.	
	Oriental Fruit Moth, Pear and Peach Scale		½ tbs. per gal. or 1 qt. in 14 gals.	Do not apply to peaches within 21 days of harvest. On peaches for Brown Rot, Rhizopus and Seal apply in pre-bloom, start shelter and cover periods. On plums for Brown Rot apply in full bloom and petal fall periods.

BEST AVAILABLE COPY

353

USE	INSECTS	DISEASES	RATES	REMARKS
PEACH	Codling Moth (peaches only) Aphid, Japanese Beetle, Leaf- hopper, European Red and Two		2 1/2 tbs. per gal. or 1 qt. in 28 gals.	BEST AVAILABLE COPY
	Cherry Curculio Aphids, Cherry Fruit Worm, Cherry Fruit Fly, Japanese Beetle and Leafhopper	Brown Rot Blight and Botrytis Rot	1/2 tbs. per gal. or 1 qt. in 28 gals. 1/2 tbs. per gal. or 1 qt. in 28 gals. 1/2 tbs. per gal. or 1 qt. in 28 gals. 1/2 tbs. per gal. or 1 qt. in 28 gals.	

3. HERE ARE THE RESULTS YOU MAY EXPECT!

Outstanding, easy and economical insect and disease control may be had on your valuable fruit trees with use of ferti-lome Improved Fruit Tree Spray.

STORAGE: Store in original container and place in a locked storage area.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Discard container in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse. Avoid contact with skin or clothing. Wear chemical resistant gloves when using indoors or outdoors. When using wear long pants and long-sleeved shirt and apply with the wind to your back. Wash non-disposable gloves thoroughly with soap and water before removing.

STATEMENT OF PRACTICAL TREATMENT

- If In Eyes - Flush eyes with plenty of water. Call a physician.
- If Swallowed - Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid vomiting.
- If On Skin - Remove contaminated clothing. Wash affected areas with soap and water.
- If Inhaled - Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this material to conform to the chemical description on the label and is suited for the purposes stated on the label. The limit of our liability, including all claims, shall be the pre-purchase price paid by the user or buyer.

Voluntary Purchasing Groups, Inc.
EPA Est. No. 7401-TX-1

Wortham, Texas 75418
EPA Reg. No. 7401-237

NET CONTENTS FL. OZS.