


**\$495**

**Coverage  
3,000 Sq. ft.**

**SYSTEMIC ACTION**



*Nov 8, 1973*

*7401234*



**PLANTAIN**



**HENBIT**



**DANDELION**



**CLOVER**



**CHICKWEED**

**ferti·lome**

**ferti·lome**

**ferti·lome**

KILLS MOST BROADLEAF AND VINING TYPE WEEDS IN ST. AUGUSTINE, COMMON BERMUDA,  
 CENTIPEDE, BAHIA AND OTHER LAWN GRASSES.  
**WEED & FEED** Systemic  
 KEEP OUT OF REACH OF CHILDREN

# Systemic WEED & FEED

\* KILLS MOST BROADLEAF AND VINING TYPE WEEDS IN ST. AUGUSTINE,  
 COMMON BERMUDA, ZOYSIA, RYE, BLUEGRASSES, BENTS, CENTIPEDE,  
 BAHIA AND OTHER LAWN GRASSES

**Active Ingredients:**

*Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid	.050%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	.165%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	.019%
Inert Ingredients:	99.766%

Used at 40 lbs. on 3,000 sq. ft. equivalent to:

- \*2,4-D Acid - 0.042% - 0.24 lbs./acre or 0.84 lbs. per ton
- \*\*MCPP Acid - 0.137% - 0.79 lbs./acre or 2.75 lbs. per ton
- \*\*\*Dicamba Acid - 0.016% - 0.09 lbs./acre or 0.33 lbs. per ton

**CAUTION**

KEEP OUT OF REACH OF CHILDREN

Keep away from food and food products. Avoid inhalation or contact with eyes, nose, mouth, or clothing. Wash after use. Do not use on dishandra or lippia ground covers. Do not re-use empty bag. Destroy empty bag. Do not burn. Do not apply in areas overlaid by roots of desirable trees and shrubs.

Avoid application on garden plants, flowers and shrubs. Avoid drift to desirable plants. Do not use on newly seeded lawns. Do not contaminate irrigation ditches or water used for irrigation.

Store in a dry place, away from plants, seeds, fertilizers, insecticides or fungicides.

A "WEEDER-FEEDER" FOR LAWN GRASSES  
 NET WT. 40 LBS.

KILLS MOST BROADLEAF AND VINING TYPE WEEDS IN ST. AUGUSTINE, COMMON BERMUDA,  
 CENTIPEDE, BAHIA AND OTHER LAWN GRASSES.  
**WEED & FEED** Systemic  
 KEEP OUT OF REACH OF CHILDREN