


7401-257

4-14-75

1 of 3

Best Available copy (Front Panel)



ferti-lome

MALATHION SPRAY

Protects Roses, Flowers, Ornamentals, Fruit Trees and Vegetables from insect damage

2 teaspoons make 1 gallon of spray

Active Ingredients:

Malathion (0,0-diethyl dithiophosphate of diethyl mercaptosuccinate).....	50%
Aromatic Petroleum Derivatives Solvent.....	38%
Inert Ingredients.....	12%
Total.....	100%

NET CONTENTS 16 FL. OZS. (ONE PINT)

EPA Est. 7401-TX-1

CAUTION (18 point)
KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS (12 Point)

(Back Panel)


1. WHAT IS THIS PACKAGE FOR?

Contains one of the home gardener's oldest "insect tools" that gives quick, effective control of listed insects.

2. HOW DO YOU USE IT?

Mix 2 teaspoons in 1 gallon of water (or 1 pint in 48 gallons of water for larger jobs). Apply thoroughly to all plant surfaces, to fruit and to vegetables. Repeat in 10-14 days if necessary. Do not apply to fruit or vegetables within 7 days of harvest.

ORNAMENTALS. (Roses, Azalea, Camellia, Gardenia, Carnations, Mums, Evergreens) Controls Aphids, Lace Bugs, Spider Mites, Whiteflies, Adult Japanese Beetle, Birch leaf miners, Bagworms, Tent caterpillar, Scale. For Scale apply when crawlers have settled in foliage. Do not apply to ferns.



(Back Panel)

FRUIT TREES:

CROP	INSECTS
APPLES	Wooly apple, Green apple and Rosey apple aphids; Codling and Bud moth; Plum Curculio; Tent Caterpillar; Mealey bugs; Forbs Scale.
PEACHES	Plum Curculio; Green Peach, Black Peach, Black Cherry aphids and Rusty plum aphids; Japanese beetle; Terrapin and Cottony peach scale; Oriental fruit moth.
PEARS	Codling moth; Plum curculio; Mealy bugs; Mites; Pear psylla; Fruit Tree and red banded leaf roller.
APRICOTS	Aphids; Codling moth; Orange tortrix; Terrapin and Soft brown scale.
CHERRIES	Black cherry aphid; Bud moth; Fruit tree leaf leaf roller; Cherry fruit fly.

NOTE: For the orchard pests Codling moth, Oriental fruit moth, and Plum curculio, apply at petal fall and every 10 to 14 days thereafter until control is achieved.

VEGETABLES:

CROP	INSECTS
BEANS	Aphids; Cucumber, Japanese and Mexican bean beetles; Leafhoppers; Spider mites.
CABBAGE	Aphids.
CUCUMBER, SQUASH	Aphids; Spider mites; Pickleworm.
CELERY	Aphids; Spider mites.
MELONS	Aphids; Spider mites.

3. HERE ARE THE RESULTS YOU MAY EXPECT!


This effective spray can be a great help to you in solving listed insect control problems.

CAUTION: May be harmful if swallowed, inhaled or absorbed through the skin. Do not take internally. Do not breathe fumes or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

ENVIRONMENTAL CAUTION: This product is toxic to fish. Keep out of lakes streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agriculture Extension Service.



3018

Page 5.

(Back Panel)

ENVIRONMENTAL CAUTION (continued) Food utensils such as a teaspoon or measuring cup should not be used for food purposes after use with insecticides.

Do not reuse container for any purpose. Destroy empty container. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

VOLUNTARY PURCHASING GROUPS, INC.

BONHAM, TEXAS

EPA Reg. No. 7401-257

