

### 1. WHAT IS THIS PACKAGE FOR?

Contains one of the home gardener's oldest insect tools that gives quick, effective control of listed insects.

### 2. HOW DO YOU USE IT?

Mix 2 teaspoons in 1 gallon of water for 1 pint in 48 gallons of water for larger jobs. Apply thoroughly to all plant surfaces to fruit and to vegetables. Repeat in 10-14 days if necessary. Do not apply to fruit or vegetables within 7 days of harvest.

**ORNAMENTALS:** Roses, Azalea, Camellia, Gardenia, Carnations, Marigolds, Evergreen. Controls: Aphids, Lace Bugs, Spider Mites, Whiteflies, Adult Japanese Beetle, Birch leaf miners, Bag worms, Tent caterpillar, Scale. For Scale apply when crawlers have settled on foliage. Do not apply to ferns.

### FRUIT TREES

FRUIT	INSECTS
APPLE	Woolly apple, Green apple and Rosey apple aphids, Codling and Bud moth, Plum Curculio, Tent Caterpillar, Mealy Bugs, Fruit Scale.
PEACH	Plum Curculio, Green Peach, Black Mealy Bug, Cherry aphids and Rusty plum aphids, Japanese beetle, Tent caterpillar, Peach scale, Oriental fruit moth.
PLUM	Trioxys moth, Plum curculio, Mealy Bugs, Mites, Pear psylla, Fruit Tree and red-banded leaf roller.
ORANGE	Aphids, Codling moth, Orange tortrix, Pear psylla and Soft brown scale.
CHERRY	Black cherry aphid, Bud moth, Fruit Tree and cherry fruit fly.

For control of insects on trees, spray with Oriental fruit moth, cherry fruit fly, and plum curculio at petal fall and every 10 to 15 days thereafter until control is achieved.

### VEGETABLES

CROP	INSECTS
BEANS	Aphids, Spider Mite, Japanese and Mexican Bean beetles, Leafhoppers, Spider Mites.
CABBAGE	Aphids.
CUCUMBER, Squash	Aphids, Spider Mites, Flea beetles.
CELERY	Aphids, Spider Mites.
MELONS	Aphids, Spider Mites.

### 3. HERE ARE THE RESULTS YOU MAY EXPECT!

This effective spray can be used on just the types of insects listed on the control problems.

**CAUTION:** May be harmful if swallowed or inhaled or absorbed through the skin. Do not take internally. Do not breathe fumes or spray mist. Avoid contact with skin or clothing. Wash thoroughly after handling. Avoid contact with food and livestock.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

**ENVIRONMENTAL CAUTION:** This product is toxic to fish. Keep out of streams, rivers, ponds, lakes, and flowing water. Do not allow runoff to reach streams, rivers, ponds, lakes, and flowing water. Do not allow runoff to reach streams, rivers, ponds, lakes, and flowing water. Do not allow runoff to reach streams, rivers, ponds, lakes, and flowing water.

This product is highly toxic to birds and should be kept out of their reach. Do not use in areas where birds may be disturbed. Keep your hands clean. Avoid contact with eyes.

Food utensils and dishes should be washed thoroughly before use. Do not use this product on dishes or dishes.

Do not use this product on any fish. Do not use this product on any fish. Do not use this product on any fish. Do not use this product on any fish.

VOLUNTARY PURCHASING GROUP, INC.  
BOHNAM, TEXAS

2-10-69 (1) EPA Reg. No. 7401-217 AA



# ferti·lome



# MALATHION SPRAY

5/2/69

- PROTECTS ROSES, FLOWERS, ORNAMENTALS, FRUIT TREES AND VEGETABLES FROM INSECT DAMAGE
- 2 TEASPOONS MAKE 1 GALLON OF SPRAY

#### Active Ingredients

Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate)  
Aromatic Petroleum Derivatives Solvent

50%  
38%  
12%

#### Inert Ingredients:

Total


100%

NET CONTENTS 16 FL OZS (ONE PINT)

EPA Est. 7401-TX-1

**CAUTION KEEP OUT OF REACH OF CHILDREN  
SEE BACK PANEL FOR ADDITIONAL CAUTIONS**

BEST AVAILABLE COPY



CROP	INSECTS
APPLES	Woolly apple, Green apple and Rosey apple aphids, Codling and Bud moth, Plum Curculio; Tent Caterpillar; Mealy bugs; Forbs Scale
PEACHES	Plum Curculio, Green Peach, Black Peach, Black Cherry aphids and Rusty plum aphids, Japanese beetle, Terrapin and Cottony peach scale. Oriental fruit moth
PEARS	Codling moth, Plum curculio, Mealy bugs, Mites, Pear psylla, Fruit Tree and red handed leaf roller
APRICOTS	Aphids, Codling moth, Orange tortrix, Terrapin and Soft brown scale
CHERRY	Black cherry aphid, Bud moth, Fruit tree leaf roller, Cherry fruit fly

NOTE: For the orchard pests Codling moth, Oriental fruit moth, and Plum curculio, apply at petal fall and every 10 to 14 days thereafter until control is achieved.

thoroughly after foodstuffs

**DO NOT USE, PO ENVIRONMENT,**

out of lakes, st likely to occur. Drift from areas of equipment, a specified on lat

This product is on residues on from your Coop Food utensils su ne used for foo

Do not reuse co

Buyer assumes r teral ent in str

**VOLL**

1-12-71

