
Hi-Yiek

Decimate Concentrate **Plus Surfactant**



-

the soil. once.

CONTAINER DISPOSAL: Do not reuse container. Wash thoroughly with water and detergent, crush if possible and discard in a safe blace. NON-WARRANTY: Follow directions carefully Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this muterial not in

strict accordance with directions given herewith. GENERAL INFORMATION: Hi-Yield DECIMATE CONCENTRATE PLUS SURFACTANT is an effective herbicide particularly for grassy weeds when applied preplant and post-emergent to cotton, Learing citrus, listed agricultural plantings, non-crop areas and lawn and ornamental turf. It is unnecessary to add any other surfactant to the spray. solution. Best results are obtained on young actively growing weeds (temperature above 70° F.). The phytotoxic properties are quickly inactivated on contact with the soil.

MIXING INSTRUCTIONS: Hi-Yield DECIMATE CONCENTRATE PLUS SURFACTANT is completely water soluble. Fill the spray tank about half full with water and add required amount of herbicide. Finish filling tank with water using mild agitation. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period.

COTTON: Preplant Application: For control of weeds and grasses such as johnsongrass, nutsedge, dallisgrass, cocklehur, ragweed, sandbur sandspur), crabgrass and puncture vine. Mix 6-1-3 pints Hi-Yield

READ ENTIRE LABEL BEFORE USING THIS PRODUCT

CAUTION: Keep out of reach of children. Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into

ANTIDOTE: If taken internally, induce vomiting and call physician at

Do not feed treated forage to livestock or graze treated areas. Do not contaminate waters used for domestic consumption, or by animals, wildlife and aquatic life, or for irrigation purposes

DECIMATE CONCENTRATE PLUS SURF Do not add any surfactant to sprc spray solution per acre. Apply once weeds have emerged. Cotton may be

COTTON: Post Emergence Directed ! weeds and grasses such as johnsong bur, ragweed, sandbur (sandspur), cr 1 3 pints in 100 gallons of water. D solution. Apply spray solution at 40 c tion in 40 inch rows, apply 1 gallon width per acre. Direct the spray so plant. Nozzles should be placed so foliage. Where practical, withhold (10) day. A second application may son to control perennials such as je the second application 1 to 3 weel Slight burning and a reddish disco occasionally occur following the recotton plant will develop normally.

Apply only when cotton is 3 inches apply after first bloom. Do not fee graze treated areas.

SEARING CITRUS (Except Florida):

HI-YIELD DECEMATE CONCENTRATE PL rected spray application in citrus orch tangerine, lemon and lime orchards. of 3 10 to 6 10 gallons per acre. A TRATE PLUS SURFACTANT at the rate water. Apply as a directed spray in trees. Spray unwanted vegetation to occurs, repeat as required; however, year. Do not allow spray solution (bark. Use a shield if necessary for i AGRICULTURAL PLANTINGS: HI-Yield SURFACTANT is useful as a directed uous fruit, nut and citrus orchards si

READ ENTIRE LABEL BEFORE USING THIS PRODUCT

_ . .

CAUTION: Keep out of reach of children. Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

Do not feed treated forage to livestock or graze treated areas. Do not contaminate waters used for domestic consumption, or by avimals, wildlife and aquatic life, or for irrigation purposes

CONTAINER DISPOSAL: Do not reuse container. Wash thoroughly with water and detergent, crush if possible and discard in a safe place.

NON-WARRANTY: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this muterial not in strict accordance with directions given herewith.

GENERAL INFORMATION: Hi-Yield DECIMATE CONCENTRATE PLUS SURFACTANT is an effective herbicide particularly for grassy weeds when applied preplant and post-emergent to cotton, kearing citrus, listed agricultural plantings, non-crop areas and lawn and ornamental turf. It is unnecessary to add any other surfactant to the spray solution. Best results are obtained on young actively growing weeds (temperature above 70° F.). The phytotoxic properties are quickly inactivated on contact with the soil.

MIXING INSTRUCTIONS: HI-Yield DECIMATE CONCENTRATE PLUS SURFACTANT is completely water soluble. Fill the spray tank about half full with water and add required amount of herbicide. Finish filling tank with water using mild agitation. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period.

COTTON: Preplant Application: For control of weeds and grasses such as johnsongrass, nutsedge, dallisgrass, cocklebur, ragweed, sandbur (sandspur), crabgrass and puncture vine. Mix 6-1-3 pints Hi-Yield DECIMATE CONCENTRATE PLUS SURFACTANT in 100 gallons of water. Do not add any surfactant to spray solution. Apply 40 gallons of spray solution per acre. Apply once if cotton planting is delayed and weeds have emerged. Cotton may be planted immediately.

COTTON: Post Emergence Directed Spray Application: For Control of weeds and grasses such as johnsongrass, nutsedge, dallisgrass, cocklebur, raqweed, sandbur (sandspur), crabarass and puncture vine. Mix 6 1/3 pints in 100 gallons of water. Do not add any surfactant to spray solution. Apply spray solution at 40 gallons per acre. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per acre. Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the cotton foliage. Where practical, withhold irrigation for approximately ten (10) day. A second application may be required during growing season to control perennials such as johnsongrass and nutsedge. Apply the second application 1 to 3 weeks later.

Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment, but the cotton plant will develop normally.

Apply only when cotton is 3 inches in height to first bloom. Do not apply after first bloom. Do not feed treated foliage to livestock or graze treated areas.

SEARING CITRUS (Except Florida):

Hi-Yield DECIMATE CONCENTRATE PLUS SURFACTANT is useful as a directed spray application in citrus orchards, such as oranges, grapefruit tangerine, lemon and lime orchards. It should be applied at the rate of 3, 10 to 6 '10 gallons per acre. Mix Hi-Yield DECIMATE CONCEN-TRATE PLUS SURFACTANT at the rate of 3 10 gallon in 75 gallons of water. Apply as a directed spray in interspaces and around base of trees. Spray unwanted vegetation to just short of run-off. If regrowth occurs, repeat as required; however, do not exceed 3 applications per year. Do not allow spray solution to contact fruit, leaves, stems or bark. Use a shield if necessary for nursery plantings or young trees. AGRICULTURAL PLANTINGS: Hi-Yield DECIMATE CONCENTRATE PLUS SURFACTANT is useful as a directed application in non-bearing deciduous fruit, nut and citrus orchards such is apple, peach, pear, plum,

prune, apricot, cherry; almond, walnut, orange, grapefruit, tangerine, lemon and lime orchards. It should be applied at the rate of 3/10 to 6/10 adllons per acre.

Mix Hi-Yield DECIMATE CONCENTRATE PLUS SURFACTANT at the rate of 3, 10 gallon in 50 to 100 gallons of water. Apply as a directed spray in interspaces and around base of trees or vines. Spray unwanted vegetation to just short of run-off. If regrowth occurs repeat as required; however, do not exceed 3 applications per year.

Do not allow spray solution to contact leaves, stems or bark. Use a shield, if necessary, for nursery plantings or young trees. Do not apply around trees or vines from which fruit will be harvested within one year of treatment.

NON-CROP: HI-Yield DECIMATE CONCENTRATE PLUS SURFACTANT s useful for control of johnsongrass, nutsedge, dallisgrass, cocklebur, barnyard grass, crabgrass, sandbur (sandspur), ragweed, puncture vine and certain other weeds on drainage ditchbanks, rights-of-way, fence rows, storage yards and similar non-crop areas.

Mix Hi-Yield DECIMATE CONCENTRATE PLUS SURFACTANT at a rate of 1 3 to 2 3 gallon in 100 gallons of wate: Spray unwanted vegetation sufficiently to provide complete coverage. Use spray equipment that gives good low volume coverage. If regrowth occurs, repeat as required. LAWN AND ORNAMENTAL TURF: HI-Yield DECIMATE CONCENTRATE

PLUS SURFACTANT is useful for control of dallisgrass, sandbur (sandspur), bahiagraiss, nutsedge, crabgrass, chickweed and wood sorrel with little or no injury to tolerant lawn grasses. On new lawns, do not treat until after three mowings. Tolerant grasses may be temporarily discolored. Zoysiagrass, bluegrass and bermudagrass are quite tolerant. Do not use on St. Augustine or centipede. Mow lawns 1 to 1-1/2 inches high before treatment. Mix 9/10 fl. oz. (5.4.10 teaspoons) of Hoyield DECIMATE CONCENTRATE PLUS SURFACTANT in 2-1/2 gallons of water and apply to an area of 1000 sq. ft. Spray thoroughly to wet all undesirable plants. Repeat applications, 10 to 14 days apart, may be needed for good control.

VOLUNTARY PURCHASING GROUPS, INC.

Bonham, Texas

14M 1 45


EPA Reg. No. 7401-260 AA.

RIO ANN WEED LER DILID

٠.

,

ļ



College of Relation Charter and Science of Elliphication Bracket of the An - Plant - Ke there to Base they (have do) Inste

SAFE FOR ESTABLISHED LAWNS when the mental as the server some grasses have in an south the assistant to keep start of a first base about this would be refered.

KEEP OUT OF REACH OF CHILDREN SEE BACK PANEL FOR ADDITIONAL CAUTIONS ACENE INGREDIENTS

计环日日的时间日日日



CAUTION

ACCEPTED 5.5 M 175 s and a F in the LAWN · marriet-s. WEED KILLER Contains 2,4-D and Silvex FORMULATED WITH VERMICULITE FOR A WEED FREE LAWN Here the late of the test of the test of the test

LAWN H10

A

. . .

A Milston Monan + + + may