

7401-267

03/22/1976

1/2


ACCEPTED
 MAR 22 1976
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registration under
 EPA Reg. No. **7401-267**

Hi-Yield[®]

ACCEPTED
 MAR 22 1976
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registration under
 EPA Reg. No.

5%
Malathion
Dust

DEC 13 1976
 RECEIVED
 FEDERAL BUREAU OF INVESTIGATION
 U.S. DEPARTMENT OF JUSTICE



2/2

5% Malathion Dust

DIRECTIONS: For Use Against Certain Listed Insects Infesting Range Cattle, Dairy Cattle, Horses, Swine, Goats, Sheep, Poultry and Poultry Houses.

PRECAUTIONS IN USING THIS PRODUCT: Do not treat animals under one month of age. When applying dust, avoid contamination of feed, food containers and watering troughs, and watering fountains. Do not apply to animals not listed on this label.

ANIMAL	PEST	DOSE	REMARKS
BEEF CATTLE HORSES	Lice, Ticks and Hornfly	4 table- spoons per head	Apply on back and neck of animals.
DAIRY CATTLE (nonlactating)	Hornfly	3 table- spoons per head	Apply on back and neck of animal. Do not apply during lactation or within 2 weeks of freshening.
DAIRY CATTLE (lactating)	Hornfly	3 table- spoons per head	Apply on back and neck of animal. Do not apply within 5 hours of milking. Repeat application at 10-14 day intervals if needed.
POULTRY (Chickens, Ducks, Geese, Turkey)	Northern fowl mite, Poultry lice, Chicken red mite		Thorough application directly to birds. NOTE: For dust box treatment for 100 birds. 12 lbs. per 6 cu. ft. of dust box. Remove box when empty.
GOATS, SHEEP	Lice, Ticks, Keds		Thorough application to animal by hand or power duster. Do not apply to lactating goats.
SWINE	Lice		
POULTRY HOUSES	Northern fowl mite, Chicken red mite, Poultry lice	5 lbs. per 100 sq. ft.	Apply dust to litter, floor, walks, nests, roosts, and adjacent areas, using rotary hand duster, puff duster or by sprinkler can.

5% Malathion Dust