

Hi-Yield

**Livestock
Spray**



276

PRECAUTIONS IN USING THIS PRODUCT: Do not treat calves under 1 month of age. Do not dip or spray animals within 28 days of slaughter. Do not dip or spray within 2 weeks of freshening. Do not dip or spray animals when weather is cold or stormy as injury may result from chilling. Do not dip or spray overheated, thirsty or sick animals. **DO NOT USE ON DAIRY ANIMALS IN DAIRY BARNs, MILK ROOMS OR POULTRY HOUSES. DO NOT USE ON LACTATING DAIRY ANIMALS OR GOATS. DO NOT SPRAY DAIRY ANIMALS UNDER 3 MONTHS OF AGE.**

AS A SPRAY:

CATTLE: Lice, Ticks, Horn Flies, Mosquitos, Sarcoptic Mange — Mix 1 cup per 5 gal. of water (or 2½ pints per 25 gal. of water). Spray animal thoroughly; do not overspray. A spray of about 2 quarts of dilute spray per adult animal affords protection for 2-3 weeks. Repeat as necessary. For lice one treatment will usually be sufficient for a season. **Do not use on dairy cattle.**

HOGS: Lice — Mix 1 cup per 5 gal. of water (or 2½ pints per 25 gal. of water). Spray animal thoroughly. One treatment is usually sufficient for a season.

SHEEP AND GOATS: Lice and Sheep Keds (ticks) — Mix 1 1/3 tablespoons per gal. of water (or 3/8 cup for 5 gal. of water). Spray animal thoroughly using high pressure in order to penetrate fleece. **Ticks:** Mix 1 3/5 tablespoons per gal of water (or 5 tablespoons per 5 gal. of water). Spray thoroughly as above. Repeat at 2-3 weeks or as necessary.

AS A DIP:

For Control of Listed Pests on Cattle: Use 1 gal. Hi-Yield LIVESTOCK SPRAY NO. 2 per 100 gals. water. Fill vat first with water, then add required amount of Hi-Yield LIVESTOCK SPRAY NO. 2. Mix thoroughly for 3 minutes or longer. Any time dipping operation is stopped for 30 minutes, agitate thoroughly before resuming dipping. Completely submerge animal's body once or twice. Duck head at least twice. **Do not use on calves or other young animals under three months of age. Do not use on dogs, cats or other household pets or poultry. Do not dip dairy cattle.**

If young animals between 3 and 6 months of age must be treated use spray only. Always water animals before dipping so that they will not drink the dipping solution. Upon prolonged standing or filth contamination in dipping vats, the active chemical ingredients used will sometimes separate. Do not use such separated dipping solution, since death to animals may occur. Clean vat and re-charge with fresh solution.

CABLE TYPE BACK RUBBERS for Range Cattle to Control Horn Flies — Use 1 pint of Hi-Yield LIVESTOCK SPRAY NO. 2 in 1 gal. No. 2 fuel oil or furnace oil. **DO NOT USE ON DAIRY ANIMALS.** Do not treat beef cattle within 30 days of slaughter. Replenish units every 3-5 weeks.

WARNING: Keep out of reach of children. May be fatal if swallowed. May be absorbed through the skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash hands thoroughly with soap and warm water after handling the concentrate or applying the diluted product to animals. Do not contaminate feed or foodstuffs. Do not use inside dairy barns, milk rooms or milk processing rooms. Do not use in poultry houses. Food utensils such as tablespoons or measuring cups should not be used for food purposes after using with pesticides.

DO NOT USE, POUR, OR SPILL NEAR HEAT OR OPEN FLAME

Environmental Caution: This product is toxic to fish and wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy by perforating or crushing. Bury or discard in a safe place away from water supplies.

Sevier's guarantee shall be limited to the terms of the label and subject thereto the buyer assumes any risks to persons or property arising out of use or handling and accepts the product on these conditions.

**VOLUNTARY PURCHASING GROUPS, INC.
BONHAM, TEXAS**

EPA Reg. No. 105-101-01 AA