PRECAUTIONARY STATEMENTS ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning or disposal of wastes.

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

1. WHAT IS THIS PACKAGE FOR?

ferti-lome Dipel® Dust is a biological insecticide intended for use against the listed insects.

2. HOW DO YOU USE IT?

Apply when pests are first noticed and repeat as necessary for control. Thorough plant coverage is necessary for control.

CELERY, LETTUCE, MELONS, POTATOES, SPINACH, TO-MATOES, CUCUMBERS: To control Cabbage Looper, apply 25 to 50 lbs. per acre (or 1/2-1 oz. per 50 ft. of row one foot wide).

BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, KALE, MUSTARD GREENS, TURNIP GREENS: To control Cabbage Looper and Imported Cabbage Worm, apply 15 to 25 lbs. per acre (or 1/3-1/2 oz. per 50 ft. of row one foot wide.)

TOBACCO: To control Hornworm apply 12 to 25 lbs, per acre (or 1/4-1/2 oz. per 50 ft. of row one foot wide).

3. HERE ARE THE RESULTS YOU MAY EXPECT!

Provides effective control of specified worms.

buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

28M-2-118



ferti·lome



Dipel Dust

Biological Insecticide

Under Pungit as agn

Active Ingredient:

Bacillus thuringiensis, Berliner, 320 International Units of Potency per mg. 0.15 billion International Units per pound of this product.

®Dipel is the registered trademark of Abbott Laboratories, North Chicago, Illinois.

EPA Reg. No. 7401-290

EPA Est. 7401-TX-1

KEEP OUT OF REACH OF CHILDREN CAUTION

See Side Panel For Additional Precautionary Statements

NET WEIGHT | LB. (454 grams)

VOLUNTARY PURCHASING GROUPS, INC.
BONHAM, TEXAS