

WHITE - UNVARNISH

**CAUTION**  
**STATEMENT OF PRACTICAL TREATMENT**

Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS**

**CAUTION:** May be harmful if swallowed. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Keep out of reach of children.

**TO PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning or equipment or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Sevin is an outstanding insecticide that controls a wide variety of insect pests. Sevin is not effective against mites; where mites are a problem use a suitable miticide. Hi-Yield Mater-Tater And Other Vegetable Dust is compatible with commonly used miticides, fungicides and insecticides. Do NOT use with lime or other alkaline material.

**FOR SMALL MEASUREMENTS**

- 57 standard level tablespoonfuls = 1 lb.
- 3 1/2 standard level tablespoonfuls = 1 oz.
- 4 1/2 ounces = 1 cupful

Apply Hi-Yield Mater-Tater And Other Vegetable Dust when insects first appear, preferably before damage from feeding occurs, or as indicated. For best results apply thoroughly to upper and lower leaf surfaces. If necessary, repeat applications every 7 to 10 days, or as indicated. Apply with conventional dusting equipment. Do NOT dust too heavily or crop injury may result.

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

**POTATOES, TOMATOES, EGGPLANTS AND PEPPERS:** For control of Colorado Potato Beetle, Flea Beetles and Leafhoppers: Apply 1 1/2 to 2 lbs. per 1,000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus and Stink Bugs: Apply 2 1/2 lbs. per 1,000 sq. ft.

**BEANS:** For control of Mexican Bean Beetle: Apply 2 lbs. per 1,000 sq. ft. For control of Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Flea Beetles and Japanese Beetle: Apply 1 1/2 lbs. per 1,000 sq. ft. For control to Corn Earworm, Armyworms and Velvet Bean Caterpillar: Apply 1 1/2 to 2 lbs. per 1,000 sq. ft. For control of Lima Bean Pod Borer, Lygus and Stink Bugs: Apply 2 1/2 lbs. per 1,000 sq. ft.

**CORN:** For control of Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles and Japanese Beetle: Apply 1 1/2 to 2 lbs. per 1,000 sq. ft. For control of Larvae in Whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking Silks and Ears: Apply at 2 to 3 day intervals, starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. May be applied up to harvest on Sweet Corn. Do NOT apply within 7 days of harvest on corn used for fodder or forage.

**CUCUMBERS AND SUMMER SQUASH:** For control of Pickleworm, Melonworm, Cucumber Beetles, Squash Bugs and Flea Beetles: Apply 1 1/2 lbs. per 1,000 sq. ft.

**LETTUCE:** For control of Armyworm, Cabbage Caterpillar, small Cabbage Looper, Corn Earworm, Flea Beetle, Harlequin Bug, Leafhopper, Six-spotted Leafhopper, Spittlebug, Lygus and Stink Bugs: Apply 2 lbs. per 1,000 sq. ft. Make first application when insects are first noted and repeat as needed at 5 to 7 day intervals.

**OKRA:** For control of Corn Earworm and Stink Bugs: Apply 2 lbs. per 1,000 sq. ft. Make first application when insects are first noted and repeat at 5 to 7 day intervals, or as necessary.

**BLUEBERRIES:** For control of Cherry and Cranberry Fruitworm: Apply 2 lbs. per 1,000 sq. ft.

**CRANBERRIES:** For control of Fruitworms, Fireworms, Leafhoppers and Cutworms: Apply 2 1/2 to 3 lbs. per 1,000 sq. ft.

**STRAWBERRIES:** For control of Meadow Spittlebug and Leaf Roller: Apply 2 1/2 lbs. per 1,000 sq. ft.

**ORNAMENTALS:** Arborvitae, Azalea, Boxwood, Dogwood, Rose, Spirea, Yew: For control of Japanese Beetle, Bagworm, Aphid, Flea Beetle, Leafhoppers, Tent Caterpillar and exposed Thrips: Apply 2 1/2 lbs. per 1,000 sq. ft. Make first application when insects are first noted and repeat at 7 to 10 day intervals, as necessary. Do NOT use on Boston Ivy or Virginia Creeper.

**SEVIN** may be applied up to and including the day of harvest on Beans, Blueberries, Sweet Corn, Cucumbers, Okra, Summer Squash, Tomatoes, Eggplant, Peppers, Potatoes and within 1 day of harvest on Cranberries and Strawberries and within 5 days of harvest on Head Lettuce, or within 14 days of Leaf Lettuce. Allow 7 days between last application and harvest on Corn for use as forage and fodder as feed for dairy and meat animals. Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable Sevin residues when Sevin is used in accordance with label directions. When Sevin has been applied in accordance with these directions for use, feeding corn forage will not result in residues in milk or meat.

**STORAGE:** Store in original container and place in a locked storage area.

**DISPOSAL:** Do not reuse container. Wrap and put in trash.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.


\*SEVIN, is the registered trademark of Union Carbide Corp. for carbaryl insecticide.

VOLUNTARY PURCHASING GROUPS, INC.

BONHAM, TEXAS 75418

# Hi-Yield<sup>®</sup>

## Mater-Tater And Other Vegetable Dust



### An Insecticide For Use On Certain Small Fruits, Ornamentals and Vegetables.

Active Ingredient:  
 \*Carbaryl (1-naphtha N-methylcarbamate) ..... 1.75%  
 Inert Ingredients: ..... 98.25%  
 EPA Est. 7401-TX-1  
 EPA Reg. No. 7401-291

NET WEIGHT 1 LB.  
**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
 See Back Panel For Additional Cautions

PH 04

7401-291

6/5/85

73121