

7401-291

4/16/2003

1/5

[FRONT PANEL]

ACCEPTED

APR 16 2003
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 7401-291

Hi-Yield®

MATER-TATER
And Other Vegetable Dust

- An Insecticide For Use On Listed Home Ornamentals and Fruit and Vegetable Gardens

ACTIVE INGREDIENT:

Carbaryl: 1-naphthyl N-methylcarbamate 1.75%

INERT INGREDIENTS: 99.25%

TOTAL: 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Cautions

NET WEIGHT _____

[BACK/SIDE PANEL]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Carbaryl is an insecticide that controls a wide variety of insect pests. Carbaryl is not effective against mites; where mites are a problem use a suitable miticide.

Hi-Yield® Mater-Tater And Other Vegetable Dust is compatible with commonly used miticides, fungicides and insecticides. Do NOT use with lime or other alkaline material.

FOR SMALL MEASUREMENTS

- 57 standard level tablespoonfuls = 1 lb.
- 3 ½ standard level tablespoonfuls = 1 oz.
- 4 ½ ounces = 1 cupful

Apply Hi-Yield® Mater-Tater And Other Vegetable Dust when insects first appear, preferably before damage from feeding occurs, or as indicated. For best results apply thoroughly to upper and lower leaf surfaces. If necessary, repeat applications every 7 to 10 days, or as indicated. Apply with conventional dusting equipment. Do NOT dust too heavily or crop injury may result.

Subsequent planting of feed and feed crops in treated soil must be made with crops on this or other carbaryl labels.

HOME GARDEN FRUITS AND VEGETABLES

Follow Pre-Harvest Intervals (PHI) listed in () following each individual crop and plant.

POTATOES (7), TOMATOES (3), EGGPLANTS (3) and PEPPERS (3): For control of Colorado Potato Beetle, Flea Beetles and Leafhoppers. Apply 1 ½ to 2 lbs. per 1,000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus and Stink Bugs: Apply 2 ½ lbs. per 1,000 sq. ft.

BEANS (3): For control of Mexican Bean Beetle: Apply 2 lbs. per 1,000 sq. ft. For control of Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Flea Beetles and Japanese Beetle: Apply 1 ½ lbs. per 1,000 sq. ft. For control of Corn Earworm, Armyworms and Velvet Bean Caterpillar: Apply 1 ½ to 2 lbs. per 1,000 sq. ft. For control of Lima Bean Pod Borer, Lygus and Stink Bugs: Apply 2 ½ lbs. per 1,000 sq. ft.

[BACK/SIDE PANEL CONTINUED]

CORN (2): For control of Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles and Japanese Beetle: Apply 1 ½ to 2 lbs. per 1,000 sq. ft. For control of Larvae in whorl (budworms damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking Silks and Ears: Apply at 2 to 3 day intervals, starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control.

CUCUMBERS (3) and SUMMER SQUASH (3): For control of Pickelworm, Melonworm, Cucumber Beetles, Squash Bugs and Flea Beetles. Apply 1 ½ lbs. per 1,000 sq. ft.

LETTUCE (14): For control of Armyworm, Cabbage Caterpillar, Small Cabbage Looper, Corn Earworm, Flea Beetle, harlequin Bug, Leafhopper, Six-spotted Leafhopper, Spittlebug, Lygus and Stink Bugs: Apply 2 lbs. per 1,000 sq. ft. Make first application when insects are first noted and repeat as needed at 5 to 7 day intervals.

OKRA (3): For control of Corn Earworm and Stink Bugs: Apply 2 lbs. per 1,000 sq. ft. Make first application when insects are first noted and repeat at 5 to 7 day intervals, or as necessary.

BLUEBERRIES (7): For control of Cherry and Cranberry Fruitworm: Apply 2 lbs. per 1,000 sq. ft.

CRANBERRIES (7): For control of Fruitworms, Fireworms, Leafhoppers and Cutworms: Apply 2 ½ to 3 lbs. per 1,000 sq. ft.

STRAWBERRIES (7): For control of Meadow Spittlebug and Leaf Roller: Apply 2 ½ lbs. per 1,000 sq. ft.

ORNAMENTALS: Arborvitae, Azalea, Boxwood, Dogwood, Rose, Spirea, Yew: For control of Japanese Beetle, Bagworm, Aphid, Flea Beetle, Leafhopper, Tent Caterpillar and exposed Thrips: Apply 2 ½ lbs. per 1,000 sq. ft. Make first application when insects are first noted and repeat at 7 to 10 day intervals, as necessary. Do NOT use on Boston Ivy or Virginia Creeper.

STORAGE AND DISPOSAL

STORAGE: Keep out of reach of children and pets. Store in original containers only. Store in cool, dry place and avoid excess heat.

DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

[BACK/SIDE PANEL CONTINUED]

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION: May be harmful if swallowed. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Keep out of reach of children.

USER SAFETY REQUIREMENTS	
<ul style="list-style-type: none"> * Wear long-sleeved shirt, long pants, shoes plus sock and household latex or rubber gloves when mixing and applying this product. * Wear a hat and eye protection when making overhead applications. * Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use. * Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet. * Do not allow children or pets to contact treated area until sprays have dried. 	

FIRST AID N-methyl Carbamate	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 or 1-800-900-4044 for emergency medical treatment information.

NOTE TO PHYSICIAN: Carbaryl is a N-methyl carbamate that inhibits cholinesterase. Atropine is an antidote. DO NOT use 2-PAM, opiates or cholinesterase-inhibiting drugs

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from area treated. Apply this product only as specified on the label. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

NOTE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

VOLUNTARY PURCHASING GROUPS, INC.
P.O. BOX 460
BONHAM, TX 75418

EPA REG. NO. 7401-291
EPA EST. NO. 7401-TX-1