

PM 04

7401-293

11/14/79

3

Best Available Copy

PS 181



Kill-a-bug Lawn Granules

Hi-Yield

Special Kill-a-bug Lawn Granules

Containing DURSBAN®



Kill-a-bug Lawn Granules

BEST DOCUMENT AVAILABLE

Hi-Yield

Special Kill-a-bug Lawn Granules

Containing DURSBAN®

This product is recommended to control Certain White Grubs¹, Chinch Bugs, and Wireworms (Leads Mistle), Anns, Armyworms, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Fleas, Bembidid (Billbug), and Hyperodes (Turf grub) Weevils Infesting Home Lawns and Other Ornamental Turf Areas.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent of its labeling.

Apply uniformly to home lawns and other ornamental and recreational turf grass areas using any hand or power-operated gravity or rotary type spreader designed for application of granular materials. Treat when insects first appear at the rate of 12½ pounds per 5000 sq. ft. To help obtain uniform distribution, apply one-half the required amount of product in one direction, then apply the balance at a right angle immediately after application. Thoroughly water the treated turf to wash the insecticide into the zone of insect habitation. Consult the following table to determine amount and water needed for another sized areas. Re-treat as necessary.

RATES TO USE

Area to be Treated	Pounds of Dursban-1G	Gallons of Water to Sprinkle on Turf after Treatment
1000 sq. ft.	2½	30
5000 sq. ft.	12½	150
10700 sq. ft.	25	325

SPREADER SETTING

Make	2 ½ lbs. Per 5000 Sq. Ft.	5 lbs. Per 10000 Sq. Ft.
Fort-Lane CYCLONE	3½	4½
Fort-Lane DELUXE	6	7
Scott	6½	7½
Scars	6	7

NOTE: Above spreader settings are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting if necessary to ensure correct rate of application.

¹To control certain white grubs (larvae of scarab beetles) and brownish beetle larvae that injure lawns, apply this insecticide at the rate of 12½ pounds per 5000 sq. ft. Apply the granules in the early morning, followed immediately by thorough watering. The insecticide will reach the turf. For best results treat when grubs are actively feeding in the soil surface, usually during late July and August. Consult your Agricultural Extension Service specialist for proper time to treat in your area.

²To control Hyperodes (billgrub) weevils use twice the rate shown in the table. Treat infested problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

STORAGE AND DISPOSAL

Storage: Keep away from food, feedstuffs, and domestic water supply. Keep container closed.

Disposal: Do not reuse empty container. Wrap container and put in trash collection.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

DURSBAN® trademark of The Dow Chemical Company, U.S. Patent 2,815,000.

WILLIAMS-SCHWAB & COMPANY, INC., CHASING GROUPE, INC., BOWMAN, TEXAS 75418

NET WT. 12 1/2 LBS.

ACCEPTED

NOV 14 1979

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for use including registration number 7401-293

032