

Fleas, Chiggers, Clovers Mites,)



Hi-Yield[®]

DURSBAN^{*}

Spray

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

LAWNS & TURF: To control Chinch Bugs, Sod Webworms (Lawn Moth Larvae), Ants, Brown Dog Ticks, Crickets, Cutworms, Earwigs & Grasshoppers infesting lawns and other turf areas, mix this product with water according to directions in table below and apply as a coarse droplet spray. Lawn should be moist at time of application. Spray when pests first appear and retreat as needed.

| DOSAGE & DILUTIONS FOR USE ON LAWNS & TURF | | |
|--|----------------------------------|---------------------------|
| Area to Be Treated | Amount of Hi-Yield DURSBAN SPRAY | Dilute with Water to Make |
| 100 sq. ft. | 7 teaspoons | 3 gallons |
| 500 sq. ft. | 12 tablespoons (6 fl. ozs.) | 15 gallons |
| 1,000 sq. ft. | 24 tablespoons (12 fl. ozs.) | 30 gallons |

To control certain White Grubs (larvae of European Chaffer, Southern Chaffer and Japanese Beetle) use two to four times the amounts of Hi-Yield DURSBAN SPRAY shown in above tables. Apply in enough water to obtain uniform coverage of the area treated followed immediately by thorough watering to wash the insecticide into the turf. Treat when grubs are young and actively feeding near the soil surfaces, usually during late July & August. Consult your local Agricultural Extension Service Specialist for proper time to treat in your area. To control Turfgrass (Hyperodes) weevils use twice the amount of Hi-Yield DURSBAN SPRAY shown in the above table. Spray suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

NOTE: To help prevent infestations of buildings by Ants, Crickets, Earwigs and Sowbugs (Pillbugs), treat a band of soil 6 to 10 feet around and adjacent to the building, also the building foundation to a height of 2-3 feet, where pests are active or may find entrance. Use the same dosage & dilution recommended for control of these pests on turf shown in above table. Apply as a coarse spray to thoroughly & uniformly wet the band area.

ORNAMENTAL PLANTS: On flowers, shrubs, vines, shade & flowering trees and evergreens infested with Aphids, Bagworms, Cutworms, Grasshoppers, Leafhoppers, Mealybugs, Mites, Spittlebugs, Thrips (exposed), & Whiteflies, use Hi-Yield DURSBAN SPRAY according to the following table. Apply with a low pressure sprayer or hose-on sprayer. Avoid over-spraying to the point of run-off. Treat when pests appear & repeat at 7-10 day intervals if needed. **IMPORTANT:** Do not use on Azaleas, Camellias, Poinsettias, Roses or Variegated Ivy because of possible injury to these plants.

DOGWOOD BORER: Apply to the trunks and limbs when adults begin to emerge. **PEACH TREE BORER:** Apply to flowering trees and shrubs of the Genus Prunus in water as a trunk spray before newly hatched larvae enter trees. Apply as coarse, low-pressure spray. Thoroughly wet bark areas from ground level to scaffold limbs. Consult your State Agricultural Extension Station or Extension Service specialist for proper time to treat in your area.

- Kills Certain White Grubs, Chinch Bugs, Sod Webworms (Lawn Moth Larvae), Ants, Brown Dog Ticks and Crickets in Lawns — Aphids, Cutworms, Grasshoppers, Mites, Whiteflies and Lilac, Ash, Dogwood and Peach Tree Borer on Flowers, Shrubs, and Trees.

- Use This Concentrate For Refilling Hi-Yield Ready-To-Use Kill-A-Bug Spray Containers.

KEEP OUT OF REACH OF CHILDREN
CAUTION
 See back panel for additional precautionary statements

OCT 7 1986

7401 294

12M 04

7401-294

10/7/86

Page 122

| DOSAGE TABLE FOR ORNAMENTAL PLANTS | | |
|--|---|---------------------------------------|
| Pests Controlled | Amount of Hi-Yield DURSBAN SPRAY in water to make | |
| | 1 gallon | 3 gallons |
| Aphids, Bagworms, Grasshoppers, Mites Spittlebugs, Whiteflies | 8 teaspoons | 8 tablespoons ($\frac{1}{2}$ cup) |
| Cutworms, Mealybugs, Leafhoppers & Thrips (exposed) | 16 teaspoons | 16 tablespoons (1 cup) |
| Ash Borer, Lilac Borer, Dogwood Borer & Peach Tree Borers | 1 Pint | 3 Pints |

*To prepare a refill for HI-Yield READY-TO-USE KILL-A-BUG, pour 4 $\frac{1}{4}$ ozs. of this product in the 1 quart container and fill with clean water. For the $\frac{1}{2}$ gallon size KILL-A-BUG container, pour 8 $\frac{1}{2}$ ozs. of this product in the $\frac{1}{2}$ gal. container and fill with clean water. Shake well before using.

STORAGE: Store in original container and place in a locked storage area.

DISPOSAL: Do not reuse container. Rinse thoroughly and securely wrap in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS

HAZAROUS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Wash contaminated clothing before reuse. Avoid breathing vapors or spray mist. Keep children and pets off treated areas until spray has dried. Do not spray vegetable gardens or other plants used for food purposes. Avoid contamination of food and feedstuffs. Food utensils such as a teaspoon or measuring cup should not be used for food purposes after use with insecticides.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED - Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED - Remove victim to fresh air and apply artificial respiration if indicated.

IF ON SKIN - Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES - Flush eyes with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: Chlorpyrifos is an cholinesterase inhibitor. Treat symptomatically. Atropine is antidote. Give by injection only.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith.

*DURSBAN - Trademark of Dow Chemical Co. U.S. Patent No. 3,244,586

Voluntary Purchasing Groups, Inc.

Bonham, Texas 75418

EPA Est. No. 7401-TX-1

EPA Reg. No. 7401-294

14M-7-115F

NET CONTENTS ONE QUART

292