PRECAUTIONARY STATEMENTS **HAZARDS TO HUMANS & DOMESTIC ANIMALS** CAUTION

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash hands after using. Keep children and domestic animals off treated areas until this material has been washed into the soil. Avoid contamination of feed and foodstuffs. Do not use in vegetable gardens. Do not contaminate water used for irrigation or domestic pur noses.

STATEMENT OF PRACTICAL TREATMENT If taken internally, induce vomiting and call a physician at once. In case of contact immediately flush ever or ekin with plenty of water. Get medical ACCEPTED attention if irritation pe

NOV 16 1979

Fungicials / E.S. & Act.

فلتستنز تشاح

Under the Federal Insecticide,

as anen en

EPA Heg. No.

registered under

Keep out of lakes, pond of equipment or dispos

It is a violation of Fede with its labeling.

1. WHAT IS THIS PAG This new product bas vegetation in areas w quickly inactivated on Kiler may be used arour and paties; along fenc also excellent for use

2. HOW DO YOU USE ferti-lome Improved V growing season, but best results are obtained during warm sunny weather

when applied to actively growing vegetation. Heavy lawn grass or difficult to control perennial weeds such as Bermudagrass and Dallisgrass may require more than one application to give complete control of top growth. DULITION DATE CHART

100 <u>sq. ft.</u>	(Sprinkling can) 12 fl. oz.	2 galions
250 sq. ft. 500 sq. ft. 1,000 sq. ft. 4,000 sq. ft.	(Tank or Hose-on Sprayer*) 8 fl. oz. 1 pint 1 quert 1 gellon	1 gallon 2-3 gallons 4-6 gallons 15-20 gallons
Area to be Treated	Amount of ferti-lome Improved Vegetation Killer to Use	Amount of Water to Use
	DISUITON NATE CHART	

*Follow directions on Hose-on Sprayer.

To kill undesirable vegetation around buildings, in brick and gravel walks and patios and along fences, sidewalks, driveways, curbs and gutters, apply ferti-lome improved Vegetation Killer with a tank sprayer, hose-on sprayer or sprinkling can using the above recommended rate. Cover unwanted vegetation thoroughly to point of runoff. Foliage will usually turn brown in about five days.

To kill undesirable vegetation around trees, shrubs, ornamentals and other confined areas, apply ferti-lome improved Vegetation Killer with a tank sprayer or sprinkling can la hose-on sprayer is not recommended in this case) using the above recommended rate. Do not allow spray solution to contact the leaves, green bark, bud crowns, or broken bark on trees,

> gh before treatment. Apply ik sprayer, hose-on sprayer led rate. Cover vegetation lly turn brown in about five ennials, or partially sprayed oply five days later if green

d be raked to remove dead ied according to local prac-

reated areas. Do not apply or aluminum. Clean spray with water.

EPA Est. 7401-TX-1

CTI

When used according to directions this new product will kill undesirable vegetation in listed areas and not harm the underlying roots of trees and shrubs.

STORAGE AND DISPOSAL

Do not store near feed, foodstuffs, fertilizers, seeds, insecticides or fungicides. Do not reuse empty container. Wrap container and put in trash collection NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. VOLUNTARY PURCHASING GROUPS, INC. BONHAM, TEXAS 75418

EPA Reg. No. 7401-300

