

Do not apply after pods begin to form. Pods are to be used as food, or within 7 days of harvest if pods are not to be used as food. Do not feed vines to dairy animals or animals being finished for slaughter.

Do not apply after seeding stage in leaf lettuce. Do not apply after heads begin to form on head lettuce.

Do not apply to green or maturing onions. Do not use after pods begin to form. On cauliflower, broccoli, and Brussels sprouts, do not apply after viable parts begin to form or within 30 days of harvest. On cabbage, do not apply after head starts to form nor feed treated plants to dairy animals or animals being finished for slaughter.

No time limitation. Apply as a foliar treatment only. Do not feed treated forage to dairy animals or animals being finished for slaughter.

Do not apply within 5 days of harvest.

Do not apply within 40 days of harvest after second cover spray. Do not apply within 30 days of harvest or after second cover spray. Do not apply after fruit starts to form. No time limitations.

Do not graze dairy animals or animals being finished for slaughter on these treated plants in the season. For early season control of these insects, make two to three applications at 10 day intervals beginning when cotton has three or four leaves to some classes may

Green webworm, Plant bug including tarnished plant bug, and spotted-winged bug, thrips, and spider mites. Do not apply to plants with yellow striped stems or worm, fireworm, or Texas.

3 1/2 pts./Acre

2 1/2 pts./Acre

Grasshopper, Cutworm, Corn borer, Lesser corn ear worm, Armyworm, Webworm, Corn earworm, Flea beetle.

4 pts./Acre

3 1/2 pts./Acre

2 1/2 pts./Acre

5 pts./Acre

2 pts./Acre

7 1/2 pts./Acre

Armyworm, Webworm, Grasshopper, Cutworm, Corn borer.

3 1/2 pts./Acre

4 pts./Acre

2 1/2 pts./Acre

5 1/2 pts./Acre

1 1/2 pts./Acre

4 pts./Acre

5 1/2 pts./Acre

4 pts./Acre

2 1/2 pts./Acre

be necessary to treat earlier, to prevent loss of stand by the pest or to insure, if this is necessary, more late or three late applications. On late planted cotton two applications may be sufficient. In any case, the last application should be made at least 30 days before bollworms usually appear for complete control of the pest. When sprays are made at 5 day intervals until three applications have been made or insects are under control, three applications at 5 day intervals are necessary to give time for all grades to hatch out after first application, these emerging adult weevils must be killed before they lay eggs. If any application is washed off by rain within 24 hours repeat application immediately. Do not apply after bolls open.

Do not apply more than once after heads have begun to form. Do not apply within 28 days of harvest when using up to 2 1/2 pts./acre. Do not apply within 40 days of harvest when using rates above 2 1/2 pts./acre. Do not feed treated forage to dairy animals or animals being finished for slaughter. Do not make treated forage.

May be applied with aerial or ground sprayers. With aerial application the product should be applied in 1 to 4 gallons per acre. Avoid spray drift. Do not apply less than 3 weeks before harvest. Do not feed treated plants or forage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Do not feed to, or use as mulch for, livestock or poultry.

Do not feed treated forage to dairy animals or animals being finished for slaughter. No limitation on use of gran.

Be certain it is a suitably diluted with water on a broadcast basis. Do not use as a post-emergence treatment. Do not graze dairy animals on stock. Do not graze treated plants on stock within 4 weeks of slaughter. No limitations on the use of gran.

Do not feed treated forage to dairy animals or animals being finished for slaughter. No restriction on the use of gran.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

***GENERAL INFORMATION ON COTTON**

Sprays have given more effective and more economical control of insects attacking cotton than dust. Because of the small size of the plants, a greater concentration of the insecticide is obtained from a spray. Very often, sprays may be applied during the day when dusting would be ineffective because much of the dust blows away.

Spray applications may be made at any time winds do not exceed 15 miles per hour. Spray when plants are dry so the material will stay on the leaves. Rain or heavy dew will wash off when leaves are wet enough to drip. For early season treatment with ground equipment, 1 more application of 6 1/2 lbs. per acre over the tops of the plants, is sufficient so long as plants are entirely covered. In some instances it may be necessary to use two sprays per acre. As plants increase in size increase the amount of insecticide to 12 lbs. per acre. Three sprays should be applied at a pressure of about 60 lbs. per acre. A water spray should be used at a nozzle pressure of 20 to 30 lbs. per acre when making early season treatments.

STORAGE AND DISPOSAL

Not for use or storage in or around the home environment. Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Triple rinse container and after for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place. Residue, spray mixture or residue that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Consult Federal, State or local disposal authorities for approved alternative procedures.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

POISONOUS BY SKIN CONTACT, INHALATION, OR SWALLOWING

Do not breathe spray mist. Avoid contamination of feed or foodstuffs. Do not use on dairy animals or in dairy barns. Wash thoroughly after using. Wash all contaminated clothing with soap and hot water before reuse. During application keep unprotected persons away from areas being treated or where there may be drift.

STATEMENT OF PRACTICAL TREATMENT

INTERNAL: If swallowed, do not induce vomiting. See Physician's Note. Never give anything by mouth to an unconscious person.

EXTERNAL: Remove clothing and wash skin thoroughly with soap and water. If in eyes, flush with clean water for at least 15 minutes.

CALL A PHYSICIAN IMMEDIATELY in all cases of suspected poisoning.

PHYSICIAN'S NOTE: If swallowed, use gastric lavage. Avoid emesis. Toxaphene produces central nervous system excitation that responds to barbituric acid derivatives. Prevent or control convulsions with barbiturates preferably administered intravenously. If the barbiturate can be given before convulsions start, sodium pentobarbital is the drug of choice. Since toxaphene has a fairly long duration of action, a long acting barbiturate such as phenobarbital may be used after initial control of convulsions. Refer to "Clinical Handbook of Economic Poisons" which was prepared by the U.S. Department of Health, Education, and Welfare, Public Health Service, Communicable Disease Center, Atlanta, Georgia.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. This product is toxic to bees and should not be applied when bees are actively visiting the area. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE POUR OR STORE NEAR HEAT OR OPEN FLAME. In case of spillage, use absorbent clay, hydrated lime, or other solids to soak up any liquid. Scrape up excess and bury residue away from water supplies. Wash area with strong detergent and hot water.

NOTICE TO DUSTERS/HANDLERS: In case of fire, leak, or damaged drums, or other emergency, report at once by toll free telephone to 800-424-9300.

NON-WARRANTY: Voluntary Purchasing Groups, Inc. warrants that this pesticide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this pesticide is reasonably fit for its intended purpose. Since timing, method of application, weather, type of plant, soil conditions, mixture with other chemicals, and other factors affecting the use of this pesticide are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

EPA Reg. No. 7401-302 VOLUNTARY PURCHASING GROUPS, INC. BONHAM, TEXAS EPA Est. 7401-TX-1

For Control of Thrips and Active Insects on Cotton. Toxaphene 67-62% Active Ingredient. Do Not Apply to Fruit or Seedlings.

to prevent
of cutworms, if
two of three later
area cotton, two
sufficient in any
should be made in
tolerance usually
control of that
city always, until
been made or
and three applica-
are necessary to
to hatch out after
emerging adult
before they lay
is washed off by
repeat application
apply only both
at once after reads
Do not apply within
then using up to 2 1/2
within 60 days of
years above 2 1/2
is needed for use to
keeping for use for
beats forage
near or ground
location the prod-
3 to 6 gallons per
Do not apply less
cotton. Do not feed
age made from
daily amounts, or
for slaughter or
wash to livestock or
age to dairy are
washed for slaughter
of grain.

GENERAL INFORMATION ON COTTON

Sprays have given more effective and more economic control of insects attacking cotton than dust. Because of the small size of the plants, a greater concentration of the insecticide is obtained from a spray. Very often, sprays may be applied during the day when dusting would be ineffective because of drift to the dust away.

STORAGE AND DISPOSAL

Not for use or storage in or around the home or where it is not contained in a safe container. Do not store in a storage or disposal. Open burning is prohibited. Triple-check containers and offers for recycling, reconditioning, or disposal in approved landfills, or bury in a safe place. Pesticide spray mixture or residue that cannot be used or chemically processed should be disposed of in a not approved for petroleum or buried in a safe place away from water supply. Consult Federal, State or local disposal authorities for approved alternative procedures.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

POISONOUS BY SKIN CONTACT, INHALATION, OR SWALLOWING
Do not breathe spray mist. Avoid contamination of feed or foodstuffs. Do not use on dairy animals or in dairy barns. Wash thoroughly after using. Wash all contaminated clothing in hot water before reuse. During application keep unprotected persons away from areas being treated or where drift may be drifting.

STATEMENT OF PRACTICAL TREATMENT

INTERNAL: If swallowed, do not induce vomiting. See Physician's Note. Never give anything by mouth to an unconscious person.
EXTERNAL: Remove contaminated clothing and wash skin thoroughly with soap and water. If in eyes, flush with clean water for at least 15 minutes.
CALL A PHYSICIAN IMMEDIATELY in all cases of suspected poisoning.

PHYSICIAN'S NOTE: If swallowed, use gastric lavage and/or sorption. Toxaphene produces central nervous system excitation that responds to barbiturates and demerol. Present or control convulsions with barbiturates, preferably administered intravenously. If the barbiturate is given before convulsions start, sodium pentobarbital is the drug of choice. Since toxaphene has a fairly long duration of action, a long acting barbiturate such as phenobarbital may be used after initial control of convulsions. Refer to Clinical Pharmacology of Economic Poisons, which was prepared by the U.S. Department of Health, Education, and Welfare, Public Health Service, Communicable Disease Center, Atlanta, Georgia.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be affected. This product is toxic to bees and should not be applied when bees are actively visiting the area. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Spraying and crab may be killed at application rates recommended on this label. Do not apply where there are important resources. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE, POUR OR STORE NEAR HEAT OR OPEN FLAME.
Wipe up spillage. Use absorbent, dry, hydrated lime, or other solids to soak up any liquid. Scrape up excess and bury residue away from water. Clean wash area with strong detergent and hot water.

NOTICE TO DRUG HANDLERS: In case of fire, injury, or damaged drums, or other emergency, report at once by toll free telephone to 800-824-9300.

NON-WARRANTY: Voluntary Purchasing Groups, Inc. warrants that this pesticide conforms to the Chemical description on its label. When used in accordance with label directions under normal conditions, this pesticide is reasonably fit for its intended purpose. Since timing, method of application, weather, type of plant, soil conditions, combine with other chemicals, and other factors affecting the use of this pesticide are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all responsibility for such use.

EPA Reg. No. 7401-302 VOLUNTARY PURCHASING GROUPS, INC. SCARBOROUGH, TEXAS EPA Est. 7401 TX 1

Hi-Yield[®] Toxaphene

ACCEPTED
JAN 3 0 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7401-302

Kills Insects on Ornamentals, Vegetables and Fruits
For Control of Boll Weevil, Bollworms, Grasshoppers, Webworms, Armyworms, Cutworms, Fleahoppers, Thrips and Other Listed Insects of Cotton, Grass and Grain Sorghum.

Active Ingredients:
Toxaphene (Technical Chlorinated Camphene Compound) 68.0%
67.65% Chlorine 23.0%
Xylene Range Aromatic Solvent 2.0%
Inert Ingredients: 100.0%

STATEMENT OF PRACTICAL TREATMENT
If Swallowed: Do not induce vomiting. See "Physician's Note." Never give anything by mouth to an unconscious person. Call a physician in all cases of suspected poisoning.
If Inhaled: Remove victim to fresh air. Call a physician in all cases of suspected poisoning.
If On Skin: Wash thoroughly with soap and water.
If In Eyes: Flush with clean water for at least 15 minutes. Call a physician.

**KEEP OUT OF REACH OF CHILDREN
WARNING**
SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS
NET CONTENTS ONE GALLON (5)

7401-302