

TOP

T

ACCEPTED

NOV 20 1980

EPA Reg. No. 7313-1



ELLENT  
**IPIC**

**GREEN**

**WATIVE**

MILDEW, ROT, DECAY  
WEATHER DAMAGE  
PREVENT WARPING AND SWELLING  
BELOW GROUND  
TREAT WITH WATER

**DIRECTIONS FOR USE**

Apply this product only as specified on this label. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Olympic Wood Preservative Green is specially formulated to treat wood used at or below ground level or in contact with water. It provides protection from mildew, rot and decay as well as termites and other wood destroying insects. Its water repellency also helps keep wood from warping and swelling. Wood Preservative Green meets TT-W-572B, the Federal specification for a water repellent wood preservative.

Wood Preservative Green combines penetrating oils with copper naphthenate, a preservative chemical which imparts a characteristic green color to wood. This green coloration will gradually fade if used on wood exposed to weather. Treated surfaces are not recommended to be finish coated with stain, paint or varnish type products.

For above ground preservative treatment where a semi-transparent stain color is desired, use Olympic Weather Screen. On wood used above ground where a finish coat of paint or stain is to be applied or a clear finish is desired, use Olympic Wood Preservative Clear. For severe exposure to wood rot, pressure treated wood offers maximum protection.

**SURFACE PREPARATION:** All surfaces must be clean, dry and free of mildew and bark. Remove all dirt, dust, oil, soot or other contaminants by washing with a detergent solution. Rinse thoroughly and allow to dry completely before application. Where mildew is evident, remove it by scrubbing the surface with this solution. One cup Trisodium Phosphate (TSP) or other detergent and one quart of household bleach dissolved in three quarts water (CAUTION: TSP and household bleach are potentially harmful materials. Follow manufacturer's precautions. Use protective clothing, eye goggles and good rubber gloves.) Rinse thoroughly with fresh water and immediately apply a solution of 50% household bleach and 50% water. Allow this final treatment to dry before applying Olympic Wood Preservative Green.

**APPLICATION:** Stir before using. Olympic Wood Preservative Green should be applied liberally to achieve maximum performance. Dipping for several minutes is recommended. Two coats applied by brush or spray are acceptable. Treat ends and edges of boards thoroughly and separate while drying. DO NOT THIN OR INTERMIX WITH OTHER COATING PRODUCTS.

FOR ADDITIONAL INFORMATION—ASK YOUR DEALER

ACTIVE INGREDIENTS  
Copper Naphthenate 21.22% minimum  
(equivalent to metallic copper-2%)  
INERT INGREDIENTS 78.78% maximum  
TOTAL 100.00%

MEETS FED. SPEC. TT-W-572B  
EPA Reg. No. 7313-1 EPA Est. No. 7313-WA-1, 7313-IL-1, 7313-KY-1

BEST DOCUMENT AVAILABLE

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Hi-Yield® Lawn and Garden Fungicide is an excellent all-around fungicide that is effective in controlling a wide variety of diseases on lawns, shrubs, flowers, vegetables and fruits. Although this product will control established diseases, best results are obtained when used as a protective spray on a regular basis. This will reduce the risk and severity of disease damage at lower rates and fewer fungicide applications. Higher rates of application are needed to bring established diseases under control.

**LAWN AND TURF GRASSES** For control of Brown Patch (Rhizoctonia), Copper Spot, Slime Mold (black sooty mold), Melting Out (Helminthosporium Leaf Spot), Pink Patch (Fusarium Thread) and Rust on crown, stem and leaf. Apply 4 ounces in 15 gallons of water per 1000 sq. ft. (or 5 level teaspoons in one gallon of water per 66 sq. ft.) at 14-day intervals as a protective spray or before diseases become well established. When conditions favor the occurrence of diseases, use 6 to 8 ounces in 15 gallons of water per 1000 sq. ft. (2 1/2-3 1/2 tablespoons in one gallon of water per 66 sq. ft.) at weekly intervals. As a protective spray for Dollar Spot (Sclerotinia) and Pythium Blight, use 6 to 8 ounces in 15 gallons of water at 5-day intervals. Repeat applications until the threat of disease is over. For control of Snow Mold (Fusarium), apply during winter using 6 to 8 ounces in 15 gallons of water at 2 to 6-week intervals.

**DOGS** Do not graze treated areas. Do not feed clippings to livestock.

**ROSES AND FLOWERS** Make a full coverage spray, diluting 1 1/2 tablespoons in one gallon of water. Apply when plants are leafed out well and spray at 7 to 10-day intervals throughout the growing season. Use this dilution rate to control the diseases mentioned below.

**Roses.** To control Black Spot, apply when plants are well leafed out at weekly intervals during growing season. Gladiolus. To control flower and leaf spots caused by Botrytis and Curvularia. Spray at weekly intervals before flower spikes appear prior to appearance of disease and at 2 to 4 day intervals after diseases appear. Do not apply after spikes appear. Carnation, Iris and Hollyhock. To control Leaf Spot start applying when plants are well leafed out and repeat at weekly intervals. Lily, Tulips and Dahlia. Apply to control Botrytis blight. Pansies, Hollyhock and Flowering Dogwood. Apply to control Rust and Anthracnose. Ornamental Crabapples. Apply to control Scab and Cedar Apple Rust. Petal Blight on Azalea, Camellia, Rhododendron. Spray at 2 to 4 day intervals while flowers are opening, thoroughly covering flowers and ground beneath bushes.

**VEGETABLES** Tomatoes. To control Gray Leaf Spot, Late and Early Blight, Gray Leaf Mold and Anthracnose, use 8 ounces of Hi-Yield® Lawn and Garden Fungicide per 25 gallons of water (2 level tablespoons per gallon of water). Spray plants adequately providing thorough coverage. Start spraying upon emergence of seedlings and repeat applications throughout the season at weekly intervals. Do not apply within 5 days of harvest. Potatoes. To control Late and Early Blight, use same rate as for tomatoes. Start spraying plants when 4 to 6 inches tall and repeat application throughout the season at 7 to 10-day intervals. Sweet Corn. To control Helminthosporium leaf blight, use same rate as for tomatoes and apply when disease first appears and repeat at 4 to 7-day intervals. Do not apply within 7 days of harvest. Do not (continued on next page)

feed treated sweet corn forage or fodder to livestock. Cucumbers, Summer Squash, Cantaloupe and Watermelons. To control Downy Mildew, Scab, Anthracnose, Alternaria and Cercospora leaf spots, Gummy stem blight. Spray thoroughly using the same rate as for tomatoes. Be sure that both upper and lower leaf surfaces are covered. Begin spraying when plants are in 2-leaf stage and repeat at 7-day intervals. Do not apply within 5 days of harvest.

**FRUIT TREES** Apples, Pears, Crabapples and Quince. To control bitter rot, black rot, brown rot, scab, cedar apple rust, fly speck and sooty blotch use 8 ounces in 25 gallons of water (2 tablespoons per gallon of water). Apply every one or two weeks throughout the season. For scab, begin at green tip or delayed dormant stage, for cedar apple rust begin at first cover. Do not apply within 15 days of harvest (30 days for apples). Do not graze livestock in treated areas. Consult your local or State Agricultural Extension Service for timing and number of sprayings. Grapes. For control of Black rot, bunch rot, downy mildew and dead arm use 8 ounces in 17 gallons of water (3 tablespoons per gallon of water) to provide complete coverage. Start spraying when shoots are 1/2 to 1 1/2 inches long and reapply when shoots are 3 to 5 inches long, and again when shoots are 8 to 10 inches long. Continue spraying at 10 to 14-day intervals. Do not apply within 66 days of harvest. In California, do not apply after fruit is

# Hi-Yield® (Maneb) Lawn and Garden Fungicide

**Controls Many Diseases  
on Lawns, Shrubs,  
Flowers, Vegetables  
and Fruits**

A Wettable Powder that

**CONTROLS THE FOLLOWING PLANT DISEASES:** Brown Patch, Dollar Spot, Leaf Spot, Snow Mold, Early & Late Blight, Downy Mildew, Anthracnose, Scab, Gray Leaf Mold, Copper Spot, Melting Out, Slime Mold, Rust, Pythium Blight, Pink Patch, Cedar Apple Rust, Black Spot, Petal Blight, Botrytis Blight, Bitter Rot, Black Rot, Brown Rot, Sooty Blotch, Fly Speck.

**ACTIVE INGREDIENTS.**

A co-ordination product of Zinc ion and Manganese Ethylene Bisdithiocarbamate .....	80%
Manganese ** .....	16%
Zinc ** .....	2%
Ethylene bisdithiocarbamate ion (C4H6N2S4) .....	62%
<b>INERT INGREDIENTS .....</b>	<b>20%</b>
<b>Total .....</b>	<b>100%</b>

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

**STOP DIS**

Do not reuse purpose. Store container and p

**PRECAUTION HAZARDS DOME**

May cause irrit and skin. Avoid met. Avoid clothing. Keep Store in a cool product to be storage. This m changes which effectiveness an be flammable. **FIRST AID:** In water and get lowed, give 1 o and induce vo of throat with vomiting or gi an unconscio attention.

**ENVIRON**

This product is leaves, streams when weather areas treated. by cleaning of water.

EPA R  
EPA E  
VOLUNT  
OF  
BOR  
NET  
(22

7401-303

27

# Hi-Yield<sup>®</sup> (Maneb) Lawn and Garden Fungicide

**Controls Many Diseases  
on Lawns, Shrubs,  
Flowers, Vegetables  
and Fruits**

A Wettable Powder that

**CONTROLS THE FOLLOWING PLANT DISEASES: Brown Patch, Dollar Spot, Leaf Spot, Snow Mold, Early & Late Blight, Downy Mildew, Anthracnose, Scab, Gray Leaf Mold, Copper Spot, Molding Rot, Slime Mold, Rust, Pythium Blight, Pink Patch, Cedar Apple Rust, Black Spot, Petal Blight, Botrytis Blight, Bitter Rot, Black Rot, Brown Rot, Sooty Blotch, Fly Speck.**

**ACTIVE INGREDIENTS.**

A co-ordination product of Zinc ion and Manganese Ethylene Bisdiethiocarbamate

Manganese Ethylene Bisdiethiocarbamate	80%
Manganese	16%
Zinc	2%
Ethylene bisdiethiocarbamate ion (C <sub>4</sub> H <sub>8</sub> N <sub>2</sub> S <sub>4</sub> )	62%
<b>INERT INGREDIENTS</b>	<b>20%</b>
<b>Total</b>	<b>100%</b>

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

**STORAGE AND  
DISPOSAL**

Do not reuse empty container for any purpose. Store in a cool dry place. Wrap container and put in trash collection.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS**

**CAUTION**

May cause irritation to nose, throat, eyes and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Keep away from fire or sparks. Store in a cool, dry place. Do not allow product to become wet or overheated in storage. This may lead to certain chemical changes which will reduce the fungicide's effectiveness and create vapors which may be flammable.

**FIRST AID:** In case of skin contact, wash with water. If in eyes, wash with plenty of water and get medical attention. If swallowed, give 1 or 2 glasses of warm water and induce vomiting by touching back of throat with fingers. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

EPA Reg. No. 7401-303  
EPA Est. No. 7401-TX-1

**VOLUNTARY PURCHASING  
GROUPS, INC.  
BONHAM, TEXAS**

**NET WT. 8 OZS.  
(227.0 grams)**

27A

**USE**

Apply in a manner

to obtain the best results. Although the best results are obtained with regular applications, the damage at lower rates and less frequent applications are not serious.

**Lawns:** Brown Patch, Black Sooty Mold, Pink Patch (Red Patch). Apply 4 ounces in 15 gallons of water in one gallon of water as a protective spray or when conditions favor disease in 15 gallons of water.

Apply to lawn clippings to

prevent disease, during dry weather throughout the season.

**Flowers and Shrubs:** Apply 4 ounces in 15 gallons of water (2 tablespoons per gallon of water) to control blight, black rot, brown rot, scab, cedar apple rust, fly speck and sooty blotch.

**Vegetables:** Apply 4 ounces in 15 gallons of water (2 tablespoons per gallon of water) to control blight, black rot, brown rot, scab, cedar apple rust, fly speck and sooty blotch.

Apply to sweet corn forage or fodder to livestock. Cucumbers, Summer Squash, Cantaloupe and Watermelons. To control Downy Mildew, Scab, Anthracnose, Alternaria and Cercospora leaf spots, Gummy stem blight. Spray thoroughly using the same rate as for tomatoes. Be sure that both upper and lower leaf surfaces are covered. Begin spraying when plants are in 2-leaf stage and repeat at 7-day intervals. Do not apply within 5 days of harvest.

**FRUIT TREES:** Apples, Pears, Crab-apples and Quince. To control bitter rot, black rot, brown rot, scab, cedar apple rust, fly speck and sooty blotch, use 8 ounces in 25 gallons of water (2 tablespoons per gallon of water). Apply every one or two weeks throughout the season. For scab, begin at green tip or delayed dormant stage; for cedar apple rust, begin at first cover. Do not apply within 15 days of harvest (30 days for apples). Do not graze livestock in treated areas. Consult your local or State Agricultural Extension Service for timing and number of sprayings. Grapes. For control of Black rot, bunch rot, downy mildew and dead arm. Use 8 ounces in 17 gallons of water (3 tablespoons per gallon of water) to provide complete coverage. Start spraying when shoots are 1 to 1 1/2 inches long and reapply when shoots are 3 to 5 inches long, and again when shoots are 8 to 10 inches long. Continue spraying at 10 to 14-day intervals. Do not apply within 66 days of harvest. In California, do not apply after fruit.

303

27

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Hi-Yield® Lawn and Garden Fungicide is an excellent all-around fungicide that is effective in controlling a wide variety of diseases on lawns, shrubs, flowers, vegetables and fruits. Although this product will control established diseases, best results are obtained when used as a protective spray on a regular basis. This will reduce the risk and severity of disease damage at lower rates and fewer fungicide applications. Higher rates of application are needed to bring established diseases under control.

**LAWN AND TURF GRASSES** For control of Brown Patch (Rhizoctonia), Copper Spot, Slime Mold (black sooty mold), Melting Out (Helminthosporium Leaf Spot), Pink Patch (Red Thread), and Rust on crown, stem and leaf. Apply 4 ounces in 15 gallons of water per 1000 sq. ft. (or 5 level teaspoons in one gallon of water per 66 sq. ft.) at 14-day intervals as a protective spray or before diseases become well established. When conditions favor the occurrence of diseases, use 6 to 8 ounces in 15 gallons of water per 1000 sq. ft. (2 1/2-3 1/2 tablespoons in one gallon of water per 66 sq. ft.) at weekly intervals. As a protective spray for Dollar Spot (Sclerotinia) and Pythium Blight, use 6 to 8 ounces in 15 gallons of water at 5-day intervals. Repeat applications until the threat of disease is over. For control of Snow Mold (Fusarium), apply during winter using 6 to 8 ounces in 15 gallons of water at 2 to 6-week intervals.

**NOTE:** Do not graze treated areas. Do not feed clippings to livestock.

**SHRUBS AND FLOWERS** Make a full coverage spray, diluting 1 1/2 tablespoons in one gallon of water. Apply when plants are leafed out well and spray at 7 to 10-day intervals throughout the growing season. Use this dilution rate to control the diseases mentioned below.

**Roses:** To control Black Spot, apply when plants are well leafed out at weekly intervals during growing season. **Gladiolus:** To control flower and leaf spots caused by Botrytis and Curvularia. Spray at weekly intervals before flower spikes appear prior to appearance of disease and at 2 to 4 day intervals after diseases appear. Do not apply after spikes appear. **Carnation, Iris and Hollyhock:** To control Leaf Spot start applying when plants are well leafed out and repeat at weekly intervals. **Lily, Tulips and Dahlia:** Apply to control Botrytis blight. **Pansies, Hollyhock and Flowering Dogwood:** Apply to control Rust and Anthracnose. **Ornamental Crabapples:** Apply to control Scab and Cedar Apple Rust. **Petal Blight on Azalea, Camellia, Rhododendron:** Spray at 2 to 4 day intervals while flowers are opening, thoroughly covering flowers and ground beneath bushes.

**VEGETABLES:** **Tomatoes:** To control Gray Leaf Spot, Late and Early Blight, Gray Leaf Mold and Anthracnose, use 8 ounces of Hi-Yield® Lawn and Garden Fungicide per 25 gallons of water (2 level tablespoons per gallon of water). Spray plants adequately providing thorough coverage. Start spraying upon emergence of seedlings and repeat applications throughout the season at weekly intervals. Do not apply within 5 days of harvest. **Potatoes:** To control Late and Early Blight, use same rate as for tomatoes. Start spraying plants when 4 to 6 inches tall and repeat application throughout the season at 7 to 10-day intervals. **Sweet Corn:** To control Helminthosporium leaf blight, use same rate as for tomatoes and apply when disease first appears and repeat at 4 to 7-day intervals. Do not apply within 7 days of harvest. Do not

(continued on side gutter)

feed treated sweet corn forage or fodder to livestock. **Cucumbers, Summer Squash, Cantaloupe and Watermelons:** To control Downy Mildew, Scab, Anthracnose, Alternaria and Cercospora leaf spots, Gummy stem blight. Spray thoroughly using the same rate as for tomatoes. Be sure that both upper and lower leaf surfaces are covered. Begin spraying when plants are in 2-leaf stage and repeat at 7-day intervals. Do not apply within 5 days of harvest.

**FRUIT TREES:** **Apples, Pears, Crabapples and Quince:** To control bitter rot, black rot, brown rot, scab, cedar apple rust, fly speck and sooty blotch use 8 ounces in 25 gallons of water (2 tablespoons per gallon of water). Apply every one or two weeks throughout the season. For scab, begin at green tip or delayed dormant stage, for cedar apple rust begin at first cover. Do not apply within 15 days of harvest (30 days for apples). Do not graze livestock in treated areas. Consult your local or State Agricultural Extension Service for timing and number of sprayings. **Grapes:** For control of Black rot, bunch rot, downy mildew and dead arm. Use 8 ounces in 17 gallons of water (3 tablespoons per gallon of water) to provide complete coverage. Start spraying when shoots are 1/2 to 1 1/2 inches long and reapply when shoots are 3 to 5 inches long, and again when shoots are 8 to 10 inches long. Continue spraying at 10 to 14-day intervals. Do not apply within 66 days of harvest. In California, do not apply after fruit.

# Hi-Yield® (Maneb) Lawn and Garden Fungicide

**Controls Many Diseases  
on Lawns, Shrubs,  
Flowers, Vegetables  
and Fruits**

A Wettable Powder that

**CONTROLS THE FOLLOWING PLANT DISEASES:** Brown Patch, Dollar Spot, Leaf Spot, Snow Mold, Early & Late Blight, Downy Mildew, Anthracnose, Scab, Gray Leaf Mold, Copper Spot, Melting Out, Slime Mold, Rust, Pythium Blight, Pink Patch, Cedar Apple Rust, Black Spot, Petal Blight, Botrytis Blight, Bitter Rot, Black Rot, Brown Rot, Sooty Blotch, Fly Speck.

**ACTIVE INGREDIENTS:**

A co-ordination product of Zinc ion and  
Manganese Ethylene Bisdithiocarbamate ..... 80%  
Manganese \*\* ..... 16%  
Zinc \*\* ..... 2%  
Ethylene bisdithiocarbamate  
ion (C4H6N2S4) ..... 82%  
**INERT INGREDIENTS** ..... 20%  
Total ..... 100%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

**STORAGE  
DISP**

Do not reuse empty purpose. Store in a container and put it

**PRECAUTIONS  
HAZARDS TO  
DOMESTI**

**CAU**

May cause irritation and skin. Avoid breath. Avoid contact clothing. Keep away. Store in a cool, dry product to become storage. This may be changes which will effectiveness and be flammable. **FIRST AID:** In case with water. If in eyes water and get medical. Give 1 or 2 gl and induce vomiting of throat with finger. Vomiting or give an unconscious person attention.

**ENVIRONMENT**

This product is toxic lakes, streams or ponds when weather conditions are treated. Do not by cleaning of equipment.

**EPA Reg. No.**

EPA Est. No.

**VOLUNTARY GROUP**

**BONHAM**

**NET WT.**

**(227.0 g)**

7401-303

27

**BEST DOCUMENT AVAILABLE**

**INSTRUCTIONS FOR USE**

Always use this product in a manner consistent with the label.

Maneab Fungicide is an excellent all-around fungicide for controlling a wide variety of diseases on vegetables and fruits. Although this fungicide controls a wide variety of diseases, best results are obtained by spraying on a regular basis. This fungicide controls disease damage at lower rates of application. Higher rates of application are required for diseases under control.

**VEGETABLES:** For control of Brown Patch, Blight, Slime Mold (black sooty mold), Downy Mildew, Leaf Spot, Pink Patch (Red stem and leaf). Apply 4 ounces in 15 gallons of water (or 5 level tablespoons in one gallon) at 7 to 10-day intervals as a protective spray or when disease first appears. When conditions favor disease, use 6 to 8 ounces in 15 gallons of water (or 7 to 10 level tablespoons in one gallon) at 7 to 10-day intervals. Repeat applications until the disease is controlled. For control of Snow Mold (Fusarium), use 6 to 8 ounces in 15 gallons of water at 7 to 10-day intervals.

**FRUIT TREES:** Do not feed clippings to livestock.

Make a full coverage spray, diluting with water. Apply when plants are in the 2-leaf stage and repeat at 7 to 10-day intervals throughout the growing season to control the diseases listed below.

**FRUIT TREES:** Apples, Pears, Crabapples and Quince. To control bitter rot, black rot, brown rot, scab, cedar apple rust, fly speck and sooty blotch use 8 ounces in 25 gallons of water (2 tablespoons per gallon of water). Apply every one or two weeks throughout the season. For scab, begin at green tip or delayed dormant stage, for cedar apple rust begin at first cover. Do not apply within 15 days of harvest (30 days for apples). Do not graze livestock in treated areas. Consult your local or State Agricultural Extension Service for timing and number of sprayings. Grapes: For control of Black Rot, bunch rot, downy mildew and dead arm use 8 ounces in 17 gallons of water (3 tablespoons per gallon of water) to provide complete coverage. Start spraying when shoots are 1/2 to 1 1/2 inches long and reapply when shoots are 3 to 5 inches long, and again when shoots are 8 to 10 inches long. Continue spraying at 10 to 14-day intervals. Do not apply within 66 days of harvest. In California, do not apply after fruit has begun to ripen.

**FRUIT TREES:** Apples, Pears, Crabapples and Quince. To control bitter rot, black rot, brown rot, scab, cedar apple rust, fly speck and sooty blotch use 8 ounces in 25 gallons of water (2 tablespoons per gallon of water). Apply every one or two weeks throughout the season. For scab, begin at green tip or delayed dormant stage, for cedar apple rust begin at first cover. Do not apply within 15 days of harvest (30 days for apples). Do not graze livestock in treated areas. Consult your local or State Agricultural Extension Service for timing and number of sprayings. Grapes: For control of Black Rot, bunch rot, downy mildew and dead arm use 8 ounces in 17 gallons of water (3 tablespoons per gallon of water) to provide complete coverage. Start spraying when shoots are 1/2 to 1 1/2 inches long and reapply when shoots are 3 to 5 inches long, and again when shoots are 8 to 10 inches long. Continue spraying at 10 to 14-day intervals. Do not apply within 66 days of harvest. In California, do not apply after fruit has begun to ripen.

7401-303

**FRUIT TREES:** Apples, Pears, Crabapples and Quince. To control bitter rot, black rot, brown rot, scab, cedar apple rust, fly speck and sooty blotch use 8 ounces in 25 gallons of water (2 tablespoons per gallon of water). Apply every one or two weeks throughout the season. For scab, begin at green tip or delayed dormant stage, for cedar apple rust begin at first cover. Do not apply within 15 days of harvest (30 days for apples). Do not graze livestock in treated areas. Consult your local or State Agricultural Extension Service for timing and number of sprayings. Grapes: For control of Black Rot, bunch rot, downy mildew and dead arm use 8 ounces in 17 gallons of water (3 tablespoons per gallon of water) to provide complete coverage. Start spraying when shoots are 1/2 to 1 1/2 inches long and reapply when shoots are 3 to 5 inches long, and again when shoots are 8 to 10 inches long. Continue spraying at 10 to 14-day intervals. Do not apply within 66 days of harvest. In California, do not apply after fruit has begun to ripen.

**FRUIT TREES:** Apples, Pears, Crabapples and Quince. To control bitter rot, black rot, brown rot, scab, cedar apple rust, fly speck and sooty blotch use 8 ounces in 25 gallons of water (2 tablespoons per gallon of water). Apply every one or two weeks throughout the season. For scab, begin at green tip or delayed dormant stage, for cedar apple rust begin at first cover. Do not apply within 15 days of harvest (30 days for apples). Do not graze livestock in treated areas. Consult your local or State Agricultural Extension Service for timing and number of sprayings. Grapes: For control of Black Rot, bunch rot, downy mildew and dead arm use 8 ounces in 17 gallons of water (3 tablespoons per gallon of water) to provide complete coverage. Start spraying when shoots are 1/2 to 1 1/2 inches long and reapply when shoots are 3 to 5 inches long, and again when shoots are 8 to 10 inches long. Continue spraying at 10 to 14-day intervals. Do not apply within 66 days of harvest. In California, do not apply after fruit has begun to ripen.

27

# Hi-Yield<sup>®</sup> (Maneb) Lawn and Garden Fungicide

## Controls Many Diseases on Lawns, Shrubs, Flowers, Vegetables and Fruits

A Wettable Powder that

**CONTROLS THE FOLLOWING PLANT DISEASES:** Brown Patch, Dollar Spot, Leaf Spot, Snow Mold, Early & Late Blight, Downy Mildew, Anthracnose, Scab, Gray Leaf Mildew, Copper Spot, Mosaic, Mottling Out, Slime Mold, Rust, Pythium Blight, Pink Patch, Cedar Apple Rust, Black Spot, Petal Blight, Botrytis Blight, Bitter Rot, Black Rot, Brown Rot, Sooty Blotch, Fly Speck.

**ACTIVE INGREDIENTS**  
A co-ordination product of Zinc ion and Manganese Ethylene Bisdithiocarbamate ..... 80%  
Manganese .. 16%  
Zinc .. 2%  
Ethylene bisdithiocarbamate ion (C<sub>4</sub>H<sub>6</sub>N<sub>2</sub>S<sub>4</sub>) .. 62%  
**INERT INGREDIENTS** ..... 20%  
Total ..... 100%

### KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

### STORAGE AND DISPOSAL

Do not reuse empty container for any purpose. Store in a cool dry place. Wrap container and put in trash collection.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

May cause irritation to nose, throat, eyes and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Keep away from fire or sparks. Store in a cool, dry place. Do not allow product to become wet or overheated in storage. This may lead to certain chemical changes which will reduce the fungicide's effectiveness and create vapors which may be flammable.  
**FIRST AID:** In case of skin contact, wash with water. If in eyes, wash with plenty of water and get medical attention. If swallowed, give 1 or 2 glasses of warm water and induce vomiting by touching back of throat with fingers. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

EPA Reg. No. 7401-303  
EPA Est. No. 7401-TX-1  
**VOLUNTARY PURCHASING GROUPS, INC.**  
BOWHAM, TEXAS  
**NET WT. 8 OZS.**  
(227.0 grams)

28

