

Hi-Yield®

Maneb Lawn & Garden Fungicide

Controls Many Diseases
on Lawns, Shrubs,
Flowers, Vegetables
and Fruits

A Wettable Powder that
CONTROLS THE FOLLOWING DISEASES AS SPECIFIED:
Brown Patch, Helminthosporium Blight, Melting Out, Dollar
Spot, Black Spot, Petal Blight, Botrytis Blight, Alternaria
Blight, Leaf Spot, Anthracnose, Early and Late Blight, Gray
Leaf Spot (Stemphylium), Septoria Leaf Spot, Downy Mildew,
Rust, Pythium Fruit Rot, Black Rot, Bitter Rot, Botryosphaeria
(White Rot), Cedar-apple Rust, Flyspeck, Scab, Sooty Blotch,
Brown Rot, Shotholes, Peach Leaf Curl.

• 1½ Tablespoons Makes 1 Gallon Spray For Shrubs,
Flowers, Vegetables and Fruits.

ACTIVE INGREDIENT:
Maneb (Manganese ethylenebisdithiocarbamate) 80%
(Manganese equivalent as metallic 16.5%)
INERT INGREDIENTS: 20%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel For Additional Precautionary Statements

ACCEPTED

DEC 14 1983

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the purposes
registered under
EPA Reg. No. 7460-703

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Hi-Yield Maneb Lawn & Garden Fungicide is effective for controlling a wide variety of diseases on lawns, shrubs, flowers, vegetables and fruits. When used as directed, this product can control listed diseases and thus help plants grow stronger, healthier and more productive.

Although this product will control established diseases, best results are obtained when used as a protective spray on a regular basis. Where higher and lower rates are given, use lower rates of application for protective sprays and higher rates of application when disease conditions are severe.

LAWN & TURF GRASSES: Apply the amount recommended after each disease listed below in enough water (15 gallons) to cover 1,000 sq. ft. Brown Patch, Helminthosporium Blight and Melting Out—Use 3-5 ozs. Begin at first sign of disease and repeat applications at 7-14 day intervals. Leaf, Stem and Stripe Rusts—Use 3-5 ozs. Begin when rust pustules are first seen. Repeat at 7-14 day intervals. Dollar Spot—Use 6-8 ozs. Begin at first sign of disease and repeat at 7-14 day intervals.

NOTE: Do not graze livestock on treated areas. Do not feed clippings to livestock.

SHRUBS & FLOWERS: Spray to cover all plant surfaces thoroughly. Apply when plants are leafed out. Use 1½ tablespoons per gallon of water to make spray mix.

Roses—Black Spot—Begin when first leaves unfold. Repeat at 7-10 day intervals.

Azaleas, Camellias—Petal Blight—Apply to litter and soil beneath plants, also to flowers of azaleas. Begin when flower buds swell. Repeat at 3-5 day intervals.

Geraniums, Hydrangeas, Lilies—Botrytis Blight—Begin when new growth starts, before disease appears or at first sign of disease. Repeat at 7-10 day intervals. Carnations, Pansies—Alternaria Blight or Leaf Spot, Anthracnose, Botrytis Blight—Begin when new growth starts. Repeat weekly. Dogwood—Anthracnose—Begin when buds open.

Repeat when bracts fall, 4 weeks later and in late summer.

VEGETABLES & FRUITS: Use 1½ level tablespoon per gallon of water. Spray to thoroughly cover upper and lower surfaces of foliage. Unless otherwise directed, repeat applications every 7-10 days as needed. Spray more frequently if disease conditions are severe. Number in () indicates the number of days to observe between last application and harvest for vegetables.

Tomatoes(5)—For Anthracnose, Early & Late Blight, Gray Leaf Spot (Stemphylium), Septoria (Leaf) Spot. Begin application when tomatoes on first clusters are well formed. Do not use on young plants in greenhouse.

Beans(4) (Green & Dry)—For Downy Mildew, Rust. Begin when plants are small. Repeat every 5-7 days.

Cucumbers(5), Squash(5)—For Anthracnose, Downy Mildew, Pythium Fruit Rot. Begin when disease first appears or runners start to run.

(continued on side gusset)

Pepper(0), Eggplant(0)—For Anthracnose, Fruit Rot—Apply as fruit begins to form. Repeat weekly through harvest.

Carrots(0)—For Leaf Spot diseases. Start spraying when plants are 6 weeks old or disease threatens.

Spinach(10)—For Downy Mildew, White Rust. Remove residues at harvest by washing.

Apples—For Black Rot, Bitter Rot, Botryosphaeria (White Rot), Cedar-apple Rust, Flyspeck, Scab and Sooty Blotch—Begin at prebloom, calyx, or first cover stages. Repeat at 7-14 day intervals as needed. Do not graze livestock in treated areas. Do not apply within 15 days of harvest in Arkansas, Delaware, Illinois, Indiana, Kansas, Kentucky, Maryland, Missouri, New Jersey, North Carolina, Ohio, Pennsylvania, South Carolina, Tennessee, Virginia and West Virginia. Do not apply within 30 days of harvest in other states.

Peaches—Brown Rot, Scab, Shotholes. Begin at red bud. Repeat in early bloom, full bloom, petal fall, and 7-14 day intervals up to 2 weeks before harvest. Do not apply within 14 days of harvest. Peach Leaf Curl—Apply in spring dormant period. Do not apply within 2 days of harvest. If applied within 14 days of harvest, remove residues by brushing.

STORAGE

Do not contaminate storage or sleep place.

Pesticide Disposal: Dispose of container in cover and discard in trash. Container Disposal: Discard bag in trash.

PRECAUTIONS/HAZARDS

CAUTION: May irritate and skin. May be swallowed. Avoid mist. Avoid contact with clothing. Protectively wear long-sleeved shirt, hat and boots) mowing and loading. A food and feedstuff.

STATEMENT OF PESTICIDE ON SKIN—RINSE clothing and wash soap and water. (If with plenty of water, if irritation persists.)

ENVIRONMENT

This product is toxic directly to waters. water by cleaning disposal of wastes.

PHYSICAL OR CHEMICAL

Do not use, pour, or open flame.

Buyer assumes all or handling of this accordance with herewith. Voluntary Inc. warrants this the chemical description for the purposes of limit of any liability purchase price paid.

Voluntary Purch Bonham
EPA Reg. N
EPA Reg. I
NET W

Hi-Yield®

ACCEPTED

DEC 14 1983

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the purposes registered under EPA Reg. No. 7401-303

Maneb Lawn & Garden Fungicide

Controls Many Diseases
on Lawns, Shrubs,
Flowers, Vegetables
and Fruits

Pepper(0), Eggplant(0) - For Anthracnose, Fruit Rot - Apply as fruit begins to form. Repeat weekly through harvest.

Carrots(0) - For Leaf Spot diseases. Start spraying when plants are 6 weeks old or disease threatens.

Spinach(10) - For Downy Mildew, White Rust. Remove residues at harvest by washing.

Apples - For Black Rot, Bitter Rot, Botryosphaeria (White Rot), Cedar-apple Rust, Flyspeck, Scab and Sooty Blotch - Begin at prebloom, calyx, or first cover stages. Repeat at 7-14 day intervals as needed. Do not graze livestock in treated areas. Do not apply within 15 days of harvest in Arkansas, Delaware, Illinois, Indiana, Kansas, Kentucky, Maryland, Missouri, New Jersey, North Carolina, Ohio, Pennsylvania, South Carolina, Tennessee, Virginia and West Virginia. Do not apply within 30 days of harvest in other states.

Peaches - Brown Rot, Scab, Shot-holes. Begin at red bud. Repeat in early bloom, full bloom, petal fall, and 7-14 day intervals up to 2 weeks before harvest. Do not apply within 14 days of harvest. Peach Leaf Curl - Apply in spring dormant period. Do not apply within 2 days of harvest. If applied within 14 days of harvest, remove residues by brushing.

A Wettable Powder that
CONTROLS THE FOLLOWING DISEASES AS SPECIFIED:
Brown Patch, Helminthosporium Blight, Melting Out, Dollar Spot, Black Spot, Petal Blight, Botrytis Blight, Alternaria Blight, Leaf Spot, Anthracnose, Early and Late Blight, Gray Leaf Spot (Stemphylium), Septoria Leaf Spot, Downy Mildew, Rust, Pythium Fruit Rot, Black Rot, Bitter Rot, Botryosphaeria (White Rot), Cedar-apple Rust, Flyspeck, Scab, Sooty Blotch, Brown Rot, Shot-holes, Peach Leaf Curl.
• 1½ Tablespoons Makes 1 Gallon Spray For Shrubs, Flowers, Vegetables and Fruits.

ACTIVE INGREDIENT:
Maneb (Manganese ethylenebisdithiocarbamate) 80%
(Manganese equivalent as metallic 16.5%)
INERT INGREDIENTS: 20%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel For Additional Precautionary Statements

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Store in a cool, dry place.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse bag. Discard bag in trash.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION: May irritate eyes, nose, throat and skin. May be harmful if inhaled or swallowed. Avoid breathing dust or spray mist. Avoid contact with skin, eyes or clothing. Protective clothing (long pants, long sleeve shirt, impermeable gloves, hat and boots) must be worn during mixing and loading. Avoid contamination of feed and feedstuffs.

STATEMENT OF PRACTICAL TREATMENT
IF ON SKIN - Remove contaminated clothing and wash affected areas with soap and water. IF IN EYES - Flush eyes with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to waters. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this material conforms to the chemical description on the label and for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the user or buyer.

Voluntary Purchasing Groups, Inc.
Benham Texas 75418
EPA Est. No. 7401 TX 1
EPA Reg. No. 7401 303
NET WT 9 OZS

is a violation of Federal law to use inconsistent with its labeling.

Maneb Fungicide is effective for controlling diseases on lawns, shrubs, flowers, and trees. When used as directed, this product can help plants grow stronger, healthier, and more productive.

For control of established diseases, best used as a protective spray on a regular basis. When used as directed, this product can help plants grow stronger, healthier, and more productive.

Apply the amount recommended after mixing in enough water (15 gallons) to cover the plants. For Helminthosporium Blight and Melting Out, apply at first sign of disease and repeat applications. For Leaf, Stem and Stripe Rusts - apply when rust pustules are first seen. Repeat applications at 7-14 day intervals.

Apply to cover all plant surfaces thoroughly. Use 1½ tablespoons per gallon of water when first leaves unfold. Repeat at 7-14 day intervals.

Apply to litter and soil beneath plants. Begin when flower buds swell.

For Botrytis Blight - Begin when new growth appears or at first sign of disease. For Carnations, Pansies, Alternaria Blight, Anthracnose, Botrytis Blight - Begin when first signs of disease appear.

Apply 4 weeks later and in late summer. Use 1½ level tablespoons per gallon of water. Thoroughly cover upper and lower surfaces of leaves. Repeat applications every 7-14 days, more frequently if disease conditions are severe. () indicates the number of days before harvest for vegetables.

For Anthracnose, Early & Late Blight, Gray Leaf Spot, Leaf Spot. Begin application when leaves are well-formed. Do not use on young plants.

For Downy Mildew, Rust. Begin when first signs of disease appear. Repeat every 5-7 days.

For Anthracnose, Downy Mildew. Begin when disease first appears or at first sign of disease.

See back panel for additional precautionary statements.