PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals.

CAUTION

May cause eye irritation. Avoid contact to eyes. Do not get on skin or clothing. Harmful if swallowed. Keep children and pets off of treated area until spray had dried.

Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with this product.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed -Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If Inhaled -Remove victim to fresh air. Apply artificial respiration if indicated.

If On Skin -Remove contaminated clothing and wash affected areas with soap and water.

If In Eyes -Flush eyes with fresh water for 15 minutes. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly into water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

STORAGE AND DISPOSAL

Avoid contamination of food or feed. When empty container is discarded, wrap container and put in trash collection.

Important: Do not apply this product when rainfall is expected within 6 hours. Do not use in galvanized or unlined steel containers. Do not use on windy days or when weather favors drift. Protect from freezing.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

KILLS

Quackgrass
Fescue
Bahiagrass
Creeping
Beggarweed
Witchweed
Ragweed
Foxtail
Sandbur
Smartweed
Milkweed
Cocklebur
Pigweed
Peppergrass
Velvetleaf
Shattercane
Panicum
Knotweed
Frenchweed
Butter Cup
Mallow
Wild Aster
Wild Carrot
Yarrow
Puncturevine
Morning Glory
English Daisy
Chicory
Fleabane
Lambsquarters
Spanishneedles
Stitch Wort
Wood Sorrel
Wintercress
Stinging Nettle
Oxalis
Wild Barley
Maiden Cane
and many others

Hi-Yield®

Systemic

Weed and Grass Killer

Before		After

- 8 ozs. Makes 1 Gallon of Spray
- Kills Plants on Contact, Roots and All.
- Can Be Used In Sensitive Areas Such As Around Shrubs, Flower Beds, and Trees.

ACTIVE INGREDIENTS:
Isopropylamine salt of *glyphosate.. 5.00%
INERT INGREDIENTS 95.00%
Total 100.00%

*N-(phosphonomethyl) glycine

NET CONTENTS 32 FL. OZS. (ONE QUART)

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

KILLS

Bermudagrass
Johnsongrass
Crabgrass
Nutgrass
Dallis Grass
Henbit
Spurge
Wild Onion
Knotweed
Docks
Clever
St. Augustine
Bluegrass
Purslane
Chickweed
Ryegrass
Poison Oak
Poison Ivy
Thistle
Mullein
Bendweed
Kochia
Cat Tails
Nightshade
Dogbane
Napiergrass
Moneywort
Galingsoga
Self-heal
Vervains
Shepherd's Purse
Wild Radish
Sheep Sorrel
Mustards
Healall
Dog Fennel
Cats-ear
and many others

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with it's labeling.

This product is a Non-selective Herbicide that will eliminate undesirable vegetation in lawns, driveways, parking areas, between stepping stones and bricks, around fences, patios, flower beds, gardens, buildings, trees, and shrubs without harm to untreated plants and vegetation. One application will kill most weeds and grasses. See list on both sides of front panel of this label for a partial list of weeds and grasses controlled.

To prepare a spray solution, mix 8 fl. ozs. (1 cup) of this concentrate with 1 gallon of clean water.

Make a full coverage application being sure to thoroughly wet plant surfaces. Avoid cutting treated vegetation for at least 7 days. If certain hard to kill weeds and grasses are not completely killed in 14 to 21 days repeat application.

For spot treatment in lawns, flower beds, ornamental beds, around trees and shrubs, apply carefully to avoid contact with desirable vegetation. If spray or drift contacts desirable vegetation, wash off immediately with water.

For areas where renovation and reseeding is desired, make a thorough coverage of lawn area. Treated area should begin to turn brown within 7 days, then reseed or sod as desired.

*To prepare a refill for Hi-Yield Ready-To-Use Systemic Weed and Grass Killer, pour 3¼ ozs. of this product into the 1 Quart container and finish filling with clean water.

*To prepare a refill for Hi-Yield Systemic Vegetation Killer, pour 13 ozs. of this product into the 1 Gallon container and finish filling with clean water.

VOLUNTARY PURCHASING GROUPS, INC.
BONHAM, TEXAS

EPA Est. No. 7401-TX-1   EPA Reg. No. 7401-306

2 of 2