

7401-311

1-15-1982

1/2

<p align="center">U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567) WASHINGTON, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION</p> <p align="center"><i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i></p>	EPA REGISTRATION NO. 7401-311	DATE OF ISSUANCE 15 January 1982
	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT Ferti-Lone Fire Blight Spray	
NAME AND ADDRESS OF REGISTRANT (Include ZIP code)		
<p align="right"><i>Best Available copy</i></p> <p align="right">161 56306 18 1</p> <p>Voluntary Purchasing Groups, Inc. P.O. Box 460 Conlan, TX 75418</p>		
<p>NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p> <p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p> <p>Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with the provisions of Section 3(c)(7)(A) of the Act, since you have agreed that you will submit and/or cite all data required for registration/reregistration of your product under FIFRA Section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.</p> <p>If these conditions are not complied with, the registration will be subject to cancellation in accordance with Section 6(e) of the Act.</p> <p>Henry M. Jacoby Product Manager (1) Fungicide-Herbicide Branch Registration Division (TS-767)</p> <p><input type="checkbox"/> ATTACHMENT IS APPLICABLE</p>		
SIGNATURE OF APPROVING OFFICIAL		DATE

PD = N/A

NO VARNISH

ACCEPTED

JAN 15 1982

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product is registered under EPA Reg. No. 7401-311

2 level tablespoons of ferti-lome FIRE BLIGHT SPRAY per 2½ gallons water (200 ppm concentration). Replant in soil free from disease organisms, and apply ½ tablespoon per 2½ gallons water (50 ppm concentration) as foliage spray at weekly intervals.

3. HERE ARE THE RESULTS YOU SHOULD EXPECT!

When used according to directions this product will control Fire Blight and Bacterial Wilt, Stem Rot, Leaf Spot, and Crown Gall on Fruit Trees, Flowers and Shrubs.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS**

May cause allergic skin reaction. Avoid contact with skin and eyes. Do not breathe dust or spray mist. Wear a dust mask and rubber gloves. Wash thoroughly after handling. NOTE: ferti-lome Fire Blight Spray is a nonsterile grade of streptomycin and is not to be used for medicinal purposes.

STATEMENT OF PRACTICAL TREATMENT
In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS
This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste.

STORAGE & DISPOSAL
Do not reuse empty container. Wrap container and put in trash collection.

*Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

VOLUNTARY PURCHASING GROUPS, INC.
Bonham, Texas
EPA Est. No. 7401-TX-1
EPA Reg. No. 7401-311



ferti-lome



Fire Blight Spray

- For Apples, Pears, Pyracantha, Chrysanthemums, Philodendron, Dieffenbachia, and Roses.
- 1 Tablespoon Makes 2½ Gallons of Spray.

ACTIVE INGREDIENT:
Streptomycin Sulfate 21.2%

INERT INGREDIENTS: 78.8%

100.0%

*Contains 17% Streptomycin derived from 21.2% Streptomycin Sulfate.

**KEEP OUT OF REACH
OF CHILDREN
CAUTION
SEE BACK PANEL FOR
ADDITIONAL PRECAUTIONARY
STATEMENTS**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. WHAT IS THIS PACKAGE FOR?
Use this product to control Fire Blight and Bacterial Wilt, Stem Rot, Leaf Spot, and Crown Gall on Fruit Trees, Flowers and Shrubs.

2. HOW DO YOU USE IT?
FRUIT: APPLES & PEARS (Areas Other Than West Coast): To control FIRE BLIGHT in home plantings, apply 1 level tablespoon in 2½ gallons of water (equivalent to 100 ppm or 4 ozs. per 50 gallons of water) in foliar and blossom sprays. Make first spray at the start of blossoming period. Continue spray applications every 3 to 4 days during blossom time. Apply additional sprays every 5 to 7 days after blossom period when weather conditions favor spread of fire blight. Do not apply when fruit is visible.

PYRACANTHA: To control FIRE BLIGHT, apply a solution of 1 level tablespoon of ferti-lome FIRE BLIGHT SPRAY per 2½ gallons water (100 ppm concentration). Begin spraying at start of blossom period. Repeat at 3 to 4 day intervals during bloom and at 5 to 7 day intervals after bloom if weather favors disease spread.

CHRYSANTHEMUMS (Cuttings): To control BACTERIAL WILT, soak cuttings for 4 hours in a solution containing ½ tablespoon of ferti-lome FIRE BLIGHT SPRAY per 2½ gallons of water (50 ppm concentration) and plant as usual.

DIEFFENBACHIA: To control BACTERIAL STEM ROT, spray plants at 5 to 7 day intervals with a solution of 1 level tablespoon of ferti-lome FIRE BLIGHT SPRAY in 2½ gallons water (100 ppm concentration) or soak cuttings for 20 minutes in a solution of 2 level tablespoons in 2½ gallons of water (200 ppm concentration) and plant in sterile rooting medium.

PHILODENDRON: To control BACTERIAL LEAF SPOT, remove all decayed leaves and spray with a solution of 2 level tablespoons of ferti-lome FIRE BLIGHT SPRAY in 2½ gallons of water (200 ppm concentration) as preventative or at first sign of water soaked leaves. Repeat at 4 to 5 day intervals.

ROSES: To control CROWN GALL, remove infested plant and cut out gall tissue. Soak root system and cut surfaces of infested area for 15 minutes in a solution of

c/c