

7401-318

(Front Panel)

3-10-87

1/3

ferti-lome(R)

KILLS

KILLS

- Henbit
- Chickweed
- Clover
- Spurge
- Crabgrass
- Sandspur
- Wiggrass
- Pigweed
- Yellow Nutsedge
- Knotweed
- Thistle
- Barnyardgrass
- Lambsquarters
- Wild Oaks
- Fern
- and many
others

- Ragweed
- Cudweed
- Moneywort
- Dichondra
- Florida
Betony
- Witchweed
- Morning
Glory
- Spurweed
- Kochia
- Annual
Bluegrass
- Narrowleaf
Vetch
- Swinecress
- Cransbill
- Nightshade
- Velvetleaf
- Smutgrass
- and many
others

Premergent

Weed & Grass

Control

ACCEPTED

MAR 10 1987

Kills 38 Weeds & Grasses.

7401-318

For St. Augustine, Dormant Bermudagrass, and Centipede Grass

1 1/2 Tablespoons Covers 500 sq.ft.

Active Ingredients:

Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine-.....	40.8%
Related Compounds.....	2.2%
Inert Ingredients:.....	57.0%

Keep Out Of Reach Of Children

CAUTION

See Back Panel For Additional Precautionary Statements

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. WHAT IS THIS PRODUCT FOR?

Use this product to effectively kill unsightly weeds in lawns and recreational areas. See the front panel for a partial list of weeds controlled.

2. HOW DO YOU USE IT?

Mix with water and apply uniformly to the lawn. Apply with a pump-up or compressed air type sprayer or with a hose-on type sprayer. ferti-lome Premergent Weed & Grass Control will kill both emerged weeds and weeds from seeds. Rain or watering within 2 or 3 days of application may decrease the effectiveness on emerged weeds. However, for control of weeds from seeds, rainfall or watering is necessary within 7 to 10 days after treatment.

ST. AUGUSTINE & CENTIPEDE GRASSES, ferti-lome Premergent Weed & Grass Control may be applied to established St. Augustine and Centipede grasses during both the dormant and the growing season. Best results are usually obtained in the early Spring of dormant periods when weeds are small or have not emerged.

DORMANT BERMUDAGRASS: Apply to Bermudagrass during the dormant season. Applications to Bermudagrass during the growing season will cause injury. Follow rates on table below.

DICHONDRA, MONEYWORT: The best control for these weeds can be obtained by making an application in early April followed with a second application in July. Do not apply to growing Bermudagrass. Applications for Spurweed or Florida Betony generally will give control or suppression of the other weeds listed. However, as a general rule ferti-lome Premergent Weed & Grass Control will give best control when applied to young tender weeds or just prior to weed emergence.

FLORIDA BENTONYL: This weed emerges in the fall, so an application in mid to late February would give the best control.

SPURWEED: The best control of spurweed can be obtained by applying spray when spurweed has emerged (December and January).

RATE OF APPLICATION : Determine the total area to be sprayed and base rate of application in the chart below. Avoid overlapping spray pattern while treating. Shake contents well before mixing.

AREA TO BE TREATED	AMOUNT OF PRODUCT	SUGGESTED AMOUNT OF WATER
500 sq.ft.	1 1/2 tbs.	1 gal.
1,000 sq.ft.	3 tbs.	2 gal.
3,000 sq.ft.	4 1/2 ozs.	6 gal.
5,450 sq.ft.	8 ozs.	10 gal.
10,900 sq.ft.	1 pint	20 gal.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

This very economical and easy-to-use product will effectively rid lawns and recreational areas of annoying and unsightly problem weeds.

STORAGE: Store in original container and place in a locked storage area.
DISPOSAL: Do not reuse container. Wrap and put in trash.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful or fatal if swallowed. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep children and pets off the treated area until spray has dried. Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with this product.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed- Call a physician or Poison Control Center immediately. If possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
If Inhaled - Remove victim to fresh air. Apply artificial respiration if indicated.
If on Skin - Remove contaminated clothing and wash affected areas with soap and water.
If in Eyes - Flush eyes with plenty of water. Call a physician immediately.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Atrazine can travel (seep or leach) through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

NOTE: Do not apply within the active root zones of vegetables or desirable ornamental plants such as Camellias, Azaleas, Boxwoods, etc.. Do not plant any crop (flower or vegetable) to treated area for 18 months or injury may result.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Voluntary Purchasing Groups, Inc.

Bonham, Texas 75418

A Est. No. 7401-TX-1

EPA Reg. No. 7401-318

NET CONTENTS _____ Fl. OZS.

