

(Front Panel)

Hi-Yield

Lindane

Spray

ACCEPTED
SEP 28 1983
U.S. Environmental Protection Agency
Washington, D.C. 20460
EPA Reg. No. 7461-321

- * Kills Ticks, Chiggers, Fleas, Leafhoppers, Borers, Bark Beetles, Leaf Miners, Aphids, Thrips, Cucumber Beetles, Lacebugs, Turpentine Beetles, Roaches, Ants, Sarcoptic Mange, Scab Mites, Lice, Stable Flies, Mosquitoes, & Keds.
- * Kills Insects on Lawns, Shrubs, Flowers, Livestock & In Livestock Quarters.
- * Kills Borers on Fruit & Shade Trees.

ACTIVE INGREDIENTS:

Gamma Isomer of Benzene	
Hexachloride from Lindane.....	10.0%
Aromatic Petroleum Derivatives	85.0%
INERT INGREDIENTS:	5.0%
	<u>100.0%</u>

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

RECORDED IN INDEX

1401 2
(Right Panel)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is formulated to effectively kill destructive and disease spreading insects in lawns, and on shrubs, trees, flowers and livestock. Kills Ticks, Chiggers, Fleas, Leafhoppers, Borers, Bark Beetles, Leaf Miners, Aphids, Thrips, Cucumber Beetles, Lacebugs, Turpentine Beetles, Roaches, Ants, Scarcotic Mange, Scab Mites, Lice, Flies, Mosquitoes, and Keds.

LAWNS

TICKS, CHIGGERS, FLEAS, LEAFHOPPERS: Use 2 pints on 10,000 sq. ft. or 6 tablespoons on 1,000 sq. ft. in sufficient water for good coverage. Repeat application if necessary.

SHADE TREES, ORNAMENTAL SHRUBS & FLOWERS

APHIDS, THRIPS, LEAFMINERS, CUCUMBER BEETLES: On Boxwood, Azaleas, Roses, Chrysanthemum, Larkspur, and other flowers, shrubs and shade trees, use two pints per 100 gallons of water or 2 teaspoonsful to 1 gallon of water.

LACEBUGS: On Azalea, Pyracantha and Rhododendron, use 3 pints per 100 gallons water or 3 teaspoons to 1 gallon. Apply in spring when immature insects first appear and retreat as necessary.

PINE BARK BEETLES, TURPENTINE BEETLES: Use 2 pints in 5 gallons of water and apply to trunk of pines to a height of 6 feet at the rate of 1 gallon per 50 sq. ft. of bark surface. Apply at first sign of attack and repeat in 3 to 6 months if necessary.

DOGWOOD BORER: Use 6 pints per 100 gallons of water or 2 tablespoons to 1 gallon and apply to trunk and lower limbs in May and June before borers enter the trunk. Apply thoroughly, particularly to injured areas and old borer wounds. Repeat applications 2 to 3 times at 10 day intervals.

FRUIT TREES

ROUNDHEADED BORER & FLATHEADED BARK BEETLES: On Apple, Peach and Cherry, use 6 pints to 100 gallons of water or 2 tablespoons to 1 gallon of water. Apply to trunk, crotch to ground. Soak bark to runoff. Apply to bark of tree and large limbs in spring. Do not apply to fruit or foliage. Do not apply within 60 days of harvest.

PEACH TREE BORER: On Peach, use 6 pints to 100 gallons of water or 2 tablespoons to 1 gallon of water. Apply to trunk, crotch to ground. Soak bark to runoff. Apply to bark of tree and large limbs in spring. Do not apply within 60 days of harvest.

LIVESTOCK

CAUTION: Do not use on dairy animals. Do not use on emaciated, sick, overheated or lactating animals. Do not use on poultry. Do not treat animals in cold stormy weather. Do not spray animal within 30 days or dip animals within 60 days of slaughter. Thoroughly stir dipping vats before using. If any separation of emulsion occurs, do not use dip.

BEEFCATTLE: LICE & TICKS: Use 2½ pints per 100 gallons of water. Thoroughly spray entire animal. **SARCOPTIC MANGE AND SCAB MITES** - Spray, using 4 pints per 100 gallons of water. Make only 2 application, 7 days apart. Do not apply to calves under 6 months of age at this dosage.

SWINE: LICE & SARCOPTIC MANGE MITES - Use 4 pints per 100 gallons of water and apply as a spray or dip. For mange mites, repeat treatment in 2 weeks if necessary. Do not treat sows within two weeks of farrowing nor for at least three week thereafter. Do not apply to swine under three months of age.

SHEEP & GOATS: TICKS, KEDS, LICE- Use 4 pints per 100 gallons of water as a spray or use 2 pints per 100 gallons in a dip. Repeat treatment in 2 to 3 weeks if necessary.

BEST DOCUMENT : 1401

7401-321

(Right Panel Cont.)

RESIDUAL SPRAYS

STABLE FLY, MOSQUITOES, ROACHES, ANTS: Inside and outside of barns, hog pens, stables and other buildings, use 3 to 4 pints to 10 gallons of water and apply at the rate of 1 gallon to 1,000 sq. ft. of surface. Thoroughly cover resting surfaces for flies and mosquitoes. Apply to cracks, crevices, and other hiding places for control of roaches and ants. Repeat applications as necessary. Do not apply in milkrooms or poultry houses. Do not contaminate animal feed or water.

TICKS & CHIGGERS: Use 4 quarts per 100 gallons of water and spray weeds and under brush harboring pests. Repeat applications as necessary. Do not use on food crops or grazing areas.

(Left Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep children and pets off treated areas until spray has dried.

Food utensils such as a teaspoon or measuring cup should not be used for these purposes after use with pesticides.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

STATEMENT OF PRACTICAL TREATMENTS

- If swallowed - Call a physician or Poison Control Center immediately. If possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
- If inhaled - Remove victim to fresh air. Apply artificial respiration if indicated.
- If on skin - Remove contaminated clothing and wash affected areas with soap and water.
- If in eyes - Wash eyes with plenty of water. Call a physician immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

7401-321 4

(Left Panel Cont.)

STORAGE AND DISPOSAL

Do not contaminate food or feed by storage or disposal. Do not reuse empty container. Wrap container and put in trash collection.

*Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Voluntary Purchasing Groups, Inc.
Benham, Texas 75418

WPA Est. No. 7401-Ta-1

EPA Reg. No. 7401-321

Net Contents One Quart