PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation of spray mist. May be absorbed through skin. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed and food stuffs. Keep children and pets off treated areas until this material is washed into the soil and grass is dry

Food utensils such as teaspoons and measuring cups should not be used for food numposes after use with nesticides

STATEMENT OF PRACTICAL TREATMENT

if swallowed

If in eyes

Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vorniting or give anything by mouth to an unconscious

If inhaled

Remove victim to fresh air. Apply artificial respiration if indicated Remove contaminated clothing and wash affected areas

If no skip

with soap and water Wash eyes with plenty of water. Call a physician if irrita-

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not graze treated areas to livestock. Apply this product only as specified on this label

WARRANTY-CONDITION OF SALE: DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of Voluntary Purchasing Groups, Inc. or the seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result. ALL SUCH RISKS ARE ASSUMED BY THE BUYER.

Voluntary Purchasing Groups, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use as modified by the above. Voluntary Purchasing Groups, Inc. makes no other warranties, expressed or implied, including FITNESS or MER-CHANTABILITY. In no case shall Voluntary Purchasing Groups, Inc. or the Seller be hable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by Voluntary Purchasing Groups, Inc. and is accepted as such by the Buyer.

> Voluntary Purchasing Groups, Inc. Bonnam, Texas 75418

EEA 5 it No. 7401 TX 1

Net Contents One Gallon

EPA Red No 7:01 324




ferti-lome



Quik-Kill™ Concentrate Weed & Grass Killer

 Kills Weeds & Grasses On Contact, In Flower Beds. Lawns, and Around Trees, Shrubs, Patios, Walks, Drive-ways, and Fences.



Sroad Spectrum Weed Killer For Use on Hose-on Wank Type Sprayers.

ACTIVE INGREDIENTS:

Sodium Cacodylate..... Dimethylarsinic Acid (Cacodylic Acid) INERT INGREDIENTS

Total Arsenic (as elemental) all in water soluable form 12.75% Product Contains 2 48 lbs. Cacodylic Acid Equivalent per gallon &

KEEP OUT OF REACH OF CHILDREN

CA ION

SEE BACK PANEL FOR ADDITIONAL ECAUTIONARY STATEMENTS



73 39%

100 00%

It is a violation of Federal law to use this product in a manner inconsistent with its

 WHAT IS THIS PACKAGE FOR.

This product will effectively kill unwanted weeds and glasses on contact in and around lawns, flower beds, fences, sidewalks, drivewals, buildings, trees & shrubs, right-of-ways, vacant lots, roadsides and industrial plant sites without the production of the productio harming untreated vegetation. It is ideal for post-emergent weed control and is non-selective in its actions. It is quickly inactivated on contact with soil thus leaving no residual soil activity to harm desired plant growth.

2. HOW DO YOU USE IT? Z. HOW DO YOU DON!!?
This completely water soluble product can be used in any spray equipment that gives good coverage. It is spray ank about half full with water and add the required amount of herdicide with agitation. Finish filling the lank with water and apply. To apply with a hose on sprayer, add required amount of product in serayer jar and finish filling jar with water to make desired number of gallons of spray. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period. Although this product is only moderately corrosive, do not use in galvanized or aluminum equipment

LAWN & GARDEN AREAS: To kill unwanted weeds and grasses in and around lawns and garden areas, mix 11/2 to 21/2 ozs. of this product per gallon of water (1-2 gallons per 100 gallons of water) and spray to wet thoroughly surfaces of unwanted weeds and grasses. For best results apply during warm, sunny weather. Heavy tawn grasses and difficult to control perennial weeds such as Berantidagrass and Dallis Grass may require more than one application to give complete control of top growth. For spot treatments in lawns, flower beds, and around trees and strubs, apply carefully to avoid contact with desirable vegetation. LAWN RENOVATIONS: Mow lawn to 1 inch high. Mix 10 fl. ozs. of product in 4 gallons of water and apply to 1,000 sq. ft. Spray foliage thoroughly. Foliage will usually turn brown in about five days. Bermudagrass, deep rooted perennials, or partially sprayed foliage, may require a second treatment. Reapply five days later if green areas remain. When foliage is brown, rake to remove dead vegetation, then re-establish according to local practice. INDUSTRIAL To use in and around industrial plant sites, buildings, fences, drainage ditch banks, vacant lots. roadsides, railroad, storage areas, highways, utility and pipeline right-of-ways and other similar areas. Mix 1 to 2 gallons of this product in 100 gallons of water 11-2 Ots, per 20 pallons of water). Spray unwanted vegetation to just short of run-off, If regrowth occurs reapply as necessary. Do not allow spray to come in contact with foliage, green bark, grafted unions, or scuffed, damaged or broken bark when spraying around trees and desirable vegetation. AGRICULTURAL PLANTINGS-CITRUS (Except Florida): This product is useful as a directed application in non-bearing citrus orchards such as orange, grapefruit, tangerine, lemon and lime orchards. It should be applied at the rate of 1 $^\circ$ 2 to 2 gallons per acre. Mix 1 to 2 gallons of this product in 50 to 100 gallons of water. Apply as a directed spray in interspaces and around base of trees. Spray unwanted vegetation to just short of run-off. If regrowth occurs, reapply as regulred; however, do not exceed 3 applications per year. Do not allow spray solution to contact leaves, stems or bank. Use a shield, if necessary, for nursery plantings or young trees. Do not apply around trees from which fruit will be harvested within one year of treatment.

3. HERE ARE THE RESULTS YOU SHOULD EXPECT!

Effective and economical control of unsightly and nuisance weed buildup in fawns, industrial and agricultural areas.