PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CALITION- Harmful if swallowed. Avoid initialation of snrav mist. May be absorbed through skin. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep children and pets off treated areas until this material is washed into the soil and grass is dry

Food utensils such as teaspoons and measuring cups should not be used for food pur poses after use with pesticides

STATEMENT OF PRACTICAL TREATMENT

| IF SWALLOWED | Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vormiting by touching back of throat with finger. Do not induce vormiting or give anything by mouth to an unconscious berson. | |
|--------------|---|--|
| FINHALED | Remove victim to fresh air. Apply artificial respiration if in- dicared | |
| IF ON SKIN | Remove contaminated clothing and wash affected areas with | |

soap and wate IF IN EYES Wash eyes with plenty of water. Call a physician if irritation persists

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Do not graze treated areas to livestock. Apply this product only as specified on this label.

WARRANTY CONDITION OF SALE: DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully, It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of Vokuntary Purchasing Groups, Inc. or the seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result. ALL SUCH RISKS ARE ASSUMED BY THE BUYER.

Voluntary Purchasing Groups, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use as modified by the above. Voluntary Purchasing Groups, Inc. makes no other warranties, expressed or implied, including FITNESS or MERCHANTABILITY. In no case shall Voluntary Purchasing Groups, Inc. or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by Voluntary Purchasing Groups, Inc. and is accepted as such by the Buver

NET CONTENTS

Voluntary Purchasing Groups, Inc. Bonham, Texas 75418

EPA Est. No. 7401-TX-1

EPA Reg. No. 7401-237 /

Phytar 560 Herbicide (Contains Cacodylic Acid)

 Kills Weeds & Grasses On Contact, In and Around Lawns, Flower Beds. Driveways, Fences, Buildings, Trees & Shrubs, Roadsides, Right-of-Ways, Vacant Lots and Industrial Plant Sites.

22.73%

3.88%

73.39%

100.00%

| For Post-Emergent | Weed Control. |
|---------------------------------------|---------------|
|---------------------------------------|---------------|

ACTIVE INGREDIENTS:

| Socium Cacodylate | | | | | | |
|---|------|------|------|------|--|------|
| Dimethylarsinic Acid (Cacodylic Acid) | | | | | | |
| INERT INGREDIENTS: | | | | | | |

Total Arsenic (as elemental) all in water soluble form 12.75% Product Contains 2.48 lbs. Cacodylic Acid Equivalent per gallon.

> KEEP OUT OF REACH OF CHILDREN CAUTION SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its lab-

Apply this product to effectively kill unwanted weeds and grasses on contact in and around lawns tiower beds, tences, sidewalks, driveways, buildings, trees & shrubs right of ways vacant lots, roadsides, and industrial plant sites without harming un treated vegetation. It is ideal for post-emergent weed control and is non-selective in its actions. It is quickly inactivated on contact with soil thus leaving no residual soil activity to harm desired plant growth

This complete - water soluble product can be used in any spray equipment that gives good coverage. Fill a spray tank about haif full with water and add the required amount of herbicide with agitation. Finish filling the tank with water and apply. To apply with a hose on spraker, add required amount of product in sprayer (ar and finish filling the (ar with water to make the desired number of gallons of spray. After use, clean equipment thoroughly by tushing with water. Do not store spray solution in tank for a prolonged period. Although this product is only moderately corrosive ido not use in galvanized or aluminum en coment

LAWN & GARDEN AREAS: To kill unwanted weeds and grasses in and around lawns and garden areas, mix 1% biss to 2': ozs. of this product per gallon of water (1.2 gallons per 100 gallons of water) and spray to wet thoroughly surfaces of unwanted weeds and grasses. For best results apply during warm, sunny weather. Heavy lawn grasses and difficult to control perennial weeds such as Bermudagrass and Dallis Grass may require more than one application to give complete control of top growth. For spot treatments in lawns, flower beds, and around trees and shrubs, apply carefully to avoid contact with casirable vegetation. LAWN RENOVATIONS Mow lawn to 1 inch high.

Mix 10 fl. ozs of product in 4 gallons of water and apply to 1,000 sq. ft. Spray foliage thoroughly. Follage will usually turn brown in about five days. Bermudagrass, deeprooted perennials, or partially sprayed foliage, may require a second treatment. Reapply five days later if green areas remain. When foliage is brown, rake to remove dead vegetation, then re-establish according to local practice

INDUSTRIAL: To use in and around industrial plant sites, buildings, fences, drainage ditch banks, vacant lots, roadsides, railroad, storage areas, highways, utility and pipeline right-of-ways and other similar areas. Mix 1 to 2 gallons of this product in 100 gallons of wates (1-2 Qts, net 20, gallons of water). Spray unwanted vegetation to just short of run-off. If regrowth occurs reapply as necessary. Do not allow spray to come in contact with foliage, greer bark, grafted unions, or scuffed, damaged or broken bark when spraying arounal trues and austrable vegetation.

AGRICULTUFAL PLANTING CITR'IS (Except Florida): This product is useful as a directed application in non-bearing citrus orchards such as orange, grapefruit, tangerine, lemon and lime orchards. It should be applied at the rate of 1 ½ to 2 gallons per acre. Mix I to 2 gallons of this product in 50 to 100 gallons of water. Apply 55 a directed spray in intryspaces and around bage of trees. Spray unwanted vegetation to just short of runoffull regrowth occurs reapplisas required; however, do not exceed 3 applications per yes? Do nate low splay solution to contact leaves, stems or bark. Use a shield, if hecessary, for dursery plantidge or young trees. Do not apply around trees from which True will be harvested within one year of treatment

DISPOSAL Do not reuse empty container, Wrap container and put in trash collection