

true leaf stage or when conditions are favorable for disease development. Re apply at 7 day intervals or as necessary under severe conditions to maintain control. CANTALOUPE, MUSKMELON, HONEYDEW (O). To control Anthracnose & Downy Mildew. Apply when plants are in first true leaf stage or when conditions are favorable for disease development. Re apply at 7 day intervals. Under severe conditions shorten application interval. POTATO (O). To control Early & Late Blight. Apply when plants are 6 inches high or when disease threatens and continue at 7-10 day intervals or as needed to maintain control.

3. HERE ARE THE RESULTS YOU MAY EXPECT!
This superior product will control a broad spectrum of plant diseases for beautiful flowers and lawns and for healthier, more productive vegetable gardens.

STORAGE & DISPOSAL

Do not reuse empty container. Wrap empty container and put in trash collection.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes and clothing. Avoid breathing of dust. Wash thoroughly after handling and before eating and smoking. Avoid contamination of feed or foodstuffs. Keep children and pets off treated areas until this material is washed into the soil.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. **IF ON SKIN**, remove contaminated clothing and wash affected areas with soap and water. **IF IN EYES**, flush eyes with plenty of water and get medical attention if irritation persists.

NOTE May produce temporary allergic side effects characterized by redness of eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reactions should consult a physician. Affected persons respond to treatments with antihistamines or steroid creams and/or systemic steroids.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

*Covered by U.S. Patent No. 3,290,353 and No. 3,331,735.

Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

VOLUNTARY PURCHASING GROUPS, INC.
Bonham, Texas

1982 JUN 11 1982

NET CONTENTS ONE POUND



ferti-lome



Disease Control for Flower and Vegetable Gardens

ACCEPTED
JUN 11 1982
General Insecticide,
Rodenticide Act,
registered under
EPA Reg. No. 7401-331

- For Effective, Broad-Spectrum Control of Diseases on Flowers, Vegetables & Lawns.
- See Side of Front Panel For Plant Diseases Controlled.

ACTIVE INGREDIENT

Chlorothalonil

(tetrachloroisophthalonitrile)* 5.0%

INERT INGREDIENTS 95.5%

KEEP OUT OF REACH OF CHILDREN

CAUTION
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

CONTROLS

- Black Spot
- Blossom Blight
- Powdery Mildew
- Ascochyta
- Ray Blight
- Downy Mildew
- Rust
- Gray Mold
- Botrytis Blight
- Leaf Spot
- Brown Patch
- Alternaria
- Leaf Spot
- Melting-Out
- Early & Late Blight
- Helminthosporium
- Leaf Blight
- Rhizoctonia
- Fruit Rot
- Copper Spot
- Target Spot
- Anthracnose
- Gummy Stem Blight
- Leaf Blight
- Scab
- Cercospora
- Leaf Spot
- Dollar Spot

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. WHAT IS THIS PACKAGE FOR?
This broad spectrum fungicide was especially formulated to effectively control damaging plant diseases on flowers, vegetables and lawns. See side of front panel for a list of diseases controlled.

2. HOW DO YOU USE IT?
Use sufficient dust to obtain full coverage on upper and lower leaf surfaces, stems and above ground parts. Do not graze treated areas or feed treated plant parts to livestock.

FLOWERS Apply at the first sign of disease and repeat at 7 to 14 day intervals. **ROSES** For control of Black Spot and Blossom Blight. **CHRYSANTHEMUM** For control of Ascochyta Ray Blight & Gray Mold. **GERANIUMS** Botrytis Blight. **IRIS** Botrytis Blight & Leaf Spot. **ZINNIA** Powdery Mildew.

LAWNS For control of Brown Patch, Dollar Spot, Leaf Spot (Blight Going out, Fading out and Melting out), Gray Leaf Spot, Stem Rust on Bluegrass and Copper Spot, apply at the first sign of disease and repeat at 7-14 day intervals. Thoroughly dust area affected and for at least 6 inches around affected area. Do not mow or water treated areas for at least 24 hours after application. For severe disease conditions shorten application intervals.

VEGETABLES Numbers in () indicate days required between last application and harvest. **BEANS (Snap) (7)** To control Rust. Apply during early bloom stage or when disease first threatens. Apply weekly or as necessary to maintain control. Do not graze treated areas or feed treated plant parts to livestock. **TOMATO (O)** To control Early & Late Blight, Gray Leaf Spot & Gray Leaf Mold. Apply when disease threatens and repeat at 7 to 10 day intervals. Under severe conditions shorten application intervals. **CORN (Sweet) (14)** To control Helminthosporium Leaf Blight. Apply when conditions favor disease development and repeat at 4 to 7 day intervals or as necessary to maintain control. Do not apply to sweet corn for processing. Do not graze treated fields, ensile treated corn, or use as feed or forage. **CUCUMBERS (O)** To control Target Spot (Florida only), Anthracnose & Downy Mildew. Apply when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals. Under severe conditions shorten application intervals. **CABBAGE CAULIFLOWER BROCCOLI BRUSSELS SPROUTS (O)** To control Alternaria Leaf Spot (South eastern states only) & Downy Mildew. Apply after transplant or set in field or shortly after emergence of field seedlings. Under severe conditions favor disease development. Repeat at 7 to 10 day intervals. **CARROT (O)** To control Early & Late Blight. Start applications when disease threatens and repeat at 7-10 day intervals as necessary to maintain control. **WATERMELON SQUASH PUMPKIN** For control of Anthracnose & Downy Mildew. Apply when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals. Under severe conditions shorten application intervals.

PRECAUTIONARY
STATEMENT
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS

WARNING

CONTAINS PERMETHRIN DISTILLATE.

Use only with adequate ventilation. Do not breathe spray, mist or vapors.

Harmful or fatal if swallowed or absorbed through skin.

Causes skin and eye irritation. Do not get into eyes, on skin or on clothing.

Wear goggles or face shield and rubber gloves when handling and cleaning. Use protective clothing when handling freshly treated leather.

Wash thoroughly with soap and water after handling and before eating or drinking.

Remove contaminated clothing and wash before reuse.

Respiratory protection may be required.

Keep away from children, pets and livestock.

STATEMENT OF PRACTICAL
TREATMENT

If on skin, wash with plenty of soap and water. Get medical attention.

If swallowed, do not cause vomiting. Call a physician immediately.

If inhaled, get fresh air. If necessary, call a physician immediately.

If in eyes, flush with plenty of water. Get medical attention.

If on clothing, remove and wash before reuse.