

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful or fatal if swallowed, inhaled or absorbed through skin. Do not apply by hand. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of food and feed products. Do not store near food or feed products. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed—Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If Inhaled—Remove victim to fresh air. Apply artificial respiration if indicated.

If on Skin—Remove contaminated clothing and wash affected areas with soap and water.

If in Eyes—Wash eyes with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and other aquatic organisms. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

May kill honeybees in substantial numbers. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use when the value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until one week after application or to take other equally effective precautions.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Voluntary Purchasing Groups, Inc.
Bonham, Texas 75418

EPA Est. No. 7401-TX-1 EPA Reg. No. 7401-334

Controls

Black Spot

Powdery Mildew

Leaf Spot

Rust

Brown Canker

Brown Rot

Scab

Septoria Leaf Spot

Sooty Blotch

Flyspeck

Kills

Mosquitoes

Lice

Scale

Elm Leaf Beetles

Boxwood Leaf Miners

Mealybugs

Ants

Strawberry Weevils

Meadow Spittlebugs

Pickleworms

Tarnished Plant Bugs

Velvet Bean Caterpillars

Leaf Rollers

Hi-Yield®

SEVIN & SULFUR DUST

ACCEPTED

JUN 5 1985

7401-334

- Kills Insects on Vegetables, Flowers, Shrubs, Trees and Lawns.
- Controls Diseases on Vegetables, Flowers, Shrubs, Trees and Lawns.
- Controls Black Spot, Powdery Mildew, Leaf Spot, Rust and Brown Canker on Roses

Active Ingredients:
Carbaryl (1-naphthyl N-methylcarbamate) . . . 2.0%; Sulfur . . . 2.0%; Inert Ingredients: . . . 96.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel For Additional Cautions

Kills

Squash Bugs

Cabbage Worms

Cutworms

Tomato

Fruitworms

Colorado Potato Beetles

Armyworms

Japanese Beetles

Bean Leaf Beetles

Squash Vine Borer

Cucumber Beetle

Leathoppers

Corn Earworm

Corn Borer

Melonworms

Hornworms

Harlequin Bugs

Flea Beetles

Stink Bugs

Lace Bugs

Strawberry Leaf Rollers

Thrips

Aphids

Bagworms

Chinch Bugs

Cankerworms

Fleas

Ticks

Chiggers

Tent Caterpillar

Sod Webworm

Oak Leaf Miner

Millipedes

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Hi-Yield Sevin & Sulfur Dust is ideal for use in killing insects and controlling diseases on vegetables, flowers, shrubs and lawns. This insecticide, fungicide combination will provide you with an easier and more economical means of eliminating a wide variety of damaging insects and diseases.

Use Hi-Yield Sevin & Sulfur Dust when insects or disease first appear. For best results, apply thoroughly to upper and lower leaf surfaces. If necessary repeat every 7-10 days. Do not dust too heavily or crop injury may result.

VEGETABLES: Apply at the rate of 1½-2 lbs. per 1,000 sq. ft. Intervals in number of days between last application of dust and harvest of product that must be observed are indicated in () after each crop heading.

Subsequent planting of food and feed crops in treated soil must be made with crops listed on this or other carbaryl labels.

Beans (0)	Carrots (3)
Tomatoes (0)	Beets (Roots 3, Tops 14)
Potatoes (0)	Brussels Sprouts (3)
Corn (7)	Radishes (3)
Cucumbers (0)	Turnips (Roots, 3, Tops 14)
Squash (0)	Collards (14)
Okra (0)	Mustard Greens (14)
Peppers (0)	
Spinach (14)	
Peas (0)	
Lettuce (Head 3, Leaf 14)	Melons (0)
Cabbage (3)	Strawberries (1)
Eggplant (0)	Grapes (0)
	Cauliflower (3)

FLOWERS, SHRUBS & TREES: Use a sufficient amount of dust to obtain a full coverage, being sure to cover upper and lower leaf surfaces. Repeat at 7 day intervals and after rains. Apply when new growth starts in spring or when diseases or insects appear.

LAWNS: To kill Chinch Bugs, Sod Webworms (Lawn Moths), Fall Armyworms, Cutworms, Ants, Fleas, Chiggers, Ticks, Mosquitoes, Earwigs, Leafhoppers, and Millipedes. Thoroughly dust infested areas and repeat as necessary.

NOTE: Do not use on Boston Ivy or Virginia Creeper.

STORAGE: Store in original container and place in a locked storage area.

DISPOSAL: Do not reuse container. Wrap and discard in trash.

PM 04

7401-334

6/5/85

PS 181