

PM 25

7401-336  
(Front of Bag)

4/6/87

183

Suggested Retail Price

Kill Weeds and Feed Your Lawn  
Coverage--Up to 2500 Sq. Ft.

**ACCEPTED**  
APR 6 - 1987  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 7401-336

ferti-lome®

St. Augustine  
Weed & Feed

Use on St. Augustine, Zoysia, Centipede,  
and Carpetgrass Lawns Only

Do not use on Bluegrass, Fescue, Bent, Bahia, Rye, Bermuda or Dichondra lawns.  
Controls Crabgrass, Sedge, Spotted Spurge and Paspalum when applied before  
these weeds germinate.

Fertilize lawns and kill the following listed weeds in one easy step!

- |                  |                   |                     |
|------------------|-------------------|---------------------|
| Dandelion        | Plantain          | Shepherdspurse      |
| Pennywort        | Lippia            | Virginia Pepperweed |
| Chickweed        | Black Medic       | Red Sorrel          |
| Sandbur          | Evening Primrose  | Stinging Nettle     |
| Henbit           | Rabbit Tobacco    | Wild Mustard        |
| Common Purslane  | Annual Sowthistle | Dock                |
| Burclover        | Blue Toadflax     | Rescuegrass         |
| Florida Betony   | Cranesbill        | Pigweed             |
| Florida Pusley   | Dollarweed        | Venus Lookingglass  |
| Creeping Charlie | Oxalis            |                     |


ACTIVE INGREDIENT:

Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine)	0.79%
Related Compounds	0.04%
INERT INGREDIENTS	99.17%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS



42.17

(Back of Bag)

ferti-lome(R)

St. Augustine  
Weed & Feed  
20-4-4

GUARANTEED ANALYSIS

Total Nitrogen(N)..... 20.0%  
 00.50% Ammonical Nitrogen  
 19.50% Urea Nitrogen  
 Available Phosphoric Acid(P<sub>2</sub>O<sub>5</sub>)..... 4.0%  
 Soluble Potash(K<sub>2</sub>O)..... 4.0%  
 Derived from - Primary Plant Nutrient Sources: Ammonium Phosphate,  
 Urea, Muriate of Potash.  
 Chlorine, not more than.... 4.0%  
 EPA Reg. No. 7401-336  
 VOLUNTARY PURCHASING GROUPS, INC.

EPA Est. No. 7401-TX-1  
BONHAM, TEXAS 75418

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash thoroughly after handling or using. Keep children and pets off treated areas until material is washed into the soil.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Atrazine can travel (seep or leach) through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand or loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

NET WT. 16 LBS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. WHAT IS THIS PACKAGE FOR?

This product is especially formulated to promote deep green, thick turf growth and eliminate unsightly weeds.

2. HOW DO YOU USE IT?

To prevent growth of Crabgrass, Sedge, Spotted Spurge, and Paspalum, apply in January, February or March before these weeds germinate. These weeds will not be controlled if they are present in the lawn at the time of application. Apply in Fall to prevent winter weeds such as Henbit, Chickweed, Rabbit Tobacco and Evening primrose.

To kill other listed weeds, apply anytime weeds are present and actively growing. Repeat in 2 months if needed.

If soil is dry, water thoroughly the day before treatment. Treat entire lawn evenly. Immediately after application, water to wash particles off grass blades.

NOTE: Don't apply under the branch spread (rootzone) area of trees and shrubs in compact, slowly draining soils (Particularly in the Dallas/Ft. Worth area).

Don't use on Centipede or Carpetgrass lawns during the period of winter dormancy (approximately October 15 to March 15), except in Florida. Don't make applications

to St. Augustine, Zoysia, Centipede or Carpetgrass lawns containing other grasses. Don't apply to newly seeded Zoysia, Centipede or Carpetgrass lawns until after the new grass has been mowed at least twice. Don't plant sprigs for one month, or sow new grass seed for six months following application of this product. Don't apply on bedding plants or ornamental trees and shrubs.

ONE BAG COVERS  
2500 Sq. Ft.

<u>SPREADER</u>	<u>SPREADER SETTING*</u>
ferti-lome CYCLONE/BROADCAST**	4 1/2
ferti-lome DELUXE/DROP	9 3/4
Central(Sears, Wards, J.C. Penny)	9 3/4
Scotts/DROP	10

\* These settings can be used only as a guide; they are not absolute. The speed at which the applicator walks, spreader condition and turf texture will alter the rate of fertilizer application.

\*\*Due to the wide swath, use broadcast lawn spreaders in open areas only: avoid trees, shrubs, and flowers. To prevent over application and possible injury, spreader passes should be spaced an adequate distance apart to assure uniform coverage.

3. HERE ARE THE RESULTS YOU SHOULD EXPECT!

Green, luxurious weed free lawns can be expected with this effective, easy-to-use product.

STORAGE: Store in original container and place in a locked storage area.

DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse bag. Discard bag in trash.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

Voluntary Purchasing Groups, Inc.  
Bonham, Texas

