PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION: May be harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Keep children and pets off treated areas until this material is washed into soil and area is dry. Do not apply by hand. Avoid contamination of food or feedstuffs.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED—Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Never give anything by mouth to an unconscious person.

IF INHALED—Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN-Remove contaminated clothing and wash affected areas with soap and water. IF IN EYES-Flush eyes with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically, Atropine is antidotal. Give by injection only.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment of disposal of wastes. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

*DURSBAN-trademark of Dow Chemical Co., U.S. Patent No. 3,244,586

Voluntary Purchasing Groups, Inc. Bonham, Texas

EPA Est. No. 7401-TX-1 EPA Reg. No. 7401-347

Net Contents One Pound



DURSBAN*

Garden

JUL 1 5 1987 Dust

7401-347.

- The Easy Way to Kill Destructive Insects.
- Kills Destructive Insects on Flowers, Shrubs, Trees, Vines, Evergreens & Vegetables.
- Helps Prevent Infestation of Buildings by Ants, Crickets, Fleas, Brown Dog Ticks, Millipedes, Earwigs, Sowbugs and Many Others.

ACTIVE INGREDIENTS:

Chlorpyrifos 10, 0-diethyl 0-(3, 5, 6-trichloro-	
2-pyridyl) phosphorothioate	0.5%
INERT INGREDIENTS:	99.5%

KEEP OUT OF REACH OF CHILDREN CAUTION SEE SACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Hi-Yield DURSBAN* GARDEN DUST is a new, easy to use product for effectively killing destructive insects on flowers, shrubs, vegetables and around buildings.

FLOWERS, SHRUBS, TREES, VINES & EVER-GREENS: To control Aphids, Bagworms, Cutworms, Grasshoppers, Tent Ceterpillars, Fall Webworms, Mealybugs, Mites, Spittlebugs, Thrips, Whiteflies, Orangestriped Qakworm and Yellownecked Caterpillar. Uniformly Just upper and Jower leaf surfaces and infested limb areas. Treat when pest appear and repeat at 7-10 day intervals. if needed.

VEGETABLES: SWEET CORN, POPCORN & FIELD CORN: To control Armyworms, Billbugs, Corn rootworms, Cutworms, Grasshoppers, Lesser cornstalk borer and Wireworms. For Corn rootworms, Cutworms, and Wireworm treatment, uniformly apply 1 lb. of dust per 300 sq. ft. of soil surface. BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, RADISHES & RUTABGAS: To control Root Maggots. Uniformly apply 1% lbs. dust per 300 feet of row at planting time. Throughly work dust into top layer of soil. For transplanted crops, apply 1¼ lbs, per 300 feet of row to the soil at the base of the plant. For radishes and rutabagas do not make more than one application per season. Do not use rutabaga tops for food or feed purposes.

SWEET POTATOES: To control Flea beetles & Wireworms. Uniformly apply 1% lbs. per 300 sq. ft. of soil surface before planting. Work dust into soil to a depth of 4 to 6 inches. Plant as soon as possible after treatment. Do not make more than one application per season.

OUTSIDE BUILDINGS: To help prevent infestation of buildings by ants, crickets, earwigs, millipedes, fleas, Brown dog ticks and sowbugs (pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to buildings, where pests are active and may find entrance. Uniformly apply at a rate of 1 lb. per 200 sq. ft. After application, lightly water the treated band.

DISPOSAL.

Do not reuse empty container. Wrap container and put in trash collection.

7/15/82