

ferti-lome (R)

Triple-Action

insecticide-miticide-fungicide

For Shrubs, Lawns, Flowers, Vegetables,
Fruit & Shade Trees.

Kills Leaf Chewing & Sap Sucking Insects

Controls Diseases

2 Tablespoons Make 1 Gallon of Spray

ACTIVE INGREDIENTS:

Methoxychlor: 1,1,1-trichloro-2,2-bis(p-methoxyphenyl)ethane..... 9.60%

Malathion: 0,0-dimethyl phosphorodithioate of diethyl mercaptosuccinate.... 8.07%

Copper Oleate**..... 3.00%

INERT INGREDIENTS*..... 79.33%


* Contains Xylene Range Aromatic Solvent

** Copper Expressed as metallic 0.27%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS



Back Panel:

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

1. WHAT IS THIS PACKAGE FOR?

This product is especially formulated for all-purpose garden and lawn uses. This Insecticide-miticide-fungicide combination controls leaf chewing and sap sucking insects and controls listed diseases on shrubs, lawns, vegetables, flowers, fruit and shade trees.

2. HOW DO YOU USE IT?

Mix 2 tablespoons (loz.) in one gallon of water. Apply to thoroughly wet upper and lower leaf surfaces and stems. Spray when insects or their damage first appear or when disease appears. Repeat at 7-10 day intervals as needed.

VEGETABLES- Do not apply within 7 days of harvest.

CROP	INSECTS CONTROLLED	DISEASE CONTROLLED
Beans, Peas	Aphids, Cowpea Curculio, Mexican Bean Beetle, Lygus Bug, Cucumber Beetle, Fall Armyworm, Bean Leaf Beetle	Powdery Mildew
Tomatoes	Aphids, Colorado Potato Beetle, Mites, Fall Armyworm, Flea Beetle, Mealybug, Fruitworm, Leafminer	Powdery Mildew
Cucumber, Melon	Aphids, Cucumber Beetle, Pickleworm, Spider Mites, Fall Armyworm, Squase Bug, Flea Beetle.	Powdery Mildew
Cabbage, Brussel Sprouts, Broccoli, Mustard, Turnips	Cabbageworms, Aphids, Leafhoppers, Flea Beetle, Japanese Beetles, Fall Armyworms	Powdery Mildew

FRUIT TREES-May be applied up to 7 days of harvest on apples, pears, plums and 21 days of harvest on peaches. NOTE: The following application schedule for orchard pests: Codling Moth, Apple Maggot, and Plum Curculio must be followed. Apply at petal fall and every 10-14 days thereafter until control is achieved.

CROP	INSECTS CONTROLLED	DISEASE CONTROLLED
Apples, Pears	Aphids, Mites, Apple Maggots, Codling Moth, Japanese Beetles, Plum Curculio, Tent Caterpillar, Forbes Scale	Powdery Mildew
Peaches, Plums	Aphids, Mites, Fruitworms, Fruit Fly, Japanese Beetles, Plum Curculio, Rose Chafer, Tent Caterpillar, Canderworm, Terripan & Cottony Cushion Scale	Leaf Blight, Leaf Curl, Powdery Mildew and Shothole
Flowers, Shrubs, Shade Trees	Aphids, Bagworms, Thrips, Leafhoppers, Japanese Beetles, Mites, Leafminers, Whiteflies, Mealybugs, Tent Caterpillars, Tarnished Plant Bug Scale, Rose Slugs, Rose Chafers, Cankerworms	Blackspot (roses) Leaf Spots, Rusts & Powdery Mildew
Lawns	Japanese beetle (Larvae), Chinch Bugs, Mosquitoes, Ants, Ground Pearls	Make a full coverage application to lawn areas. Immediately wash into soil after treatment.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing of spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed or feedstuffs. Keep children and pets of treated area until this material has dried completely. Food utensils such as teaspoons or measuring cups should not be used for food purposes after use with pesticides.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED- Call a physician or Poison Control Center immediately. Do not induce vomiting because of aspiration hazard.
IF INHALED- Remove victim to fresh air. Apply artificial respiration if indicated.
IF ON SKIN- Remove contaminated clothing. Wash affected areas with soap and water.
IF IN EYES- Flush eyes with plenty of water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated areas. Do not apply when bees are actively visiting the crop. Cover crop, or weed blooming in the treatment. Application should be timed to provide maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARDS: Do not use, pour, spill or store near heat or open flame.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

VOLUNTARY PURCHASING GROUPS, INC.
EPA EST. NO. 7401-TX-1

BONHAM, TEXAS 75418
EPA REG. NO. 7401-369

NET CONTENTS _____