

7401-378 PM 23 1063
(Front Panel)

Hi-Yield(R)

5%
CAPTAN
Dust

ACCEPTED
Per sell Cert
OCT 13 1987
Under the Federal Insecticide, Fungicide, and Rodenticide Act
7401-378

*Controls diseases on vegetables, shrubs, fruits, turf, and lawns.

*1 lb. covers up to 900 sq.ft.


Active Ingredients:

*Captan.....	4.889%
Related Derivatives.....	0.111%
Inert Ingredients:.....	95.000%

*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

Keep Out Of Reach Of Children
DANGER

See back panel for additional precautionary statements



DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Hi-Yield 5% CAPTAN DUST provides an effective and easy method for controlling diseases on Vegetables, Shrubs, Flowers, Lawns and Turf.

To apply, thoroughly dust upper and lower surfaces of plant foliage. Frequent light dustings are more effective than single heavy dustings. Where soil applications are required, follow directions listed after the crop heading. Begin applications before disease appears or at first sign of disease. Where lower and higher rates of application are shown, use the higher rate when disease conditions are severe.

VEGETABLES

TOMATOES: Early & Late Blight, Anthracnose, Septoria Leaf Spot, Gray Leaf Spot (*Stemphylium*)
Young Plants - Apply $\frac{1}{4}$ - $\frac{1}{3}$ lb. to 450 sq.ft. Mature Plants - Increase to $\frac{1}{2}$ -1 lb. per 450 sq.ft.
Repeat at 7-10 day intervals.

CANTALOUPEs, CUCUMBERS: Downy Mildew, Angular Leaf Spot, Anthracnose - Apply 1 lb. to 600 sq.ft.
Continue applications at weekly intervals as needed.

POTATOES (Seed Pieces): Damping-Off, Seed Rot-- Dust 1 lb. per bushel cut seed pieces.

Seed pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored for 2 to 3 days in open crates before bagging. Do not use treated seed for food or feed purposes.

STRAWBERRIES: Botrytis Rot (Gray Mold), Leaf Spots: Apply 1 lb. to 600 sq.ft. Begin at new growth. Repeat at 7 day intervals through picking and after harvest.

FLOWERS & SHRUBS

ROSES: Black Spot, Botrytis Blossom Blight. **CARNATIONS:** Alternaria Leaf Spot, Rust.

CHRYSANTHEMUM: Botrytis Flower Blight, Septoria Leaf Spot. **AZALEAS:** Petal Blight - Apply 1 lb to 800 sq.ft. Repeat at 7-14 day intervals and more frequently during frequent rains and heavy dews. For Azaleas, make applications only through bloom.

CAMELLIAS: Petal Blight - Apply $\frac{1}{2}$ lb. to 800 sq.ft. Repeat at 7-10 day intervals. Apply only through bloom.

LAWNS, TURF: Brown Patch, Damping-Off, Leaf Spots, Copper Spot, Melting Out, Seedling Blights - Apply 1 lb. to 350 sq.ft. Repeat at 7-10 day intervals throughout season. Do not graze treated areas or feed treated clippings to livestock.

ORANGE: Store in original container and place in a locked storage area.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Discard container in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse. Avoid contact with skin or clothing. Wear chemical resistant gloves when using indoors or outdoors. When using wear long pants and long-sleeved shirt and apply with the wind to your back. Wash non-disposable gloves thoroughly with soap and water before removing.

STATEMENT OF PRACTICAL TREATMENT

If In Eyes - Flush eyes with plenty of water. Call a physician.

If Swallowed- Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

If On Skin - Remove contaminated clothing. Wash affected areas with soap and water.

If Inhaled - Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

(Back Panel Cont.)

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this material to conform to the chemical description on the label and is suited for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the user or buyer.

Voluntary Purchasing Groups, Inc.

Bonham, Texas 75418

EPA Est. No. 7401-TX-1

EPA Reg. No. 7401-378

NET CONTENTS _____ LBS.

