

P.M.R. 7401-386

UN-VARNISHED CAP

1991

STORAGE AND DISPOSAL
 Storage: Store in original container and place in a locked storage area.
 Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.
 Container Disposal: Do not reuse container. Rinse thoroughly before discarding in trash.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed or foodstuffs. Keep children and pets off treated areas until this material has dried completely.

STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Never induce vomiting or give anything by mouth to an unconscious person.
IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.
IF ON SKIN: Remove contaminated clothing. Wash affected areas with soap and water.
IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.
NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidote.

ENVIRONMENTAL HAZARDS
 This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatments or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.
 Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this product to conform to the chemical description on the label and for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the user or buyer.
 *SEVIN - Registered Trademark of Union Carbide Corporation for Carbaryl insecticide.
 Voluntary Purchasing Groups, Inc.
 Bonham, Texas 75418
 EPA Est. No. 7401-TX-1 EPA Reg. No. 7401-386
 14M-2-74H Net Contents One Quart

ferti-lome

JUN 5 1985

**Gypsy Moth
 Japanese Beetle
 & Pine Tip Moth Killer** \$8.98

- Kills Gypsy Moths, Japanese Beetles & Pine Tip Moths on Roses, Flowers, Shrubs & Shade Trees.
- Kills other listed insects on Fruit Trees & Vegetables.
- 2 Tablespoons make 1 Gallon of Spray.
- 1 Quart makes 32 Gallons of Spray.



Keep out of reach of children
CAUTION
 See back panel for additional precautions

Active Ingredient Carbaryl (1 naphthyl methylcarbamate) 13.5%
 Inert Ingredients 86.5%

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

WHAT IS THIS PACKAGE FOR?
 This product provides an easy and effective method for killing Gypsy Moths, Japanese Beetles, Pine Tip Moths and many other destructive insects on Roses, Flowers, Shrubs, Shade Trees, Fruit Trees and Vegetables.

2. HOW DO YOU USE IT?
 Shake well before using. Mix 2 tablespoonful (1 oz.) per gallon of water or 1 pint in 16 gallons of water. Apply to wet all exposed plant parts. May be used in all types of sprayers.

ROSES, FLOWERS, SHRUBS, SHADE TREES: For Japanese Beetles, Gypsy Moths, Pine Tip Moths, Aphids, Bagworms, Elm Leaf Beetle, certain Leaf Miners, Rose Slugs, Periodical Cicada (17 year locust), Scale Insects, Tort Caterpillars, Blister Beetles, Leafrollers, Exposed Thrips - Spray when insects or their damage appear. Repeat weekly or as necessary. Do not use on Boston Ivy or Virginia Creeper. Injury to tender foliage may occur if plants are wet when treated or humidity is high.

FRUIT TREES (Apples, Pears & Peaches): For Apple Maggots, Plum Curculio, Apple Aphids, Leafrollers, Peach Tree Borers, Oriental Fruit Moths, Codling Moths - Make one application between bud break and bloom, a second application at petal fall, and then re-apply at 3-4 week intervals for 3-4 months as necessary. Apple Maggot - For best control, apply at 7-10 day intervals. Rosy Apple Aphid - Apply before leaves are curled. **NOTE:** To avoid foliage injury on McIntosh apples and possible fruit thinning on all varieties of apples, delay use until at least 30 days after bloom. Do not apply within 1 day of harvest.

VEGETABLES (Tomatoes, Eggplant, Peppers, Cucumbers, Summer Squash): Pickleworms, Fruitworms, Melonworms, Hornworms, Fall Armyworms, Corn Borers - Apply when insects first appear. Repeat at 7 to 10 day intervals. May be applied up to day of harvest. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

3. HERE ARE THE RESULTS YOU MAY EXPECT!
 When used as directed, this product will kill Gypsy Moths, Japanese Beetles, Pine Tip Moths and other destructive insects and thus help provide for healthier, more attractive Roses, Flowers, Shrubs, Shade Trees, Fruit Trees and Vegetables.