10/05/2000 OCT -5 2000

1/5

Marla K. Jackson Brazos Associates, Inc. P.O. Box 703524 Dallas, Texas 75370-3524

Dear Ms. Jackson:

SUBJECT:

8-Month Response to Oryzalin RED ferti-lome Weed and Feed Special EPA Registration No. 7401-415
Your Application Dated April 15, 2000

The revised labeling for the subject product, referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable with the following provisions:

On page 4, under "Directions for Use" there is a typographical error in the first question. Replace the word "if" with "is", so that it reads: "What is this package for?"

A stamped copy of the labeling is enclosed for your records. Please submit one copy of your final printed labeling incorporating this change before you release the product for shipment.

EPA has determined that you have fulfilled the labeling and product specific data requirements of the oryzalin RED. However, we cannot reregister this product at this time for the following reason:

The subject product is registered for us on turfgrass, a use for which the Agency does not have sufficient information to make a reregistration eligibility decision. Once the required data (foliar dislodgeable residues/guideline 132-la; soil dislodgeable residues/guideline 132-lb; estimation of dermal exposure/guideline 133-3; and estimation of inhalation exposure/guideline 133-4) have been reviewed by the Agency, EPA will make a reregistrationi eligibility decision for the turfgrass use.

Sincerely yours,

Joanne I. Miller Product Manager (23) Herbicide Branch Registration Division (7505C)

Enclosure

					 		
				URRENCES			
SYMBOL .	7505C						
SURNAME -	Walson	************	•	}	ĺ	ľ	
DATE	18/5/00						

Kill Weeds And Feeds Your Home Lawn

ferti-lome®

WEED & FEED SPECIAL

• For Use On Established Warm Season Turf Grasses

Bermuda

Bahia

St. Augustine

Centipede

Zoysia

 Developed For Winter And Spring Pre-Emergence Weed Control And Lawn Fertilization Programs.

ACTIVE INGREDIENTS:

Benefin: N-butyl-N-ethyl- α , α , α -trifluoro-2,6-dinitro-	
p-toluidine	1.0%
oryzalin: 3.5-dinitro-N ⁴ , N ⁴ -dipropylsulfanilamide	1.0%
INERT INGREDIENTS:	98.0 <u>%</u>
TOTAL:1	00.0%

Contains 0.172 lbs. active ingredients per 8.6 lb. bag Coverage ---- Up to 2,500 sq. ft.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Precautionary Statements

NET WEIGHT 8.6 LBS.

• ACCEPTED
with COMMENTS
IN EPA Letter Dated

OCT - 5 2000-

Under the Federal Inscolicide. Fundicide, and Rodemicide Act an amended, for the pesticide registered under EPA Reg. No.

[BACK PANEL]

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Keep all persons, children and pets out of the treated area until dusts have settled.

Personal Protective Equipement (PPE): Wear eye protection, long-sleeved shirt, long pants, shoes, socks, and chemical resistant gloves when using this product. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Users should wash hands after using and before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Wash the outside of gloves before removing.

First Aid

If In Eyes: Hold eyelids open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

If On Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If Inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL		••••
Storage: Store in original container only and preferably in a locked are	a away from	••••
children or pets.		
Disposal: Securely wrap original container in several layers of newspap	per and discard	l in
trash.	: ::	::
	****	••••
		•••••
•		****
		• • • • •

[BACK PANEL]

20 - 4 - 4

GUARANTEED ANALYSIS

Total Nitrogen (N)	20%
00.50% Ammoniacal Nitrogen	
19.50% Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	4%
Derived from-Primary Plant Nutrient Sources: Amn Urea, Muriate of Potash.	nonium Phosphate,
Chlorine, not more than	4.0%

DIRECTIONS FOR USE.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. WHAT IF THIS PACKAGE FOR?

This product is formulated to pre-emergently eliminate unsightly weeds and feed your lawn, all in one application.

2. HOW DO YOU USE IT?

Apply ferti-lome® Weed & Feed Special approximately 1 to 2 weeks prior to germination of annual grasses and weeds.

To Control Crabgrass, Foxtail and Barnyardgrasses, apply in January, February or March.

<u>Goosegrass</u> normally germinates 4 to 6 weeks later than Crabgrass and other summer grass species, therefore, applications made for goosegrass should be timed accordingly. Also, refer to "NOTE" below.

To Control <u>Poa annua</u>, (annual bluegrass), make application in early fall (September or October).

Where germination of annual grasses and weeds continue for extended periods of time, a second application can be made 3 months after initial treatment.

NOTE: The minimum allowable time between application is 3-months. Do not apply more than 34 pounds of this product per year to cover 2,500 sq.ft. Do not apply in spring to turfgrass planted the previous fall. Do not apply to new sprigged areas of Bermudagrass, St. Augustine or Centipede until turf is well established.

[BACK PANEL]

Using the spreader settings listed below, apply one 8.6 lb. bag per 2,500 sq. ft. of lawn area. A uniform coverage is important to assure complete weed control. ALWAYS WATER THOROUGHLY AFTER APPLICATION.

ONE 8.6 LB. BAG COVERS 2,500 SQ. FT.

<u>SPREADER</u>	*SETTING
ferti-lome® Cyclone/Broadcast	†
ferti-lome® Deluxe	†
Central (Sears, Wards, J.C.Penney)	†
Scotts	†

†[Spreader settings listed on label will be established based on actual application test of each spreader brand and will not exceed a rate of 1.5 lbs. a.i./acre]

*These settings can be used only as a guide, they are not absolute. The speed at which the applicator walks, spreader condition and turf texture will alter rate of fertilizer application.

NOTE: Do not apply in spring to turfgrass planted the previous fall. Do not apply to newly sprigged areas of Bermudagrass, St. Augustine or Centipede until turf is well established.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

Effective, economical weed control and a thick, green lawn may be expected with use of this product.

NOTE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this product to conform to the chemical description on the label and for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the user or buyer.

·	•••
VOLUNTARY PURCHASING GROUPS, INC.	
P.O. Box 460	•
BOHNAM, TEXAS 75418	••••
••••	• • • •
EPA Registration No. 7401-415	•
EPA Establishment No. 7401-TX-1	••••
	••••
Page 4	••••