1401-452

05-15-2009



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Leslie Oliver Agent for Voluntary Purchasing Groups, Inc. MANDAVA Associates, LLC 6860 N. Dallas Parkway, Suite 200 Plano, TX 75034

MAY 1 5 2009

Subject: Notification(s) for Label Revisions under PRN 2008-1 and 2007-4

Dear Ms. Oliver:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notices (PRN) 2008-1 and 2007-4 dated April 30, 2009 for:

EPA Registration 7401-452Hi-Yield Kill-A-Bug II Dust Containing
Deltamthrin

The Registration Division (RD) has conducted a review of the request(s) for applicability under PRN 2008-1 and 2007-4 and finds that the label changes requested fall within the scope of PRN 2008-1 and 2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identify the batch of the pesticide distributed and sold be placed on <u>nonrefillable</u> containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Nicole Williams of my staff at 703-308-5551.

Sincerely,

Linda Arrington Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

2/11

| lease read instructions on | reverse before completing form. | F | .bevorgqA mic | | | OPP Identifier Number |
|--|--|---|---|---|---|---|
| ₽EPA | o Agency | ency Registration Amendment × Other | | | | |
| | Application | n for Pesticide | - Section | I | | |
| . Company/Product Numb /401-452 | er | 2. EPA Pro George L | Suct Manager aRocca | | | posed Classification |
| . Company/Product (Name Hi-Yield Kill-A-Bug II Du | a) Ist Containing Deltamthrin | PM# 13 | | | | |
| Voluntary Purchasing (230 N. FM 87 Bonham, Texas 75418 | pplicant <i>(Include ZIP Code)</i> Groups, Inc. <i>is is a new address</i> | | n. No | ilar or identi N | cal in con | FIFRA Section 3(c)(3) position and labeling ATION 2009 |
| | | Section - II | | | | |
| Amendment - Exple Resubmission in re Notification - Explei | sponse to Agency letter dated | [] [•] | nal printed label gency lettar dat Ae Too" Applica ther - Explain ba | ation. | to | |
| I further understand that i | ormula for this product. I understand th f the amended label is not consistent wit A and I may be subject to enforcement act | h the requirements of | 40 CFR 156.10, | 156.140, 156.1 | 44, 156.146 | |
| 1. Material This Product V | Vill Be Packaged In: | | | • | | |
| Child-Resistant Packaging Yes* X No * Certification must be submitted | Unit Packaging Yes No If "Yes" No. per Unit Packaging wgt. container | Water Soluble Pac Yes X No If "Yes" Package wgt | Kaging No. per container | 2. Type of | Container Motel Plastic Glass Peper Other (S | pecity) |
| 3. Location of Net Conten | te Information 4. Size(s) Reta Container 1 lb. | il Conteiner | 5. La | | | ns panying product |
| 6. Manner in Which Label | | eph glued ed | Other | | | |
| | | Section - IV | | | | |
| 1 Contact Point (Comple | te items directly below for identification | n of individual to be a | ontacted, if nee | cessary, to pr | ocess this | application.) |
| | | Title | | | | No. (Include Area Code) |
| Name Madhu Mandava | | Agent for Voluntar | Y Purchasing G | roups, Inc. | 97 | 2-265-7924 |
| Name Madhu Mandava I certify that the sta | Certifical stements I have made on this form and any knowingly felse or misloading state | Agent for Voluntar tion all attachments there | ito are true, acc | curate and co | mplete. | 2-265-7924 8. Date Application Received (Stamped) |
| Name Madhu Mandava I certity that the stu i acknowledge that both under applicat | Certificat atoments I have made on this form and eny knowingly false or misleeding state bis law. | Agent for Voluntar tion all attachments there | nto are true, acc able by fine or | curate and co imprisonment | mplete. : or | 6. Date Application Received |

EPA Form 8570-1 (Rev. 8-94) Previous editions are obsolete.

Yellow - Applicant Copy

MANDAVA ASSOCIATES, LLC

<u>CONSULTANTS IN SCIENCE, TECHNOLOGY AND REGULATORY AFFAIRS</u> 6860 N Dallas Parkway, Suite 200, Plano, TX 75034 Telephone: (972) 265-7924 / Fax: (972)-265-7942 / E-MAIL: Madhu@Mandava.com / www.Mandava.com

January 14, 2009

Mr. George LaRocca US EPA Office of Pesticide Programs Registration Division (7504P) Ariel Rios Building 1200 Pennsylvania Avenue, N.W. Washington, D.C. 20460

SUBJECT: EPA Registration Number: 7401-452; Hi-Yield Kill-A-Bug II Dust Containing Deltamethrin; Notification Per PR Notices 2008-1 and 2007-4

Dear Mr. LaRocca:

On behalf of Voluntary Purchasing Groups, Inc. (VPG), we are submitting notification for change in Storage and Disposal Statements per PR Notice 2007-4 and Environmental Hazards Statements per PR Notice 2008-1 for Hi-Yield Kill-A-Bug II dust Containing Deltamethrin (EPA Registration Number: 7401-452).

Enclosed with this submission, please find the following:

- 1) Application for Pesticide Registration Notification
- 2) Two amended labels clearly marked
- 3) One original label

Please update the agency records in consideration of this notification.

If you have any questions regarding this submission, please contact me at (972) 265-7925.

Sincerely,

Jeslie Oliver

......

Sceres - - - -

Leslie Oliver Agent for Voluntary Purchasing Groups, Inc.

[FRONT LABEL PANEL]

Hi-Yield® Kill-A-Bug® II Dust Containing Deltamethrin

- For Indoor/Outdoor Residential Use Only
- [For Household Use]
- [Household Ornamental Garden Insect Control]
- [Controls Fire Ants, Fleas, Ticks *Termites and More]
- [Non-Staining & Odorless]
- [Long-Lasting Residual Action]
- [Long-Lasting]
- [Kills Insects for up to 8 Months]
- [Does not Clump or Absorb Moisture]
- [For Use on Ornamentals]
- [Use on Roses, Flowers & Ornamentals]
- [Controls Many Different Insect Pests]
- [To Control Imported Fire Ants on Lawns and Other Residential Areas]
- [8 oz. Treats 15 Fire Ant Mounds]
- [1 lb. Treats 30 Fire Ant Mounds]
- [2 lbs. Treats 60 Fire Ant Mounds]
- [Fast, Effective Kill of Fire Ants]
- [Fast, Effective Kill of Stinging Ants]
- [Easy To Use]
- [Convenient]
- [Kills Fire Ants on Contact]

[]-Denotes Optional Statements

*For localized area control only

ACTIVE INGREDIENT:

| *Deltamethrin [(s)-alpha-cyano-3-phenoxybenzyl-(1R,3R)-3-(2-2- | |
|--|--------|
| Dibromovinyl)-2,2-dimethyl cyclopropanecarboxylate] | 0.05% |
| INERT INGREDIENTS: | 99.95% |
| TOTAL :1 | 00.00% |

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back/Side Panel for Additional Precautionary Statements

Net Weight_____

NOTIFICATION

MAY 1 5 2009

[Back/Side Label Panel]

| Have the produc going for treatm | FIRST AID It container or label with you when calling a poison control center or doctor, or ent. |
|-------------------------------------|--|
| If swallowed | Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If in eyes | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. |
| If on skin or clothing | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. |
| If inhaled | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to mouth, if possible. Call a poison control center or doctor for treatment advice. |

PRECAUTIONARY STATEMENTS

Hazards to Humans And Domestic Animals

CAUTION: May be harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read the entire Directions for Use, Precautions and Disclaimer of Warranties before using this product. Hi-Yield[®] Kill-A-Bug[®] II Dust Containing Deltamethrin is a ready-to use insecticide dust which exhibits effective knockdown and long-lasting control against pests listed when used according to label directions. Hi-Yield[®] Kill-A-Bug[®] II Dust Containing

Deltamethrin can be used for long-lasting pest control in and around homes and their immediate surroundings. Permitted areas of use include:

| *animal quarters | *mobile homes and |
|-----------------------------------|-----------------------|
| *apartments | recreational vehicles |
| *household greenhouses *houses | *ornamentals |

This product is intended for applications with a hand duster, from the shaker can, or by other suitable means, to hiding and runway areas and other places where insects are found. Food utensils such as a teaspoon or measuring cup should not be used for food after use with pesticides.

INDOOR USE

In living areas, make applications in such a manner to avoid deposits-on exposed surfaces or introducing the material into the air.

To apply insecticide directly into cracks and crevices, use a bulbous duster, or other suitable equipment. Apply lightly and uniformly to infested areas. Pay particular attention to cracks and crevices; service ducts, false floors and ceilings; wall void; around electrical and telephone fittings and equipment; around water and sewer pipes; under and behind cabinets; refrigerators and sinks; around windows and door frames; along baseboards; in attics and crawl spaces. Apply at the rate of 1 oz. of product per 125 square feet (or $\frac{1}{2}$ lb. per 1000 square feet). Repeat after 10 days only if needed to maintain control. Do not apply more than 2 times per year.

Ants, boxelder bugs, cockroaches (including both resistant and non-resistant strains), crickets, fleas, firebrats and silverfish: Apply Hi-Yield[®] Kill-A-Bug[®] II Dust Containing Deltamethrin thoroughly to all areas where these pests crawl and hide, especially in those areas described above. For Ants, apply to ant trails, around doors and windows and other places where ants enter premises.

Termites, Carpenter Ants and Carpenter Bees: For Localized Treatment of Wood Infesting Termites, Carpenter Ants, and Carpenter Bees; apply to voids to channels in damaged wooden members of a structure, to cracks, spaces or bearing joints between wooden members of a structure; or between such members and the foundation; and in locations vulnerable to termite attack. Drilling may be necessary prior to applying dust in these locations, but it must be recognized that such drilling and treating of timber in a particular location may not result in complete coverage of infested and vulnerable voids or galleries. When drilling, care should be taken to avoid drilling into electrical wire, plumbing, gas lines or other utilities. Re-seal holes following treatment. This dust can be applied in attics and crawl spaces to prevent entry of drywood and formosan termites at a rate of 1/2 lb. per 1000 square feet of floor area. In these treatments, all vents and blowers should be closed. Inspect at 6 months and one year to ensure that colony has been killed. Repeat after 10 days only if needed to maintain control. Do not apply more than 2 times per year.

The purpose of such applications of Hi-Yield[®] Kill-A-Bug[®] II Dust Containing Deltamethrin is to kill workers or winged forms which may be present in the treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or

foundation treatment but merely a supplement. For active termite infestation, get a professional

inspection.

Centipedes, ground beetles, millipedes, scorpions, spiders and sowbugs: Apply dust along and behind baseboards, to window and door frames, corners, pipes, storage locations, attics, crawl spaces, and other areas where these pests may enter or crawl.

Wasps and Bees: It is generally advisable to treat wasp and bee nests in the evening when insects are less active and have returned to the nest. Wear protective clothing if deemed necessary to avoid stings. Using hand or power duster (actisol or similar equipment) or other suitable means, with extension tubes if necessary, thoroughly dust nets, entrance and surrounding areas where insects alight. Nests in wall voids can be located by listening with your ear against the wall. Drill a hole in the area, blow dust in and reseal. For ground wasps, dust entrance and surrounding areas. For best results check nests carefully one or two days after treatment to ensure complete kill then remove and destroy nest to prevent emergence of newly -hatched insects. If removal is not feasible, retreat nest if necessary. Do not use this product to treat overhead nests.

Webbing Clothes Moths: Use Hi-Yield[®] Kill-A-Bug ® II Dust Containing Deltamethrin as a crack and crevice treatment in closets and other storage areas where these pests are found. This is an adjunct treatment and will not control moth larvae already on the clothes. DO NOT apply this product directly to clothing.

Carpet Beetles (Black, Furniture, Common and Varied): Dust where these insects are found, for example: on and under edges of floor coverings, around baseboards, under rugs and furniture, and in closets.

Bedbugs: Apply dust thoroughly to all areas where these pests are typically found including folds and tufts of mattresses, coils or springs, cracks and hollow posts of bedsteads, upholstery of chairs and sofas, picture frame moldings and _{all} cracks and crevices in room should be treated. Allow powder to remain in contact with surface for 4-6 hours. Thoroughly vacuum treated areas, dispose of vacuum bag and place clean linens on bed.

Stored Product Pests: To aid in control of exposed stages of:

cadelles cheese mites cigarette beetles confused flour beetles dark meal worms drugstore beetles grain mites • granary weevils lesser grain borers merchant grain beetles red flour beetles rice weevils rusty grain beetles saw-toothed grain beetles spider beetles tobacco moths yellow meal worms

In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Clean up wastes materials, dust, dirt, and other debris. Thoroughly treat floors, walls and other surfaces in storage and handling areas. Apply dust in cracks and crevices, behind and under cupboards, and other difficult to reach areas where insects may hide. DO NOT apply directly on food, utensils or other areas where food is prepared, processed, served, or stored.

ORNAMENTAL PLANT PESTS For the control of:

aphids beet armyworms blister beetles boxelder bugs certain leafminers (Boxwood, Holly Oak, Spruce Needle elm leaf beetle flea beetles Japanese beetles leafrollers orange tortrix periodical cicada plant bugs rose chafer rose midges rose slugs tent caterpillars thrips weevils Leafhoppers

Shrubs and Ornamentals: Apply when insects or their damage first appear. Repeat after 10 days only if needed to maintain control. Do not apply more than 2 times per year. Apply 8 ounces per 10000 square feet; or thoroughly apply to place a thin layer of dust on both upper and lower sides of leaves, stems and flowers.

Roses, Petunias, Chrysanthemums, Geraniums, Carnations and other related flowers: TO control many leaf and flower chewing insects, apply as soon as the insect appears. Repeat after 10 days only if needed to maintain control. Do not apply more than 2 times per year. Apply a thin layer of dust on both upper and lower sides of leaves, stems and flowers.

OUTDOOR USE

Perimeter Treatment: Use Hi-Yield[®] Kill-A-Bug[®] II Dust Containing Deltamethrin as a residual treatment for control of ants, bees, centipedes, cockroaches, crickets, firebrats, fleas, ground beetles, millipedes, scorpions, silverfish, slugs, sowbugs, spiders, ticks and wasps. Make a residual treatment around windows and doors, porches, screens, eaves, patios, garages, under stairways, in crawl spaces where pests are active and may find entrance. **Fleas and Ticks in Rodent Burrows:** To control fleas and ticks in and around rodent burrows around homes and other buildings, apply dust with a shaker can. Thoroughly apply dust to adequately cover mound and around the perimeter (18" from the outer edge). Be sure to treat abandoned rodent burrows which are suspected of containing fleas and ticks. Treat for a minimum of 5 days for light rodent populations, 10 days when populations are dense. Retreatment should only be made as necessary to maintain insect control.

HOME VEGETABLE GARDENS: Use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate available duster. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest comer of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.

| TARGET PESTS | REMARKS | Application Rate: Oz. of Dust/100ft ² | Max Application/ Growing Season | [Active Ingredient/Acre] (This column is alternate text) | [Packages/100ft ^{2]} (This column is alternate text) |
|---|--|--|--|--|--|
| Aphids, armyworm, cabbage looper, cribrate weevil, leafminers, lygus bug (plant bug), pale striped flea beetles, plume moth, probas bug | Do not apply more than recommended amount [0.14 lb ai/A (active ingredient/acre)] in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest. | 2.13 | 5 | 0.028 lb. | X Packages/100 ft ² |
| Cutworms, stink bugs, thrips | Do not apply more than recommended amount [0.112 lb ai/A (active ingredient/acre)] in one growing season. Allow 5 days between applications. Do not apply within one days of harvest. | 2.13 | - 4 | 0.028 lb. | X Packages/100 ft ² |
| Aphids, armyworms, chinch bugs, click beetle, corn earworm, corn rootworm (adult), cutworms, European corn borer, flea beetles, grasshoppers, Japanese beetle (adult), June beetle (adult), Masked chafer (adult), Southern corn borer, Southwestern corn borer, stalk borer, stink bugs, webworms | Do not apply more than recommended amount [0.45 lb ai/A] in one growing season. Do not apply within 24 hours of harvest. | 2.13 | 16 | 0.028 lb | X Packages/100 ft ² |
| Aphids, armyworms, chinch bugs, click beetle, corn earworm, corn rootworm (adult), cutworms, European corn borer, flea beetles, grasshöppers, Japanese beetle (adult), June beetle (adult), Masked chafer (adult), Southern corn borer, Southwestern corn borer, stalk borer, stink bugs, webworms | Do not apply more than recommended amount [0.095 lb ai/A] in one growing season. Do not apply within 21 hours of harvest. | 1.68 | 4 | 0.022 lb | X Packages/100 ft ² |
| Armyworm, cabbage looper, Colorado potato beetle, corn earworm, cucumber beetle, cutworms, diamondback moth (larvae), Flea beetle, grasshoppers, green peach aphid, leafhopper, leafminer, melonworm, pickleworm, stinkbugs, tarnished plant bug, tomato pinworm, vegetable weevil (adult), western plant bug, whitefly (adult). | Do not apply more than recommended amount [0.168 lb ai/A] in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest. | 2.13 | 6 | 0.028 lb | X Packages/100 ft ² |
| | Aphids, armyworm, cabbage looper, cribrate weevil, leafminers, lygus bug (plant bug), pale striped flea beetles, plume moth, probas bug Cutworms, stink bugs, thrips Aphids, armyworms, chinch bugs, click beetle, corn earworm, corn rootworm (adult), cutworms, European corn borer, flea beetles, grasshoppers, Japanese beetle (adult), June beetle (adult), Masked chafer (adult), Southern corn borer, stalk borer, stink bugs, webworms Aphids, armyworms, chinch bugs, click beetle, corn earworm, corn rootworm (adult), cutworms, European corn borer, stalk borer, stink bugs, webworms Aphids, armyworms, chinch bugs, click beetle, corn earworm, corn rootworm (adult), cutworms, European corn borer, flea beetles, grasshoppers, Japanese beetle (adult), June beetle (adult), Southern corn borer, Southwestern corn borer, Stalk borer, stink bugs, webworms Armyworm, cabbage looper, Colorado potato beetle, corn earworm, cucumber beetle, cutworms, diamondback moth (larvae), Flea beetle, grasshoppers, green peach aphid, leafhopper, leafminer, melonworm, pickleworn, stinkbugs, tarnished plant bug, tomato pinworn, vegetable weevil (adult), western plant bug, | Aphids, armyworm, cabbage looper, cribrate weevil, leafminers, lygus bug (plant bug), pale striped flea beetles, plume moth, probas bugDo not apply more than recommended amount [0.14 lb ai/A (active ingredient/acre] in one growing season. Allow 3 days between applications. Do not apply more than recommended amount [0.112 lb ai/A (active ingredient/acre] in one growing season. Allow 5 days between applications. Do not apply more than recommended amount [0.112 lb ai/A (active ingredient/acre] in one growing season. Allow 5 days between applications. Do not apply within one days of harvest.Aphids, armyworms, chinch bugs, click beetle, corn carworm, corn borer, flea beetles, grasshoppers, Japanese beetle (adult), Masked chafer (adult), Southern corn borer, Southwestern corn borer, Suthwestern corn borer, Suthwestern corn borer, Suthwestern corn borer, Suthwestern corn borer, Southwestern corn b | Aphids, armyworm, cabbage looper, cribrate weevil, leadininers, lygus bug (plant bug), pale striped flea beetles, plume moth, probas bugDo not apply more than recommended armount [0.14 lb ai/A (active ingredient/acre)] in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest.2.13Cutworms, stink bugs, thripsDo not apply within 3 days of harvest.2.13Cutworms, stink bugs, thripsDo not apply more than recommended armount [0.112 lb ai/A (active ingredient/acre)] in one growing season. Allow 5 days between applications. Do not apply within one days of harvest.2.13Aphids, armyworms, chinch bugs, click beetle, corn earworm, corn toorer, flea beetles, grasshoppers, Japanese beetle (adult), Masked chafer (adult), Southern corn borer, stalk borer, stink bugs, webwormsDo not apply more than recommended armount [0.45 lb ai/A] in one growing season. Do not apply wore than recommended armount [0.45 lb ai/A] in one growing season. Do not apply within 24 hours of harvest.1.68Aphids, armyworms, chinch bugs, click beetle, corn earworm, corn rootworm (adult), cutworms, European corn borer, Stalk borer, stink bugs, webwormsDo not apply more than recommended armount [0.168 lb ai/A] in one growing season. Do not apply within 21 hours of harvest.1.68Aphids, armyworms, chinch bugs, click beetle, corn earworm, corn rootworm (adult), scutworms, stinkbugs, tarnished plant bug, tornito pinworm, vegetable weevil adadity, leadhopper, leadininer, melonworm, pickleworm, stinkbugs, tarnished plant bug, tornato pinworm, vegetable weevil (adult), we | Aphids, armyworm, cabbage looper, cribrate weevil, leafminers, lygus bug (plant bug), paic striped flae bettes, plume moth, probas bugDo not apply more than recommended amount [0.14 lb ai/A (active ingredient/acre] in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest.2.135Cutworms, stink bugs, thripsDo not apply more than recommended amount [0.112 lb ai/A (active ingredient/acre]) in one growing season. Allow 3 days between applications. Do not apply within one days of harvest.2.134Aphids, armyworms, chinch bugs, click beetle, com carworm, corn borer, flae beetles, grasshoppers, Japanese beetle (adult), Southern corn borer, flae beetles, grasshoppers, Japanese beetle (adult), Southern corn borer, fae beetles, grasshoppers, Japanese | Rate: Oz. of cabbage looper, cribrate weevil, learminers, tygus taristed flae beetles, pluer moth, probas bugDo not apply more than recommended amount [0.14 lb at/A (active ingredient/acre] in one growing season. Allow3 days between applications. Do not apply within 3 days of harvest.2.1350.028 lb.Cutworms, stink bugs, thripsDo not apply more than recommended amount [0.112 lb at/A (active ingredient/acre]) in one growing season. Allow3 days between applications. Do not apply wore than recommended amount [0.112 lb at/A (active ingredient/acre]) in one growing season. Allow3 days between applications. Do not apply more than recommended amount [0.112 lb at/A (active in one growing season. Allow3 days between applications. Do not apply more than recommended amount [0.418 bat/A] in one growing season. Do not apply more than recommended amount [0.418 bat/A] in one growing season. Do not apply more than recommended amount [0.418 bat/A] in one growing season. Do not apply more than recommended amount [0.418 bat/A] in one growing season.Do not apply within 24 hours of harvest.1.6840.028 lbAphids, armyworms, chinch bugs, click beetle, datult/, Mack chafer (adult/), Southern com borer, flae beetles, grashopers, Japanese beetle (adult), June beetle (adult/), Southern com borer, flae beetle, cutworms, flamode harder, amout [0.168 lb at/A] in one growing season. Di not apply more than recommended amount [0.168 lb at/A] in one growing season. Di not apply more harvest.1.6840.028 lbArmite commended amount [0.168 lb at/A] in one growing season. Allow 3 days be |

10/11

| Fruiting Vegetables- Eggplant, groundcherry, pepinos, peppers, tomatillo, tomato | Aphids, armyworms, cabbage looper, Colorado Potato Beetle, cucumber beetle, cutworms, diamondback moth, European corn borer, flea beetles, imported cabbageworm, leafminers, pepper weevil, stink bugs, tarnished plant bug, thrips, tomato fruitworms, (corn earworm), tomato pinworm, variegated cutworm, whitefly | Do not apply more than recommended amount [0.168 lb ai/A] in one growing season. Allow 5 days between applications. Do not apply within 1 days of harvest. | 2.13 | 6 | 0.028 lb. | X Packages/100 ft ² |
|--|---|--|------|---|-----------|--------------------------------|
| Root Vegetables- Carrot, celeriac, Chervil (turnip Rooted), Chicory | Armyworms, cabbage looper, Colorado Potato Beetle, cutworms, European corn borer, flea beetles, green peach aphid, leafhoppers, lygus bug, (plant bug), pea aphid, tarnished plant bug, | Do not apply more than recommended amount [0.14 lb ai/A] in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest. Other than radishes, applications may not be made to root vegetables whose tops or leaves are to be harvested for human consumption. | 2.13 | 5 | 0.028 lb | X Packages/100 ft ² |
| Tuberous & Corm Vegetables-Potato, sweet potato, arracacha, arrowroot, Chinese artichoke, Jerusalem artichoke, edible canna, bitter and sweet cassava, chayote, (root), chufa, dasheen, ginger, leren tanier, tumeric, yarn bean, true yam | Armyworm, cabbage looper, Colorado potato beetle, cutworms, European corn borer, flea beetle, green peach aphid, leafhoppers, lygus bug (plant bug), pea aphid, Tarnished plant bug. | Do not apply more than recommended amount [0.14 lb ai/A] in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest. Applications may not be made to tuberous or corm vegetables whose tops or leaves are to be harvested for human consumption. | 2.13 | 5 | 0.028 lb | X Packages/100 ft ² |

USE DIRECTIONS FOR FIRE ANT TREATMENT

Imported Fire Ant Mound Treatment: Apply Hi-Yield® Kill-A-Bug® II Dust Containing Deltamethrin with a shaker can. Thoroughly apply dust to adequately cover mound and around the perimeter (18" from the outer edge of the mound). A second treatment may be necessary if complete control is not obtained. DO NOT DISTURB THE MOUND PRIOR TO APPLICATION. If possible, treat all colonies in the vicinity. Also, treat colonies which have not yet constructed a mound.

NOTIFICATION

MAY 1 5 2009

(Alternate directions - Evenly sprinkle one tablespoon (approximately 1/2 oz.) of Hi-Yield® Kill-A-Bug® II Dust Containing Deltamethrin over the top of each mound. Do not water in or disturb the mound. Apply when temperatures are between 65 and 80°F. Allow 3 to 4 days for maximum control, then repeat at 7 days if needed to maintain control.)

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep out of reach of children and pets. Store in original containers only. Store in cool, dry place and avoid excess heat. **PESTICIDE DISPOSAL**: **If empty**: **Non-refillable container**. Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this product to conform to the chemical description on the label and for the purposes stated on the label. - The limit of any liability incurred shall be the purchase price paid by the user or buyer.

Released For Shipment By:

VOLUNTARY PURCHASING GROUPS, INC. P.O. BOX 460 230 FM 87 BONHAM, TX 75418 EPA Reg. No.7401-452 EPA Est. No. 7401-TX-1

www.hiyield.com