1/11

## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

SEP - 9 2014

Mr. Jesus Atencio Regulatory Assistant Voluntary Purchasing Groups, Inc. 230 FM 87 Bonham, TX 75418

Subject: Label Notification per PRN 98-10 – Adding Squash Bugs a Pest, Adding an Optional

Statement for Previously Approved Home Vegetable Garden Use, and Adding

Telephone Number for Product Questions Product Name: Hi-Yield Kill-A-Bug II Dust

EPA Registration Number: 7401-452 Application Date: August 5, 2014

Decision Number: 494212

#### Dear Mr. Atencio:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, you may contact Owen Beeder at 703-308-8899 or via email at beeder.owen@epa.gov.

Mark Suarez

Sinderel

Product Manager 13 Insecticide Branch

Registration Division (7505P) Office of Pesticide Programs

Please read instructions	on reveise before completing fo	orm.	Form Approved.	OMB No. 2070	-0060 ALATHELA ASSOCIA
<b>\$EPA</b>	United Environmental Pro Washington	tection Agency	×	Registration Amendme Other	A==/
	Арр	lication for Pestic	ide - Section	l	
1. Company/Product No 7401-452	mber	II.	A Product Manager nael Walsh		3. Proposed Classification
4. Company/Product (N Hy-yield Kill-A-Bug	<b>ame)</b> I Dust Containing Deltamethi	rin <b>PM#</b> 13			X None Restricted
Voluntary Pu 230 FM 87 Bonham, TX	f Applicant (Include ZIP Code) rchasing Groups, Inc 75418 If this is a new address	(b)(i), to: EPA			e with FIFRA Section 3(c)(3) I in composition and labeling
		Section -			
X Notification - Ex Explanation: Use ad Label Notification per PF regulations at 40 CFR 15 violation of 18 U.S.C. Se requirements of 40 CFR	response to Agency letter dated plain below.  ditional page(s) if necessary. (For Notice 98-10, amended labeling to 52.46. No other changes have been control to willfully make any false states 156.10, 156.140, 156.144, 156.14	or section I and Section II., with copy as highlighted. Th n made to the labeling or the statement to the EPA. I furth	is notification is consi e Confidential Statem ner understand that if	etion.  blow.  istent with the propent of Formula for the amended lab	ovisions of PR Notice 98-10 and EPA or this product. I understand it is a
penalties under sections		Section -	<u> </u>		
1. Material This Produc		DM-an-Callela	Darling's	T	. •
Child-Resistent Peckagi Yes* No * Certification mus	Yes No If "Yes" No	Water Soluble Yes No If "Yes" Package wgt	No. per container	P G P	ntainer Metal Mastic ilass Japer Other (Specify)
3. Location of Net Cont	_	ze(s) Retail Container	5. Lo	cation of Label I On Label	Directions
6. Manner in Which Lab	Container	Lithograph Paper glued	Other	On Labeling	accompanying product
		Section -	IV		0 6 0 B
1 Control Dei 1 /C-	plete items directly below for ide				
Name Jesus Atencio	SHEE REITS BEECHY DEIDW TOF TOE	Title Regulatory Ass		Te	lephone No. (Include Area@adc)
	statements I have made on this f at any knowingly false or mislea				ete.  6. Date Application 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6
2. Signature		3. Title  Regulatory Ass	istant		CCCC
4. Typed Name	•	5. Date			

Aug 4, 2014

Jesus Atencio

#### August 5, 2014

U.S. Environmental Protection Agency Document Processing Desk (NOTIF) Office of Pesticide Programs (7504P) Attn: Mr. Michael Walsh 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460

Subject:

Hi-Yield® Kill-A-Bug II Dust Containing Deltamethrin, EPA Reg. No. 7401-452, Notification per PR Notice 98-10

Dear Mr. Walsh,

This notification is made with the following additions to the product label:

- Added statement: For Use On Roses, Flowers, Ornamentals, Shrubs and Home Vegetable Gardens
- Add Pest: Squash Bugs
- Add Contact Information: Product Questions? 855-270-4776

Enclosed with this submission, please find the following:

- EPA Form 8570-1
- revised labels with additions highlighted
- One revised label with clean copy

Please update the agency records in consideration of this notification. If you should have any questions please contact me.

Sincerely,

Jesus Atencio Regulatory Assistant VPG, Inc. 230 FM 87 Bonham, TX 75418 (903) 583-5501 x121 jatencio@fertilome.com

#### [FRONT LABEL PANEL]

# Hi-Yield® Kill-A-Bug® II Dust Containing Deltamethrin

- For Indoor/Outdoor Residential Use Only
- [For Household Use]
- [Household Ornamental Garden Insect Control]

### NOTIFICATION

SEP - 9 2014

- [Controls Fire Ants, Fleas, Ticks \*Termites and More][Non-Staining & Odorless]
- [Long-Lasting Residual Action]
- [Long-Lasting]
- [Kills Insects for up to 8 Months]
- [Does not Clump or Absorb Moisture]
- [For Use on Ornamentals]
- [Use on Roses, Flowers & Ornamentals]
- [Controls Many Different Insect Pests]
- [To Control Imported Fire Ants on Lawns and Other Residential Areas]
- [8 oz. Treats 15 Fire Ant Mounds]
- [1 lb. Treats 30 Fire Ant Mounds]
- [2 lbs. Treats 60 Fire Ant Mounds]
- [Fast, Effective Kill of Fire Ants]
- [Fast, Effective Kill of Stinging Ants]
- [Easy To Use]
- [Convenient]
- [Kills Fire Ants on Contact]
- [Effective Knockdown and Long-Lasting Control of Listed Pests]
- [For Use On Roses, Flowers, Ornamentals, Shrubs, and Home Vegetable Gardens]

[ ]-Denotes Optional Statements

\*For localized area control only

#### **ACTIVE INGREDIENT:**

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back/Side Panel for Additional Precautionary Statements

. . . .

Net	Weight	
-----	--------	--

#### [Back/Side Label Panel]

Have the produc	<b>FIRST AID</b> t container or label with you when calling a poison control center or doctor, or
going for treatm	
If swallowed	<ul> <li>Call poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
If in eyes	<ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If on skin or clothing	<ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If inhaled	<ul> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to mouth, if possible.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>

#### PRECAUTIONARY STATEMENTS

#### Hazards to Humans And Domestic Animals

**CAUTION:** May be harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

#### **ENVIRONMENTAL HAZARDS**

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

#### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Deltamethrin can be used for long-lasting pest control in and around homes and their immediate surroundings. Permitted areas of use include:

\*animal quarters

\*mobile homes and

- \*apartments
- \*household greenhouses
- \*houses

- \*recreational vehicles
- \*ornamentals

This product is intended for applications with a hand duster, from the shaker can, or by other suitable means, to hiding and runway areas and other places where insects are found. Food utensils such as a teaspoon or measuring cup should not be used for food after use with pesticides.

[Option B: Spout Top Applicator---This product is intended for application to hiding and runway areas and other places where insects are found. Remove cap from top of spout. Clip the end of spout off to apply the product. Point directly to treatment area and squeeze bottle to apply product. Replace the cap on end of spout when finished applying.]

#### **INDOOR USE**

In living areas, make applications in such a manner to avoid deposits-on exposed surfaces or introducing the material into the air.

To apply insecticide directly into cracks and crevices, use a bulbous duster, or other suitable equipment. Apply lightly and uniformly to infested areas. Pay particular attention to cracks and crevices; service ducts, false floors and ceilings; wall void; around electrical and telephone fittings and equipment; around water and sewer pipes; under and behind cabinets; refrigerators and sinks; around windows and door frames; along baseboards; in attics and crawl spaces. Apply at the rate of 1 oz. of product per 125 square feet (or ½ lb. per 1000 square feet). Repeat after 10 days only if needed to maintain control. Do not apply more than 2 times per year.

Ants, boxelder bugs, cockroaches (including both resistant and non-resistant strains), crickets, fleas, firebrats and silverfish: Apply Hi-Yield® Kill-A-Bug® II Dust Containing Deltamethrin thoroughly to all areas where these pests crawl and hide, especially in those areas described above. For Ants, apply to ant trails, around doors and windows and other places where ants enter premises.

Termites, Carpenter Ants and Carpenter Bees: For Localized Treatment of Wood Infesting Termites, Carpenter Ants, and Carpenter Bees; apply to voids to channels in damaged wooden members of a structure, to cracks, spaces or bearing joints between wooden members of a structure; or between such members and the foundation; and in locations vulnerable to termite attack. Drilling may be necessary prior to applying dust in these locations, but it must be recognized that such drilling and treating of timber in a particular location may not result in complete coverage of infested and vulnerable voids or galleries. When drilling, care should be taken to avoid drilling into electrical wire, plumbing, gas lines or other utilities. Re-seal holes following treatment. This dust can be applied in attics and crawl spaces to prevent entry of drywood and formosan termites at a rate of 1/2 lb. per 1000 square feet of floor area. In these dreat treatments, all vents and blowers should be closed. Inspect at 6 months and one year to ensure that colony has been killed. Repeat after 10 days only if needed to maintain control. Do not apply more than 2 times per year.

The purpose of such applications of Hi-Yield® Kill-A-Bug® II Dust Containing Deltamethrin is to kill workers or winged forms which may be present in the treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplement. For active termite infestation, get a professional inspection.

Centipedes, ground beetles, millipedes, scorpions, spiders and sowbugs: Apply dust along

and behind baseboards, to window and door frames, corners, pipes, storage locations, attics, crawl spaces, and other areas where these pests may enter or crawl.

Wasps and Bees: It is generally advisable to treat wasp and bee nests in the evening when insects are less active and have returned to the nest. Wear protective clothing if deemed necessary to avoid stings. Using hand or power duster (actisol or similar equipment) or other suitable means, with extension tubes if necessary, thoroughly dust nets, entrance and surrounding areas where insects alight. Nests in wall voids can be located by listening with your ear against the wall. Drill a hole in the area, blow dust in and reseal. For ground wasps, dust entrance and surrounding areas. For best results check nests carefully one or two days after treatment to ensure complete kill then remove and destroy nest to prevent emergence of newly -hatched insects. If removal is not feasible, retreat nest if necessary. Do not use this product to treat overhead nests.

Webbing Clothes Moths: Use Hi-Yield<sup>®</sup> Kill-A-Bug ® II Dust Containing Deltamethrin as a crack and crevice treatment in closets and other storage areas where these pests are found. This is an adjunct treatment and will not control moth larvae already on the clothes. DO NOT apply this product directly to clothing.

Carpet Beetles (Black, Furniture, Common and Varied): Dust where these insects are found, for example: on and under edges of floor coverings, around baseboards, under rugs and furniture, and in closets.

**Bedbugs:** Apply dust thoroughly to all areas where these pests are typically found including folds and tufts of mattresses, coils or springs, cracks and hollow posts of bedsteads, upholstery of chairs and sofas, picture frame moldings and <sub>all</sub> cracks and crevices in room should be treated. Allow powder to remain in contact with surface for 4-6 hours. Thoroughly vacuum treated areas, dispose of vacuum bag and place clean linens on bed.

Stored Product Pests: To aid in control of exposed stages of:

cadelles
cheese mites
cigarette beetles
confused flour beetles
dark meal worms
drugstore beetles
grain mites
granary weevils
lesser grain borers

merchant grain beetles red flour beetles rice weevils rusty grain beetles saw-toothed grain beetles spider beetles tobacco moths yellow meal worms

In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Clean up wastes materials, dust, dirt, and other debris. Thoroughly treat floors, walls and other surfaces in storage and handling areas. Apply dust in cracks and crevices, behind and under cupboards, and other difficult to reach areas where insects may hide. DO NOT apply directly on food, utensils or other areas where food is prepared, processed, served, or stored.

#### ORNAMENTAL PLANT PESTS For the control of:

aphids beet armyworms

leafrollers orange tortrix blister beetles boxelder bugs certain leafminers (Boxwood, Holly Oak, Spruce Needle elm leaf beetle flea beetles Japanese beetles periodical cicada plant bugs rose chafer rose midges rose slugs tent caterpillars thrips weevils Leafhoppers

**Shrubs and Ornamentals:** Apply when insects or their damage first appear. Repeat after 10 days only if needed to maintain control. Do not apply more than 2 times per year. Apply 8 ounces per 10000 square feet; or thoroughly apply to place a thin layer of dust on both upper and lower sides of leaves, stems and flowers.

Roses, Petunias, Chrysanthemums, Geraniums, Carnations and other related flowers: To control many leaf and flower chewing insects, apply as soon as the insect appears. Repeat after 10 days only if needed to maintain control. Do not apply more than 2 times per year. Apply a thin layer of dust on both upper and lower sides of leaves, stems and flowers.

#### **OUTDOOR USE**

**Perimeter Treatment:** Use Hi-Yield<sup>®</sup> Kill-A-Bug<sup>®</sup> II Dust Containing Deltamethrin as a residual treatment for control of ants, bees, centipedes, cockroaches, crickets, firebrats, fleas, ground beetles, millipedes, scorpions, silverfish, slugs, sowbugs, spiders, ticks and wasps. Make a residual treatment around windows and doors, porches, screens, eaves, patios, garages, under stairways, in crawl spaces where pests are active and may find entrance.

Fleas and Ticks in Rodent Burrows: To control fleas and ticks in and around rodent burrows around homes and other buildings, apply dust with a shaker can. Thoroughly apply dust to adequately cover mound and around the perimeter (18" from the outer edge). Be sure to treat abandoned rodent burrows which are suspected of containing fleas and ticks. Treat for a minimum of 5 days for light rodent populations, 10 days when populations are dense. Retreatment should only be made as necessary to maintain insect control.

HOME VEGETABLE GARDENS: Use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate available duster. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest comer of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas, before reentering.

VEGETABLE	TARGET PESTS	REMARKS	Application Rate: Oz. of Dust/100ft <sup>2</sup>	Max Application/ Growing Season	[Active Ingredient/Acre] (This column is alternate text)	Packages/100ft <sup>2</sup> (This column is occide alternate text)
Artichoke, Globe	Aphids, armyworm, cabbage looper, cribrate weevil, leafminers, lygus bug (plant bug), pale striped flea beetles, plume moth, probas bug	Do not apply more than recommended amount [0.14 lb ai/A (active ingredient/acre)] in one growing season. Allow 3 days	2.13	5	0.028 lb.	X Packages/100 ft <sup>2</sup> ccc

		between applications.  Do not apply within				
Bulb vegetables-	Cutworms, squash bugs,	3 days of harvest.  Do not apply more	2.13	4	0.028 lb.	X Packages/100 ft <sup>2</sup>
garlic, leeks, onions, shallots	stink bugs, thrips	than recommended amount [0.112 lb				
		ai/A (active ingredient/acre)] in				
		one growing season. Allow 5 days				
		between applications. Do not apply within				
Corn, Sweet	Aphids, armyworms,	one days of harvest.  Do not apply more	2.13	16	0.028 lb ·	X Packages/100 ft <sup>2</sup>
	chinch bugs, click beetle, corn earworm, corn rootworm (adult),	than recommended amount [0.45 lb ai/A] in one growing				
	cutworms, European corn borer, flea beetles,	season. Do not apply within 24 hours of				
	grasshoppers, Japanese beetle (adult), June beetle (adult), Masked chafer	harvest.				
	(adult), Southern corn borer, Southwestern corn borer, stalk borer, squash					
	bugs, stink bugs, webworms					
Corn, Pop	Aphids, armyworms, chinch bugs, click beetle,	Do not apply more than recommended	1.68	4	0.022 lb	X Packages/100 ft <sup>2</sup>
	corn earworm, corn rootworm (adult), cutworms, European corn	amount [0.095 lb ai/A] in one growing season. Do not apply		,		
	borer, flea beetles, grasshoppers, Japanese	within 21 hours of harvest.				
	beetle (adult), June beetle (adult), Masked chafer (adult), Southern corn				·	
•	borer, Southwestern corn borer, stalk borer, squash					
	bugs, stink bugs, webworms					
Curcurbit Vegetables-	Armyworm, cabbage looper, Colorado potato	Do not apply more than recommended	2.13	6	0.028 lb	X Packages/100 ft <sup>2</sup>
Cantaloupes, Chayote, Chinese	beetle, corn earworm, cucumber beetle,	amount [0.168 lb ai/A] in one growing				
Wax Gourd, Citron Melon, Cucumber,	cutworms, diamondback moth (larvae), Flea beetle,	season. Allow 3 days between applications.				
Gherkin, gourds, Momordica species	grasshoppers, green peach aphid, leafhopper,	Do not apply within 3 days of harvest.				•
balsom pear, bitter melon, Chinese	leafminer, melonworm, pickleworm, squash bugs, stinkbugs, tarnished plant		i c			l .
cucumber) muskmelon, pumpkin	bug, tomato pinworm, vegetable weevil (adult),					6666 6 6 6000
squash, watermelon	western plant bug, whitefly (adult).					6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6
				<b>.</b> .	,	6 <u>6</u> 66 6 6
Fruiting Vegetables-	Aphids, armyworms, cabbage looper, Colorado	Do not apply more than recommended	2.13	6	0.028 lb.	K Packages/100 ft <sup>2</sup> c c
Eggplant, groundcherry,	Potato Beetle, cucumber	amount [0.168 lb				C C C C C C C C C C C C C C C C C C C
pepinos, peppers, tomatillo, tomato	beetle, cutworms, diamondback moth,	ai/A] in one growing season. Allow 5 days				c 666
	European com borer, flea beetles, imported	between applications.  Do not apply within				( (
į	cabbageworm, leafminers, pepper weevil, squash	1 days of harvest.				(
	bugs, stink bugs, tarnished plant bug, thrips, tomato					i ( ( (

	fruitworms, (corn	!	· · · · · · · · · · · · · · · · · · ·	1		1
	earworm), tomato					
	pinworm, variegated		ł		'	
	cutworm, whitefly	-				
Root Vegetables-	Armyworms, cabbage	Do not apply more	2.13	5	0.028 lb	X Packages/100 ft <sup>2</sup>
Carrot, celeriac,	looper, Colorado Potato	than recommended				
Chervil (turnip	Beetle, cutworms,	amount [0.14 lb ai/A]				_
Rooted), Chicory	European corn borer, flea	in one growing				
, ,	beetles, green peach aphid,	season. Allow 3 days				
	leafhoppers, lygus bug,	between applications.			1	
	(plant bug), pea aphid,	Do not apply within			1	
	tarnished plant bug,	3 days of harvest.				
	. 5	Other than radishes,				
	l	applications may not				,
		be made to root	i i	ľ	*	
•		vegetables whose				
•		tops or leaves are to	•			
		be harvested for				}
		human consumption.		<u> </u>		
Tuberous & Corm	Armyworm, cabbage	Do not apply more	2.13	5	0.028 lb	X Packages/100 ft <sup>2</sup>
Vegetables-Potato,	looper, Colorado potato	than recommended				
sweet potato,	beetle, cutworms,	amount [0.14 lb ai/A]				)
arracacha, arrowroot,	European corn borer, flea	in one growing		·		
Chinese artichoke,	beetle, green peach aphid,	season. Allow 3 days				
Jerusalem artichoke,	leafhoppers, lygus bug	between applications.				
edible canna, bitter	(plant bug), pea aphid,	Do not apply within				
and sweet cassava,	Tarnished plant bug.	3 days of harvest				
chayote, (root),		Applications may not				
chufa, dasheen,		be made to tuberous		·		
ginger, leren tanier,		or corm vegetables				
tumeric, yarn bean,		whose tops or leaves are to be harvested	-			
true yam	1	for human				
,		consumption.			,	
	<u> </u>	Consumption.		L	L	I

#### USE DIRECTIONS FOR FIRE ANT TREATMENT

Imported Fire Ant Mound Treatment: Apply Hi-Yield® Kill-A-Bug® II Dust Containing Deltamethrin with a shaker can. Thoroughly apply dust to adequately cover mound and around the perimeter (18" from the outer edge of the mound). A second treatment may be necessary if complete control is not obtained. DO NOT DISTURB THE MOUND PRIOR TO APPLICATION. If possible, treat all colonies in the vicinity. Also, treat colonies which have not yet constructed a mound.

(Alternate directions - Evenly sprinkle one tablespoon (approximately 1/2 oz.) of Hi-Yield® Kill-A-Bug® II Dust Containing Deltamethrin over the top of each mound. Do not water in or disturb the mound. Apply when temperatures are between 65 and of core 80°F. Allow 3 to 4 days for maximum control, then repeat at 7 days if needed to maintain control.)

#### STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep out of reach of children and pets. Store in original containers only. Store in cool, dry place and avoid excess heat.

PESTICIDE DISPOSAL: If empty: Non-refillable container. Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your locate.

solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**NOTE:** Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this product to conform to the chemical description on the label and for the purposes stated on the label. - The limit of any liability incurred shall be the purchase price paid by the user or buyer.

#### Manufactured By:

VOLUNTARY PURCHASING GROUPS, INC.
230 FM 87
BONHAM, TX 75418
EPA Reg. No.7401-452
EPA Est. No. 7401-TX-1

www.hi-yield.com
Product Questions? 855-270-4776

Page 8