

NuLife WEED AND FEED

10-2-3

COVERS 5,000 SQ. FEET

GENERAL DIRECTIONS

NuLife WEED and FEED contains 2,4-D and Silvex weed killers and fertilizer to beautify your lawn.

1. Controls broad-leaved weeds such as dandelions, plantains, chickweeds, clovers and most non-grass weeds.
2. Feeds the grass and assists in producing a thicker turf of deeper color.

USE AT THE RATE OF 1 LB. TO 125 SQ. FT.

MATURE LAWNS: Apply with fertilizer spreader or by hand. If by hand, using protective gloves, measure off lawn in squares to insure even coverage. Broadcast NuLife WEED and FEED close to the surface of lawn. If using fertilizer spreader be sure it is set for proper distribution. Close spreader when stopping, backing or turning to avoid double dosage.

NEW LAWNS: Burning may result if applied to tender seedling grass less than 4 or 5 months old.

SPOT TREATMENT OF WEEDS: Apply a pinch of WEED and FEED in crown of each weed. A good time to spot treat weeds is at time of mowing lawn. Use spot treatment especially after weeds are under control.

IMPORTANT

The most satisfactory results from application of NuLife WEED and FEED are obtained if the following steps are taken (preferably in this order):

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least one day before application; this will sustain moisture until the next watering (See step 4).
3. Apply when weeds are young and actively growing and preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold material and prevent dust drift onto plants which might be injured by weed killers.
4. NuLife WEED and FEED should not be washed from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

CAUTION!!

HARMFUL IF SWALLOWED. KEEP OUT OF REACH OF CHILDREN. Avoid inhalation of dust or contact with eyes, skin and clothing. Wash hands thoroughly after handling. Induce vomiting if swallowed. If material contacts the eyes, get medical attention immediately. Keep children and animals off treated areas. Do not store near fertilizers, seeds, insecticides or fungicides.

Excessive application will injure turf. When repeat treatments are necessary, allow one month between treatments. Do not use on St. Augustine grass. Carpet grass, Dichondra or clover lawns. Treatment above 80° may result in some discoloration of Bent grass. This material is injurious to many plants. It should be applied while the air is quiet to avoid drift and application on valuable plants. Vapors from this product may cause injury to susceptible plants growing in the immediate vicinity.

Environmental Note: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning equipment or disposal of wastes.

Clean fertilizer spreader several times with chemical cleaners and rinse thoroughly before using spreader for other purposes. Do not reuse this container, dispose of with refuse going to a sanitary landfill area.

NOTICE!!

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

MINIMUM GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) 10.0%

40# / 5,000 sq. ft.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least one day before application; this will sustain moisture until the next watering (See step 4).
3. Apply when weeds are young and actively growing and preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold material and prevent dust drift onto plants which might be injured by weed killers.
4. Nutlife WEED and FEED should not be washed from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

CAUTION!!

HARMFUL IF SWALLOWED. KEEP OUT OF REACH OF CHILDREN. Avoid inhalation of dust or contact with eyes, skin and clothing. Wash hands thoroughly after handling. Induce vomiting if swallowed. If material contacts the eyes, get medical attention immediately. Keep children and animals off treated areas. Do not store near fertilizers, seeds, insecticides or fungicides.

Excessive application will injure turf. When repeat treatments are necessary, allow one month between treatments. Do not use on St. Augustine grass, Carpet grass, Dichondra or clover lawns. Treatment above 80° may result in some discoloration of Bent grass. This material is injurious to many plants. It should be applied while the air is quiet to avoid drift and application on valuable plants. Vapors from this product may cause injury to susceptible plants growing in the immediate vicinity.

Environmental Note: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning equipment or disposal of wastes.

Clean fertilizer spreader several times with chemical cleaners and rinse thoroughly before using spreader for other purposes. Do not reuse this container, dispose of with refuse going to a sanitary landfill area.

NOTICE!!

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

MINIMUM GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) (Derived from Ammonium Sulfate and Sterilized Organic Matter)	10.0%
Available Phosphoric Acid (P ₂ O ₅) (Derived from Sterilized Organic Matter)	2.0%
Soluble Potash (K ₂ O) (Derived from Monite of Potash)	3.0%

ALSO CONTAINS

Magnesium (Mg) (Derived from Dolomitic Limestone)	0.70%
Sulfur (S) (Derived from Ammonium Sulfate and Sterilized Organic Matter)	9.70%

Potential Acidity: 727 lbs. Calcium Carbonate Equivalent Per Ton.

RECOMMENDED FERTILIZER SPREADER SETTINGS*	
Spread one way, being careful not to skip.	
MAKE OF SPREADER	RATE
SCOTT	5½
LAWN BEAUTY	6½
SEARS	6

*As calibration and general condition of spreader can materially affect the rate of flow, neither the manufacturer nor seller can guarantee exact coverage as indicated by the above recommended dial settings.

WITH WETTING AGENT ADDED

Nutlife Fertilizers
TACOMA, WASH. 98421

EPA Est. 7404-WA-1

015



FREIGHT SHIPPING BAG
Meeting requirements of
APPLICABLE FREIGHT CLASSIFICATION
Guaranteed by
BEMIS COMPANY, INC.
VANCOUVER, WASH.

Turf King

Makes Better Lawns

Covers
5000
sq. ft.

Kills
Weeds

Feeds
Grass

with
wetting
agent

ACCEPTED

JAN 22 1970
Comment on letter
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 7402-8

Weed & Feed

Makes Better Lawns



JAN 23 1976
Comments on Esters
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 7404-8

Weed & Feed 3-in-1

HERBICIDAL CONTENT

ACTIVE INGREDIENTS:
2,4-Dichlorophenoxyacetic Acid
2-Ethylhexyl ester of silver*

By Wt.
0.750%
0.355%

INERT INGREDIENTS:

98.895%
100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL
CAUTION STATEMENTS.

EPA No 7404-B-18527

*Eq. to 0.248% Silvex 2 (2,4,5-T-chlorophenoxy)propionic acid

25 LBS. NET • COVERS 5,000 SQ. FT.



Weed & Feed



Turf King

Weed & Feed 3-in-1

WILL NOT BURN IF USED ACCORDING TO DIRECTIONS

WEED and FEED beautifies the lawn 3 ways:

1. Controls unwanted plants - released red color with 2.4-D and Glyphosate.
2. Helps produce a thicker, greener lawn.
3. Contains Wetting Agent to increase water penetration into soil.

GENERAL DIRECTIONS

APPLICATION: At a rate of 1/8 to 20 lbs. per acre.

For Best Results:

1. Mow lawn to normal height 2 to 3 days before application.
2. Water lawn thoroughly 1 day and dry before application.
3. Apply under nozzle in young and actively growing and in early morning when there is no wind. If grass is very hot from sun, lightly water material on grass and keep present drift away.
4. Lower sprayer when mowing, raking up or cutting to avoid damage from over application.
5. Do not mow material from weed areas for 1 to 2 days after which a thorough watering should be done.

WEED CONTROL: To avoid damage to new lawns, do not spray grasses of less than 3 months old. Application may be made at any time during the growing season when temperatures will not exceed 80°. Do not apply to bare ground.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen N 16.0%

Water Soluble Ammonium Sulfate 16.0%

Available Phosphoric Acid P₂O₅ 2.0%

Water Soluble Phosphate 2.0%

Soluble Potash K₂O 0.0%

ALSO CONTAINS

Iron Fe 1.0%

Water Soluble Iron 1.0%

Potential Acidity: 723 lbs. Calcium Carbonate per ton.

RECOMMENDED FERTILIZER SPREADER SETTINGS*	
Spreader setting	1/2
Spreader speed	6 mph
TURF KING	3/4
SCOTT	5

*The above settings are recommended for spreading on lawns. For best results, spread evenly and uniformly. Do not apply to bare ground.

CAUTION

Do not apply to lawns containing St. Augustine grass. Avoid application to lawns where seed heads have been scattered on the grass. Avoid spraying if it is allowed to rain. Do not apply to lawns where seed heads have been scattered on the grass and the grass is thoroughly dry. KEEP OUT OF REACH OF CHILDREN.

Avoid applying to lawns containing St. Augustine grass. When repeated applications are made to lawns containing St. Augustine grass, do not use on St. Augustine grass lawns more than 1/2 lb. per acre per year. A temperature of 70° F. results in some discoloration of Bermudagrass, but not injury. A temperature of 70° F. has been reported to injure Zoysia grass. Do not apply to lawns containing St. Augustine grass, or Zoysia grass, or Bermuda grass, or other warm season grasses growing in the immediate area. Do not apply to lawns containing cool season grasses.

ENVIRONMENTAL NOTE: Do not apply to lawns containing St. Augustine grass. Do not feeding on treated area may result in discoloration and/or death of grass. Do not contaminate water by cleaning of equipment or person of user.

DO NOT USE ON LAWNS CONTAINING CHEMICAL PLANTS AND DO NOT THOROUGHLY BURN BURNABLE PLANTS. DO NOT BURN BURNABLE PLANTS IN AREAS WHERE BURNING IS PROHIBITED OR WHERE BURNING IS PROHIBITED.

STORE IN A COOL DRY LOCATION

100% GUARANTEED ANALYSIS

Distributed by ROUNDUP CO., Spokane, Washington 99220

Turf King

WILL NOT BURN IF USED

WEED and FEED beautifies the lawn

1. Controls dandelion, plantain, chickweed
2. Helps produce a thicker, greener turf.
3. Contains Wetting Agent to increase water

GENERAL

APPLICATION: USE AT RATE OF 1 LB. TO 20
For Best Results:

1. Mow lawn to normal height 2 to 3 days before application.
2. Water lawn thoroughly at least one day before application.
3. Apply when weeds are young and actively growing. If grass is not moist, sprinkle lightly to prevent damage to plants that might be injured by weed killer.
4. Close spreader when stopping, backing up or turning.
5. Do not wash material from weed leaves for several hours after application is completed.

WEED CONTROL: To avoid damage to new grass seed, application may be made any time during the growing season. Do not apply in rain.

GUARANTEED FERTILIZER

Total Nitrogen (N)
(Derived from Ammonium Sulfate, Organic Matter and Urea)

Available Phosphoric Acid
(Derived from Sterilized Organic Materials)

Soluble Potash (K₂O)

ALSO

Iron (Fe)
(Derived from Sterilized Organic Materials)

Potential Acidity: 723 lbs.

RECOMMENDED FERTILIZER

Spread one way, 100 ft.

Make of Spreader

TURF KING

SCOTT

*As calibration and general conditions vary, it is impossible to determine the rate of flow neither manufacturer nor distributor can guarantee coverage as indicated by the above figures.

2. Water lawn thoroughly at least one day before application.
 3. Apply when weeds are young and actively growing and in early morning when dew is on the grass. If grass is not moist, sprinkle lightly to hold material on grass and help prevent drift onto plants that might be injured by weed killers.
 4. Close spreader when stopping, backing up or turning to avoid damage from over application.
 5. Do not wash material from weed leaves for 1 to 2 days, after which a thorough watering should be done.
- WEED CONTROL:** To avoid damage to new turf apply after grass is at least 4 to 5 months old. Application may be made any time during the growing season when temperatures will not exceed 80°. Do not apply in rain.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) 16.0%

(Derived from Ammonium Sulphate, Sterilized
Organic Matter and Urea)

Available Phosphoric Acid (P₂O₅) 2.0%

(Derived from Sterilized Organic Matter)

Soluble Potash (K₂O) 0.0%

ALSO CONTAINS

Iron (Fe) 1.0%

(Derived from Sterilized Organic Matter)

Potential Acidity: 723 lbs. Calcium Carbonate per ton.

RECOMMENDED FERTILIZER SPREADER SETTINGS*

Spread one way, being careful not to skip

Make of Spreader	Rate
TURF KING	3½
SCOTT	5

*As calibration and general condition of spreader can materially affect the rate of flow, neither manufacturer nor seller can guarantee exact coverage as indicated by the above recommended dial settings.

CAUTION

Do not handle this material with bare hands. Avoid inhalation of dust or contact with eyes, skin or clothing. If material contacts the eyes, get immediate medical attention. Induce vomiting if swallowed. Keep children and pets off treated areas until this material is washed into the soil and the grass is thoroughly dry. **KEEP OUT OF REACH OF CHILDREN.**

Apply this product only as specified on this label. Excessive application will injure turf. When repeated treatments are necessary, allow one month between treatments. Do not use on St. Augustine grass, Carpet grass, Dichondra or Clover lawns. Treatments above 80° temperature will result in some discoloration of Bentgrass. Do not apply if temperature is expected above 80°F. This material is injurious to many plants and should only be applied while the air is quiet to avoid drift onto valuable plants. Vapors from this product may cause injury to susceptible plants growing in the immediate area. Do not store near fertilizers, seeds, insecticides or fungicides.

ENVIRONMENTAL NOTE: This product is toxic to fish and wildlife. Birds feeding on treated area may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of waste.

Clean fertilizer spreader several times with chemical cleaners and rinse thoroughly before using spreader for other purposes. Do not reuse or burn container. Dispose of with refuse going to a sanitary landfill area.

STORE IN A COOL DRY LOCATION

Manufactured at EPA Est. 7404-WA-1

Distributed by ROUNDUP CO., Spokane, Washington 99220

ACCEPTED WITH COMMENTS