

[7405-52]

Active Ingredients

* (5-Benzyl-3-furyl)-methyl-2,2-dimethyl-3-(2-methylpropenyl)-cyclopropanecarboxylate	0.900% w/w
Related compounds	0.068%
Aromatic petroleum hydrocarbons	0.667%
Petroleum distillate	13.650%
Inert ingredients	85.120%
	100.000%

* cis-trans isomers ratio: max. 30% (+) and min. 70% (+) trans
 † U.S. Patent No. 3,465,007

Directions for Use

Turn off fans and air conditioners. Put out all flames and pilot lig. Close doors and windows. Point valve opening away from face and eyes when releasing. Use one bomb for each 10,000 cu. ft. of unobstructed area.

1. Operate Valve. Can be used manually or valve can be locked in open position. To lock valve in open position for automatic discharge, press arrow on valve handle all the way forward, hooking the catch over the metal rim.

For Mosquitoes, Gnats and Moths (Millers). For best results, first spray distant rooms manually spraying toward ceiling filling room with mist. Place bomb on stand or table in center of unobstructed area, then lock valve open. Leave building at once and keep building closed for 2 hours before airing out. Open all doors and windows and allow to air thoroughly. Repeat when necessary.

For Crawling Insects. Roaches, Earwigs, Grain Mites, Wasps, Spiders and Centipedes. Spray thoroughly in hiding places such as cracks, crevices, moist areas, plumbing ground pipes and sinks, under refrigerators, cardboard and storage areas. Spray directly on insects where possible. Ants. Spray trails, nests and points of entry. Spray on ants where possible. Repeat when necessary.

For Use as a space spray in food processing plants only when plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. In hospitals, patients should be removed from room prior to treatment. Room should be ventilated for 2 hours after spraying. Do not return patients to room until after ventilation. Avoid contamination of food and feedstuff. Remove pets, birds and fish bowls.

WARNING: Contents under pressure. Do not puncture. Exposure to prolonged sunlight or other sources of heat may cause bursting. Do not throw into fire or incinerator. Do not expose to temperatures above 120° F. Do not spray on walls, composition tile or plastic. Do not spray over exposed foods or utensils. Avoid contamination of feed and foodstuffs.

MADE IN U.S.A.

A
F
UN
FOR
UN:

No. 50 INSECT SPRAY

LOCK VALVE FOR CONTINUOUS SPRAYING

EPA Reg. No. 7405-52




For

- | | |
|-----------------------------|-------------------------|
| BAKERIES | HOTELS |
| FOOD PLANTS | MOTELS |
| RESTAURANTS | OFFICE BUILDINGS |
| SCHOOLS | HOSPITALS |
| FOOD HANDLING PLANTS | |

ONE SPRAYING KILLS EXPOSED STAGES
 Flies, Mosquitoes, Gnats, Moths (millers),
 Wasps, Roaches, Earwigs, Grain Mites, Sowbugs,
 Spiders, Centipedes and Ants.

NET WT. 14 OZ.

**CAUTION: KEEP OUT OF REACH OF CHILDREN
 CONTENTS UNDER PRESSURE.**



[7405-52]

Active Ingredients:
 * + (5-Benzyl-3-furyl) methyl 2, 2-dimethyl-3-
 (2-methylpropenyl) cyclopropanecarboxylate 0.500% w/w
 Related compounds 0.000%
 Aromatic petroleum hydrocarbons 0.667%
 Petroleum distillate 13.630%
 Inert Ingredients: 85.120%
 100.000%

* Cis/trans isomers ratio: max. 30% (+) and min. 70%
 (+) trans.
 + U.S. Patent No. 3,465,007

Directions for Use

Shut off fans and air conditioners. Put out all flames and pilot lights. Close doors and windows. Point valve opening away from face and eyes when releasing. Use one bomb for each 10,000 cu. ft. of unobstructed area.

To Operate Valve: Can be used manually or valve can be locked in open position. To lock valve in open position for automatic discharge, press arrow on valve button all the way forward, hooking the catch over the lip of metal rim.

Flies, Mosquitoes, Gnats and Moths (Millers): For best results first spray distant rooms manually spraying toward ceiling filling room with mist. Place bomb on stand or table in center of unobstructed area, then lock valve open. Leave building at once and keep building closed for 2 hours before airing out. Open all doors and windows and allow to air thoroughly. Repeat when necessary.

For Crawling Insects: Roaches, Earwigs, Grain Mites, Sowbugs, Spiders and Centipedes: Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insects where possible. Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Caution: Use as a space spray in food processing plants only when plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. In hospitals, patients should be removed from room prior to treatment. Room should be ventilated for 2 hours after spraying. Do not return patients to room until after ventilation. Avoid contamination of food and feedstuff. Remove pets, birds and fish bowls.

WARNING: Contents under pressure. Do not puncture. Exposure to prolonged sunlight or other sources of heat may cause bursting. Do not throw into fire or incinerator. Do not expose to temperatures above 120° F. Do not spray on walls, composition tile or plastic. Do not spray over exposed foods or utensils. Avoid contamination of feed and foodstuffs.

SOLD BY



ACCEPTED

Feb-7-1972

UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND ROBERTSON
 FOR ECONOMIC POISON RESISTANCE
 UNDER NO. 7405-52

No. 50

INSECT SPRAY

LOCK VALVE FOR CONTINUOUS SPRAYING

EPA Reg. No. 7405-52

For

- BAKERIES
- FOOD PLANTS
- RESTAURANTS
- SCHOOLS
- HOTELS
- MOTELS
- OFFICE BUILDINGS
- HOSPITALS
- FOOD HANDLING PLANTS

ONE SPRAYING KILLS EXPOSED STAGES
 Flies, Mosquitoes, Gnats, Moths (millers),
 Wasps, Roaches, Earwigs, Grain Mites, Sowbugs,
 Spiders, Centipedes and Ants.



CHEMICAL Packaging CORPORATION

FT. LAUDERDALE FLORIDA 33309

LOS ANGELES CALIFORNIA 91733