3

BEST AVAILABLE COLY

LAST-BITE

SNAIL & SLUG KILLER

MEAL

A SOIL SURFACE TREATMENT FOR CONTROL OF SNAILS & SLUGS IN CITRUS GROVES- APPLE & PEACH ORCHARDS - BERRY CROPS VEGETABLE GARDENS - ORNAMENTAL GARDENS - SEED GRASSES DICHONDRA & TURFGRASS LAWNS - IN ORNAMENTAL GREENHOUSES

ACTIVE INGREDIENT:

Metaldehyde (2,4,6,8-tetramethyl -1,3,5,7-tetraoxycyclo-octane) --- 2.75% INERT INGREDIENTS: 97.25%

Total

100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

- STATEMENTS OF PRACTICAL TREATMENT -

if SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. IF ON SKIN: Wash with soap and warm water. IF IN EYES: Flush with flowing water. Get medical attention if irritation persists.

THIS PESTICIDE MAY BE FATAL TO DOGD OR OTHER PETS IF EATEN KEEP PETS OF THE ATED AREA

SEE SIDE BACK PANEL FOR AIDDITIONAL PRECAUTIONARY STATEMENTS
EPA REG. NO. 7421EPA EST. NO. 7421-CA-1

NET WEIGHT 36 OZ. (2 1/4 LBS.) Also Available in 20 LB. and 40 LB.

CALIFORNIA LIQUID FERTILIZER COMPANY 1801 Highland Avenue - Duarte, California 91010

PRECAUTIONARY STATEMENT'S
HAZARUS TO HUMANS & DOMESTIC ANIMAL
DO NOT TAKE INTERNALLY. DO NOT GET N
EYES, DO NOT GET ON SKIN. CAUTION: May be
harmful if swalfowed inhaled or absorbed through the skin. Do not
breathe dust, get in eyes, on skin or on clothing. Wash thoroug by
with soap and warm water after handling. Wash contaminated clothing with soap and warm water before re-use. If poisoning occitin,
obtain prompt medical aid. Keep granules away from, dogs and pie,
balt may be attractive to dogs. Confine pets during application or

ENVIRONMENTAL HAZARDS

prevent them from believing they are being feed.

Do not apply directly to water or wetlands (swamps, bogs, marship, and potholes). Do not contaminate water when disposing of equipment was maters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconstrent with its labelling, READ ENTIRE LABEL, USE IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE, DO NOT APPLY DIRECTLY ONTO PLANTS, Do not allow ballot contact the edible portions of any food crop.

STORAGE AND DISPOSAL

On not contaminate water, food or feed by storage or disposal. STORAGE Store product in original labeled container in a cool, dir, locked prace out of reach of children Keep container tightly closed when not in use. Do not store in direct sunlight. PESTICIDE DISPOSAL: Wastee resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Completely empty beg into application equipment. Then dispose of empty bag in a sanitary landfill, or by scineration, or, if allowed by State & focal authorities, by burning the turned stay out of smoke. HOUSEHOLD: On not reuse empty optioner, Wrea in several larvers of newpoors and dispard in trash.

IMPORTANT - HOW TO USE Peacl entire label before using, LAST-BITE Snail & Slug Killer Me

a balt product formulated for control of Snalls and Slugs. The pe will be attracted to the meal and will be liffled quickly. These peals : night feeders and their presence can be detected by their sh mucous-like trade. They seek sheller during the day in protected, or areas and generally hide in damp locations around flower bade, namentals, hedges, and especially in low-growing cover, such as N Dichondra lawns and young bedding plants. LAST-BITE Snall & St Killer Meal should be broadcast less in the day, before or after th oughly wetting the areas suspected to be infested. During periods a damp weather or continuous watering the peets multiply rapidly as it may be necessary to reapply belt at two week intervals - even mo frequently during extremely heavy infestations. New broads can usually be controlled by 3 bartings at 10 day intervals. Best results (obtained if soil is wet before use and dry soil should be mousten before application. Do not allow Meal to come in contact with parts. plant that may be used for food or feed purposes. Apply any time the day but evening applications are preferred. Do not re-wear for hours after application.

NOTICE

Seller makes no werranty, expressed or implicately seller of this product other than indicated on the laboratory seller than indicated on the laboratory seller than the manufacture of use and/or handling of this manufacture.

DEC 1 7 1990

Under the Federal Insociation
Functivide, and Ruskinderic Act
as amended, for the product
registered under EPA to 186
7421-14

APPLICATION RATE

Use 1 lb. to treat 1,100 to 1,500 Sq.Ft. For control of SNALS and SLUGS in the following areas, of soll surface equivalent to 30 to 40 pounds granules per acre. Apply when first plant damage appears or is enticipated and repeat as needed for control.

FRUIT TREE CROPS

Apply at the rate of 40 lbs. per scre.

APPLES, APRICOTS, AVOCADOS, CHERRES, PEACHES,
PEARS, and CITPLES; (Oranges, Laurons, Grapefulit, etc.): Broadcast liberally under trees, around irrigation pipes, farm buildings
and any other place where slugs and snails may congregate.

BERRY CROPS

Use up to 40 lbs. per acre.
BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, FASP-BERRIES, STRAWBERRIES, For soil surface coverage;

VEGETABLE CROPS AND GARDENS

Apply up to 30 to 40 lbs per acre.
For use around Artichoke, Asparague, Beans, Beets, Blackeyed
Peas, Broccoll, Brussele Sprouts, Cabbage, Cantaloupe, Cucumbers, Eggplant, Endive, Gartic, Ginseng, Horseradish, Kale,
Kohlrabi, Leeks, Lettuce, Melons, Mustard Greens, Okra, Onions,
Parnips, Peas, Peppers, Pimentos, Potatoes, Pumplons, Racishes, Rhubarth, Rutabagae, Salsily, Shallots, Spinach, Squash,
Sweet Potatoes, Swisa Chard, Tomatoes, Turnips, Watermelons,

ORNAMENTAL PLANTS

One pound will cover up to 1,100 Sq. Pt.
OPNAMENTAL GARDENS, PLANTED BORDERS AND GROUND
COVERS: Including Azalasa, Begonias, Camellas, Chrysanthemums, Daffodille, Daleles, Geraniums, Honeysucide, Hydrangets, Ice Plant, Impadene, Iris, Marigolds, Paneles, Petunias,
Rhod odendron, Rose, Sweet Peas, Enapdragons,
Spathiphyllum, Tulips. Water the infested area thoroughly before
application. Broadcast or sprintide LAST BITE Snall & Stug Meal
evenly over the infested areas and along edges of flower bade
adjacent to border or ground cover plants. Also apply on ground
at base of fences and walls in shedy, protected locations. Do not
re-water for 48 hours.

DICHONDRA AND/OR TURFGRASS LAWNS

One pound will cover up to 1,500 sq. ft. Water infested area thoroughly before application. Broadcast LAST-BITE Snail & Stug Killer Meal directly on lewn or turf. Spread evenly. Do not re-water for 46 hours.

ORNAMENTAL GREENHOUSES

One pound will cover up to 1,100 aq.ft. Water or interes greenhouse sof to be treated before application. Broadcast UST-BITE Shall & Slug littler Meal Riperally around greenhouse, under benches in greenhouses and around plants; any place where SNALS and SLUGS might congregate. Do not apply directly to plants unless prior experience has shown the plant to be tolerant to this product. Most plants are tolerant to UST-BITE Meal houseurs some alight tip burning has been notice on extremely limited greenhouses a second application is recommended 7 to 10 days following the first application to bring the house under control.

SEED GRASSES

Alfalfa, Berley, Hay, Oats, Wheat, Apply up to 40 lbs, say acc