

PRECAUTIONARY STATEMENTS:

HAZARD TO HUMANS & DOMESTIC ANIMALS:

CAUTION: Keep away from humans, domestic animals, and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K intramuscularly or orally, as indicated in bishydroxycoumarin over dosages. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD:

Keep Out of Lakes, Streams or Ponds.

EPA Reg. No. 7455-12

EPA Est. 995-MN-1

BEST AVAILABLE COPY

TAKE TIME



OBSERVE LABEL DIRECTIONS

 **INTERNATIONAL MULTIFOODS**
AGRICULTURAL PRODUCTS DIVISION
MINNEAPOLIS, MN 55402/DISTRIBUTOR

8118-981

ACCEPTED

JAN 13 1982

Handwritten notes and stamps:
145-12
10/2

Supersweet[®]

RODENT RID[™]

Faint, mostly illegible text, possibly containing product details or contact information.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Handwritten text: CONTROLLED SUBSTANCE

Handwritten: PM 19

Handwritten: 7455-12

Handwritten: 1 of 2



2 of 2

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS — For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar manmade structures.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS— Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow.

Remove as much food as possible.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply ¼ to ½ ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water and harborage as much as possible. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

DISPOSAL

DO NOT REUSE EMPTY CONTAINER.

WRAP CONTAINER AND PUT IN TRASH COLLECTION.

