

DIRECTIONS

DAIRY MIST

A SPRAY SOLUTION: For use in Foggers, Mist Sprayers and Hand Sprayers. Ready to use solution for the control of insects attacking livestock — Face Flies, Horn Flies, Stable Flies, House Flies, Gnats, Mosquitoes, Insect Control in Halls, Barns, Dairy Barns, Shelter Sheds, Milk Sheds, Restaurants, Warehouses, Food Processing Plants and Outdoor Areas such as Patios.

ACTIVE INGREDIENTS

BY WEIGHT

2,2-Dichlorovinyl Dimethyl phosphate	0.92%*
Related Compounds	0.08%*
Mineral Seal Oil	49.49%
Petroleum Distillates	49.49%
INERT INGREDIENTS	0.02%

Total

100.0%

*Equivalent to 1.0% Vapona Insecticide® T.M. Shell Chemical Co.

Do not reuse empty drum. Return to drum manufacturer or destroy by perforating or crushing and burying in a safe place.

WARNING

KEEP OUT OF REACH OF CHILDREN

See side panel for other cautions.

NET CONTENTS 30 GAL.

m



CATTLE (Horn and Dairy) to control Hornflies, Houseflies, Stable Flies, Face Flies, apply 1 to 2 fluid ounces of Dairy Mist Spray Solution per animal as a mist spray daily with hand or automatic sprayer. Apply spray so as to wet the hair. **DO NOT apply in excess of 2 fluid ounces per animal per day.**

FOR FOGGING: Use of a 1% dichlorvos fogging spray out of doors should be limited to use by or under the direction of trained personnel. The product should be applied at the rate of 0.05 to 0.1 pounds of actual dichlorvos per acre (5-10 pints per acre), to control Flies, Mosquitoes and Gnats. Apply dense fog when wind is calm. Do not contaminate food crops.

For fogging indoors for Flies, Gnats and Mosquitoes, use no more than one gallon every 64,000 cu. ft. When fogging indoors, eliminate all sources of ignition (pilot lights, fires, open flames) and disconnect electrical power. Before applying, reduce air movement as much as possible by closing doors, windows and other openings. Apply 1-5 seconds per 1000 cu. ft. **DO NOT** contaminate water, feed or foodstuffs, milk or milking utensils. **DO NOT** use this solution for fogging purposes in areas where animals have received direct application within 8 hours. The use of 1% dichlorvos (DDVP) spray in Food Processing Plants, Restaurants and Warehouses must be limited to application by Pest Control Operators. **DO NOT** treat food processing areas of food processing plants while in operation. **HOUSEHOLDS SHOULD BE TREATED BY A PEST CONTROL OPERATOR.**

Occupants should vacate premises before indoor treatments and should not reoccupy the premises until it has been ventilated. After application open several windows and doors to create air currents to ensure reduction of Vapona Insecticide vapors. Food should be removed and food handling equipment covered with wrapping paper during application, or washed after application is completed. During fogging or mist spraying indoors, use mask or respirator of a type passed by the U.S. Bureau of Mines for DDVP (Vapona Insecticide) protection. **DO NOT FOG INTO OR NEAR AN OPEN FLAME.**

ANIMAL BUILDINGS (Horse Barns, Dairy Barns, Shelter Sheds, Milk Sheds). To control Flies, Gnats, Mosquitoes, Roaches, Fleas, Ants, Sowbugs, Brown Dog Ticks, Ants, Spiders and Silverfish. Apply 1 pt. of 1.0% solution as a coarse, wet spray per 1,000 sq. ft. Treat around doorways, feed storage rooms, alleyways and window sills. For control of Ely Maggots in maggot breeding areas, apply at the rate of 1 to 2 pts. per 100 sq. ft.

Manufactured for



AGRICULTURAL PRODUCTS DIVISION
INTERNATIONAL MULTIFOODS CORPORATION
MINNEAPOLIS, MINNESOTA 55402

8166-1274p

Supera-weel

DAIRY MIST

quitoes, Roaches, Fleas, Ants, Sowbugs, Brown Dog Ticks, Ants, Spiders and Silverfish, treat **OUTDOOR LIVING AREAS, PICNIC GROUNDS, BACKYARD AREAS, PATIOS, LOADING DOCKS, OUTDOOR LATRINES, PARKING AREAS, REFUSE AREAS AROUND SERVICE STATIONS, OPEN AIR DRIVE-INS, ICE CREAM STANDS, GARBAGE COLLECTION AND DISPOSAL AREAS, RESTAURANTS, THEATERS, FOOD PROCESSING PLANTS, INDUSTRIAL PLANTS AND WAREHOUSES** using 1 pt. of 1% solution as a coarse, wet spray per 1,000 sq. ft.

DO NOT use in edible products areas of Food Processing Plants, Restaurants or other areas where food is commercially prepared or processed. **DO NOT** use in serving areas while food is exposed.

In treating Dog Kennels apply to outside dog runways, window sills and ledges. For control of fleas and brown dog ticks increase dosage to 1 qt. of Dairy Mist Solution per 1,000 sq. ft.

Treat infested areas, baseboards, cracks, walls, doors. **DO NOT** allow children in treated area until dry.

WARNING

May be fatal if swallowed, inhaled or absorbed through the skin or eyes. Wash thoroughly with soap and water after handling and before eating or smoking. **DO NOT** contaminate feed, water or foodstuffs. Keep out of reach of children. If swallowed call physician immediately. Do not get in eyes, on skin or on clothing. Do not breathe vapor spray mist. In case of contact flush skin and eyes with plenty of water. If eyes get medical attention. Wear a mask or respirator of a type passed by the U.S. BUREAU OF MINES for protection against DDVP. Wash contaminated clothing with soap and hot water before reuse.

NOTE TO PHYSICIAN: Antidotes are not available.

This product is toxic to fish, wildlife, and birds. Do not apply when weather conditions favor drift from areas treated. Keep out of water storage containers. Do not apply where runoff is likely to occur.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on the label.

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

E. P. A. REG. NO. 7455-21



DAIRY MIST

A SPRAY SOLUTION: For use in Foggers, Mist Sprayers and Hand Sprayers. Ready to use solution for the control of insects attacking livestock — Face Flies, Horn Flies, Stable Flies, House Flies, Gnats, Mosquitoes, Insect Control in Horse Barns, Dairy Barns, Shelter Sheds, Milk Sheds, Restaurants, Warehouses, Food Processing Plants and Outdoor Areas such as Patios.

ACTIVE INGREDIENTS

BY WEIGHT

2,2-Dichlorovinyl Dimethyl phosphate	0.92%*
Related Compounds	0.08%*
Mineral Seal Oil	49.49%
Petroleum Distillates	49.49%
INERT INGREDIENTS	0.02%
Total	100.0%

Equivalent to 1.0% Vapona Insecticide T.M. Shell Chemical Co.

Do not reuse empty drum. Return to drum manufacturer or destroy by perforating or crushing and burying in a safe place.

WARNING

KEEP OUT OF REACH OF CHILDREN

See side panel for other cautions.

NET CONTENTS 30 GAL.