

MINERAL 15-12 WITH RABON®

Supplement FEEDS

5620

20

MINRAL 15-12 WITH RABON® ORAL LARVICIDE

To prevent the development of horn flies, face flies in the manure of treated cattle.

ACTIVE INGREDIENTS:
2-Chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate

INERT INGREDIENTS:
Each Pound contains 54 grams Rabon®

GUARANTEED ANALYSIS
Calcium (Ca) Min. 14.0%
Phosphorus (P) at less than
Iodine (I), not fer. Min.
Salt (NaCl) Min. 90%

FEED INGREDIENTS: Cane Molasses, Soybean Meal, Calcium Carbonate, Dicalcium Phosphate, Potassium Sulfate, Magnesium Sulfate, Bentonite, Salt, Iron Oxide, Vitamin A Supplement, D-Ascorbic Acid, Selenium, Manganese Oxide, Iron Sulfate, Iron Carbonate, Copper Sulfate, Cobalt Carbonate, Zinc Oxide, Calcium Iodate, Magnesium Oxide, Calcium Sulfate

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If eyes wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS
This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL
Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Pesticide that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Consult local, state or local disposal authorities for approved alternative methods such as timed open burning.

EPA Reg. No. 7455-32
EPA Est. No. 7455-1A-01
NET WEIGHT 50 LBS.
U.S. Patent No. 2,988,812

5620

Manufactured by
MULTIFOODS
AGRI-CULTURAL PRODUCTS DIVISION
MINNEAPOLIS, MN 55412

SEE BACK OF LABEL FOR DIRECTIONS FOR USE

1 2 3 4 5 6 7 8 9 0 1 2 3

DIRECTIONS FOR USE: GENERAL CLASSIFICATION

It is a violation of the Federal Law to use this product in a manner inconsistent with its labeling.

Allow free-choice to cattle. Cattle should consume an average of 1/2 lb. of the mineral per 100 pounds of body weight per month. This will supply the recommended average daily intake of 70 mg. RABON® Oral Larvacide per hundred pounds of body weight. Animals should not be starved for minerals before initiation of SUPERSWEET Mineral 15-12 With RABON® Oral Larvacide program but OMIT all salt from other feed after the program is started. Do not feed other free-choice sources of salt or mineral containing salt in any form. Allow one mineral feeder for each 10-15 head and place near watering and loafing area. Put out no more than one month's supply at a time and protect from rain.

If consumption of SUPERSWEET Mineral 15-12 With RABON® Oral Larvacide is above 1/2 lb. per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below 1/2 lb. per 100 lbs. body weight per month, increase the number of feeding locations. Start feeding SUPERSWEET Mineral 15-12 With RABON® Oral Larvacide early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

SUPERSWEET Mineral 15-12 With RABON® Oral Larvacide may be fed up to slaughter and to lactating dairy cows.

SUPERSWEET Mineral 15-12 With RABON® Oral Larvacide prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to supplement control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, SUPERSWEET Mineral 15-12 With RABON® Oral Larvacide should be used in conjunction with other good management and sanitation practices.

DOSAGE

The following chart shows recommended ounces per day for various body weights:

Lbs Body Weight	Oz./ Head/Day	Lbs Body Weight	Oz./ Head/Day
700	1.9	1200	3.2
800	2.1	1300	3.5
900	2.4	1400	3.7
1000	2.7	1500	4.0
1100	2.9		

Rabon is the registered trademark of Shell Chemical Co.

0295

ACCEPTED
7455-32
MAR 5 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7455-32

A B C D E F G H I J K L M