

MINRAL 24-6 WITH RABON®

5621
Supersweet
FEEDS
MINRAL 24-6 WITH
RABON® ORAL LARVACIDE
5621

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle

ACTIVE INGREDIENTS:

| | |
|-------------------------------------|-----------|
| 2-Chloro-1-(2,4,5-trichlorophenyl)- | BY WEIGHT |
| Vinyl dimethyl phosphate* | 1.00% |
| INERT INGREDIENTS | 99.00% |

*Each Pound contains 4.54 grams RABON®

GUARANTEED ANALYSIS

| | |
|------------------------------|------------------------|
| Calcium (Ca) ... Min 22.0% | Max 26.0% |
| Phosphorus (P) not less than | Max 8.0% |
| Iodine (I) not less than | Max 0.002% |

Salt (NaCl) Min 80% Max 10.5%

FEED INGREDIENTS: Cane Molasses, Calcium Carbonate, Dicalcium Phosphate, Potassium Sulfate, Magnesium Sulfate, Bentonite, Salt, Iron Oxide, Vitamin A Supplement, D-Activated Animal Sterol, Manganese Oxide, Iron Sulfate, Iron Carbonate, Copper Sulfate, Cobalt Carbonate, Zinc Oxide, Calcium Iodate, Magnesium Oxide, Calcium Sulfate

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION**

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS
This product is toxic to fish. Keep cut flakes, streams or ponds. Do not contaminate water by clearing of equipment or disposal of wastes.

STORAGE AND DISPOSAL
Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Pesticide that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

EPA Reg. No. 7455-33
EPA Est. No. 7455-1A-01
5621

NET WEIGHT 50 LBS
U.S. Patent No. 3,101,642

5621

15

Manufactured by
MULTIFOODS
MULTIFOODS CORPORATION
1000 N. Cicero Avenue
Skokie, IL 60077

SEE BACK OF LABEL FOR DIRECTIONS FOR USE

1 2 3 4 5 6 7 8 9 0 - 2 3

**DIRECTIONS FOR USE:
GENERAL CLASSIFICATION**

It is a violation of the Federal Law to use this product in a manner inconsistent with its labeling.

Allow free-choice to cattle. Cattle should consume an average of $\frac{1}{2}$ lb. of the mineral per 100 pounds of body weight per month. This will supply the recommended average daily intake of 70 mg. RABON[®] Oral Larvicide per hundred pounds of body weight. Animals should not be starved for minerals before initiation of SUPERSWEET Mineral 24-6 With RABON[®] Oral Larvicide program but OMIT all salt from other feed after the program is started. Do not feed other free-choice sources of salt or mineral containing salt in any form. Allow one mineral feeder for each 10-15 head and place near watering and loafing area. Put out no more than one month's supply at a time, and protect from rain.

If consumption of SUPERSWEET Mineral 24-6 With RABON[®] Oral Larvicide is above $\frac{1}{2}$ lb. per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below $\frac{1}{2}$ lb. per 100 lbs. body weight per month, increase the number of feeding locations. Start feeding SUPERSWEET Mineral 24-6 With RABON[®] Oral Larvicide early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

SUPERSWEET Mineral 24-6 With RABON[®] Oral Larvicide may be fed up to slaughter and to lactating dairy cows.

SUPERSWEET Mineral 24-6 With RABON[®] Oral Larvicide prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, SUPERSWEET Mineral 24-6 With RABON[®] Oral Larvicide should be used in conjunction with other good management and sanitation practices.

DOSAGE

The following chart shows recommended ounces per day for various body weights.

| Lbs. Body Weight | Oz./Head/Day | Lbs. Body Weight | Oz./Head/Day |
|------------------------|--------------|------------------------|--------------|
| 700 | 1.9 | 1200 | 3.2 |
| 800 | 2.1 | 1300 | 3.5 |
| 900 | 2.4 | 1400 | 3.7 |
| 1000 | 2.7 | 1500 | 4.0 |
| 1100 | 2.9 | | |

Rabon is the registered trademark of Shell Chemical Co.

5621

| | |
|---|---------|
| ACCED | 7455-33 |
| MAR 5 1981 | 16 |
| Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for use as a registered insecticide. | |
| EPA Reg. No. 7455-33 | |

A B C D W U G H - J K L M