

7467-51

5-4-77

113

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567) WASHINGTON, D.C. 20460	EPA REGISTRATION NO. 7467-51	DATE OF ISSUANCE MAY 1 1977
	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT	

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
 (Under the Federal Insecticide, Fungicide,
 and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Best Available Copy

Valley 20 241 Hill
 Attention: Mr. Hill
 Chemicals Division
 Post Office Box 1310
 Harlingen, Texas 79560

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Note that this submission was processed and accepted under the 1947 Federal Insecticide, Fungicide, and Rodenticide Act. If such time as re-registration is required or amendments are proposed, the Registration, Re-registration and Classification procedures, as published in the Federal Register on July 2, 1973, will be applied. Refer to Section 2.2.22 of that document. Refer also to PL Notice 75-1 and 75-4.

Robert J. Taylor
 Product Manager (25)
 Fungicide-Herbicide Branch
 Registration Division (WH-567)

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

DESTROY EMPTY CONTAINER - PERFORATE - CRUSH AND BURY

CAUTION

KEEP OUT OF REACH OF CHILDREN

Do Not Get in Eyes, on Skin or on Clothing.
Wash hands thoroughly with soap and warm water after handling and before eating or smoking.
Wear clean clothing daily.
Do not contaminate feed, food stuffs or water.
Do not allow spray to drift to other crops, plants or trees that may be defoliated.
Wash and clean spray equipment after each day's use and when spraying is completed. Do not allow wash water to contact desirable plants.
Do not contaminate water by cleaning of equipment or disposal of wastes.
Apply this product only as specified on this label.
Rinse and drain empty containers into the spray tank at least three successive times; then puncture, crush and bury the container in a safe place.
NEVER REUSE.
Harmful if swallowed.

GENERAL INFORMATION

READ THE ENTIRE LABEL. USE ONLY IN ACCORDANCE WITH THE DIRECTIONS GIVEN.
CLIMAX-3 is designed to reduce the moisture content in grain sorghum prior to harvest.
APPLICATION: CLIMAX-3 may be applied by aircraft or ground equipment. Thorough coverage of the crop is essential. Low temperatures require more time for CLIMAX-3 to act and requires the higher rate.

NEVER RE-USE CONTAINER OR KEEP ON PREMISES

Net
Contents
5 Gals.

VALCO
CHEMICAL

CLIMAX-

COTTON DEFOLIANT AND GRAIN SORGHUM DES

ACTIVE INGREDIENTS:

Sodium Chlorate

INERT INGREDIENTS:

A CONCENTRATED LIQUID SOLUTION CONTAINING 3 POUNDS OF SODIUM CHLORATE WITH A FIRE INHIBITOR. DOES NOT CONTAIN BORON IN A

EPA Reg. No. 7467-51

® Trade Mark Reg. U. S. Patent Office U. S. Reg. No. 7467-37

EPA Establishment No. 07467TX01

CONDITIONS OF SALE

Neither the manufacturer nor the seller makes any guarantee, representation or warranty on this material, except that it substantially conforms with the chemical description on this label. From non-conformance thereto shall be limited to direct damages, shall not exceed the purchase price to buyer or any other person resulting from the handling, storage or use of this product from residues or decomposition compounds remaining after use. The buyer and/or user assumes the suitability of this material for his particular purposes, on the basis of his own investigation and liability resulting from the handling, storage, or use of this product whether used or not. No person is authorized to vary or waive any of these conditions.

CAUTION: Keep out of reach of children
See Left Panel for additional Cautions.

Manufactured by **VALCO CHEMICAL DIVISION**
VALLEY CO-OP OIL MILL

DO NOT STORE NEAR FOOD OR FEED PRODUCTS

DIRECTIONS FOR USE

GRAIN, SORGHUM AND MILQ: To reduce the moisture content in grain while the crop is standing, use 1 1/2 to 2 gallons of CLIMAX-3 per acre in 5 to 6 gallons of total solution per acre by air or 1 1/2 to 2 gallons of CLIMAX-3 per acre in 20 to 30 gallons of total solution per acre with suitable ground equipment. Make application 7 to 10 days before anticipated harvesting date. Use the lower rate when grain sorghum and weather are right for rapid desiccation. Swath width should not exceed wing span; altitude and speed of aircraft should be regulated to give best coverage. Do not apply during windy weather as spray drift to susceptible nearby crops may cause damage. Do not apply during very high temperatures.

DO NOT graze treated areas or feed treated fodder, forage or residual seeds within 14 days of application.

COTTON: STAGE of plant growth and prevailing weather conditions markedly affect the defoliation process. Best results are obtained when soil fertility and moisture are adequate, the boll set is heavy, all bolls are mature, vegetative growth has stopped and there is little or no second growth. LEAVES SHOULD BE GREEN AND TURGID. Bolls less than 35 days old may not mature after application of this product. Apply 2 to 3 weeks before anticipated picking date on clear days when temperatures will go above 70° F. Do not apply under conditions of extreme heat during middle of day. Low temperatures require more time and the higher rate for acceptable defoliation.

DO NOT graze treated area or feed gin waste to livestock.

APPLICATION: One to 1 1/2 gallons of CLIMAX-3 per acre in enough water to give thorough coverage may be applied by aircraft or ground equipment.

NOTE: DO NOT apply within 7 days of harvest.

For airplane application mix required amount of CLIMAX-3 in 5 to 10 gallons of total solution per acre. Use the lower rate when conditions for good defoliation prevail. Swath width should not exceed wing span; altitude and speed of aircraft should be regulated to give best spray coverage. Do not apply during windy weather as spray may drift to susceptible nearby crops and cause damage.

For ground equipment, use 10 to 20 gallons per acre of ready-to-apply material per acre. Prevailing conditions will govern the rate of CLIMAX-3 to use.

KEEP AWAY FROM HEAT AND OPEN FLAME

3/3