

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS**

**WARNING:** Poisonous if swallowed, inhaled or absorbed through skin. Do not swallow or get in eyes or on clothing. Do not breathe vapors or mist. Keep out of reach of children and domestic animals.

Wear clean rubber gloves, clean protective clothing, and goggles or face shield. Replace gloves frequently and destroy used gloves.

Keep all unprotected persons out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Workers entering field within 16 hours of treatment should be protected with clean protective clothing. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water for at least 15 minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before re-use. Discard and destroy articles made of leather which become contaminated because they can not be decontaminated.

**ANTIDOTE AND FIRST AID TREATMENT**

**INTERNAL:** If toxaphene is swallowed, do not induce vomiting. See "Physician's Note." Never give anything by mouth to an unconscious person.

**EXTERNAL:** Remove contaminated clothing and wash skin thoroughly with soap and water. If in eyes, flush with clean water for at least 15 minutes.

**CALL A PHYSICIAN IMMEDIATELY:** In all cases of suspected poisoning.

**PHYSICIAN'S NOTE:** If swallowed, use gastric lavage. Avoid aspiration. Toxaphene produces central nervous system excitation that responds to barbituric acid derivatives. Prevent or control convulsions with barbiturates, preferably administered intravenously. If the barbiturate can be given before convulsions start, sodium pentobarbital is the drug of choice. Since toxaphene has a fairly long duration of action, a long-acting barbiturate such as phenobarbital may be used after initial control of convulsions. Refer to "Clinical Handbook on Economic Poisons," which was prepared by the U.S. Department of Health, Education, and Welfare, Public Health Service, Communicable Disease Center, Atlanta, Georgia.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and other wild life. Keep out of lakes, ponds, and streams.

This product is toxic to bees and should not be applied when bees are actively visiting the area. Do not apply in any manner not specified on the label. Birds and other wildlife in treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL AND CHEMICAL HAZARDS**  
**DO NOT USE, POUR, OR STORE NEAR HEAT OR OPEN FLAME.** In case of spillage, use absorbent clay, hydrated lime, or other solids to soak up any liquid. Scrape up excess and bury residue away from water supplies. Wash area with strong detergent and hot water.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Triple rinse containers and dispose of in an approved landfill or bury in a safe place. Consult federal, state or local authorities for approved alternate procedures.

**ACCEPTED**

FEB 26 1981

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No.

Net  
Contents  
Gal.



## TOXAPHENE LIQUID - 6

ACTIVE INGREDIENTS:	By Wt.
Toxaphene (Technical Chlorinated Camphene, Chlorine Content 67-69%) .....	88.00%
Xylene Range Aromatic Petroleum Derivative Solvents .....	37.38%
INERT INGREDIENTS: .....	4.61%
Total .....	100.00%

CONTAINS 8 POUNDS TOXAPHENE PER GALLON

EPA EST. No. 07467-TX-01  
EPA Reg. NO. 07467-55

### STATEMENT OF PRACTICAL TREATMENT

CALL A PHYSICIAN IN ALL CASES OF SUSPECTED POISONING.

If swallowed, Do not induce vomiting. See "Physician's Note."

Never give anything by mouth to an unconscious person.

If on skin, Remove contaminated clothing and wash skin thoroughly with soap and water.

If in eyes, Flush with clean water for at least 15 minutes.

See side panels for additional precautionary statements.

Keep Out Of Reach Of Children!

# WARNING

Manufactured by

**VALCO CHEMICALS DIVISION**

Harlingen, Texas 78550

VALLEY CO-OP OIL MILL

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with directions on the label.

**COTTON-** For SEEDLING PLANTS against Cutworms, use 2½ to 5½ pints per acre and spray plants as well as the soil around the plants. Against Thrips use 2 to 2½ pints per acre.

During EARLY FRUITING STAGE against overwintered Boll Weevil, Cotton Fleahopper, Lygus Bugs, and Cotton leafworm, use 2½ pints per acre, increasing to 3 pints during MID-FRUITING SEASON, and to 5½ pints during Boll Weevil Migration or when weevils are heavy. Begin spraying cotton when injury or insects threaten to become important and continue every four to five days (7 days for Fleahopper and Lygus Bug). Mix the required dosage per acre of insecticide with the number of gallons of water per acre required for good plant coverage with the sprayer used, usually 3 gallons per acre.

Do not graze or feed gin waste to dairy animals or animals being finished for slaughter if fields are treated late in the season.

**VEGETABLE CROPS** - Apply in at least 3 gallons in water per acre at the rate per acre given below for the crops and insects. To avoid excessive residues on edible portions, observe maximum rates and waiting period given.

**Beans** - Against Flea Beetles, Leafhoppers, Thrips, Loopers, use 1½-5½ pints. Do not apply after pods begin to form. Do not feed treated vines to dairy animals or to animals being finished for slaughter.

**Cabbage, Broccoli, Cauliflower** - Against Flea Beetles, use 4 pints per acre; Cutworms (spray soil) use 2 pints; Cabbage Worms use 2½-4 pints per acre. Do not use after edible parts begin to form, or within 30 days of harvest. Do not feed treated plants to dairy animals or to animals being finished for slaughter.

**Corn** - Against Cutworms, use 2½ to 4 pints per acre and spray plants as well as the soil around the plants. Do not feed treated forage to dairy animals or to animals being finished for slaughter.

**Carrots** - Against Flea Beetles use 4 pints; Cutworms use 2½-4 pints per acre. Apply as foliar treatment only. Do not feed treated foliage to dairy animals or to animals being finished for slaughter.

**Eggplant** - Against Flea Beetles and Cucumber Beetles use 1½-4 pints per acre and not within 5 days before harvesting.

**LETTUCE** - Against Cucumber Beetles, Cutworms, Loopers, use 2½-5½ pints per acre. Do not apply after seedling stage for **ARDE** lettuce, nor after heads begin to form on head lettuce.

**Onions** - Against Thrips, use 2½ pints per acre. Do not apply to green or spring onions.

**Peas** - Against Cowpea Curculio and Loopers, use 2-4 pints per acre. Do not apply after pods begin to form if pods are to be used as food, or within 7 days of harvest if pods are not to be used as food. Do not feed vines to dairy animals or animals being fattened for slaughter.

**PEPPERS** - Against Cutworms (spray soil), Flea Beetles, Leafhoppers and Pepper Weevil, use 1½-4 pints per acre. Do not apply within 5 days of harvest. *thor*

**Tomatoes** - Against Cutworms (spray soil), Flea Beetles, Hornworms, and Stink Bugs, use 2½ to 6½ pints per acre. Do not apply within 3 days of harvest.

#### CONDITIONS OF SALE

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material, not in strict accordance with the directions given on the label.