



Please Read Entire Label Before Using
Citrus (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangelos, Tangerines) - Citrus flat, Citrus red (purple), citrus rust, six-spotted, Texas citrus and Yuma mites.

Refer to State Agricultural Experiment Station or State Agricultural Extension Specialist for proper timing and number of applications.

DO NOT USE ON AVOCADOS or EGG PLANT STRAWBERRIES - Six-spotted, strawberry (Atlantic) and two-spotted mites. Apply when mites appear and repeat as necessary. For cyclamen mite apply 2 or 3 thorough, high-pressure, drenching sprays (400 gal. per acre) at 10 to 20 day intervals when new growth starts in spring, or use as a spot treatment in infested areas during production season.

ROSES, ORNAMENTALS, FLOWERS, NURSERY STOCK & SHADE TREES - For the control of mites common on greenhouse or outdoor plants. Use 1½ - 2 tsp. in 1 gal. of water. Apply when mites are present and repeat as necessary. Do not apply to Chinese holly (ilex cornuta) as leaf spotting may result.

To avoid excessive residue at harvest do not exceed the maximum dosage, or the time restriction between last application and harvest for each crop group listed in the table.

Gallonage of spray will vary with size and number of trees or plants. In general, Kelthane applications should be made when the various mite species appear in threatening numbers and repeated as necessary for adequate seasonal control, except as noted herein or recommended locally. Repackaged and Distributed by

CHEM PAK COMPANY
 Miami, Florida 33143

EPA Reg. No. 7478-27 EPA Est. No. 7478-Ga-1

KELTHANE

Home Garden
**AGRICULTURAL
 MITICIDE**

EMULSIFIABLE CONCENTRATE

KILLS MITES

DOES NOT KILL BEES OR BENEFICIAL INSECT PREDATORS

ACTIVE INGREDIENTS:

1,1-Bis(chlorophenyl)-2,2,2 trichloro-ethanol* 18.5%
 Xylene 73.0%

INERT INGREDIENTS

8.5%
 Total 100.0%

*Equivalent to 1.6 lbs Active Ingredient per gallon.

Keep out of reach of children.
 See Additional Cautions on Inside Panel.

NET CONTENTS 1 PINT (16 Fluid Ounces)

DIRECTIONS FOR USE: Kelthane gives both quick and residual kill. Does not harm bees. It is compatible with most commonly used insecticides and fungicides. Do not use with lime. Kelthane should be applied thoroughly using sufficient gallonage to cover all parts of the plant, especially the undersides of leaves. Summary of Kelthane EC Dosage Rates & Limitations Between Last Application & Harvest

C R O P	Per Gal.	Days before Harvest
Strawberries - Two-spotted and similar mites	2-4 tsp.	2
Cyclamen mite	3 tsp.	2
(Do not exceed 1½ qts. to 100 gal. water - Do not exceed 6 qts. per acre.)		
Roses, ornamentals, flowers, nursery, (full coverage)	1½-2 tsp.	
Citrus (all)	1½-2 tsp.	7

(Do not exceed 2½ gal. Kelthane EC per acre. Use 1 qt. to 100 gal. water.)

tsp = teaspoon / TBS = tablespoon

2 tsp. to a gallon - 1 qt. to 100 gal. water.

CAUTION: Avoid skin contact. Wash thoroughly after using. Avoid inhalation of mists. Harmful if swallowed. If taken internally get medical attention. Do not store near feed or foodstuffs.

Do not use, pour, spill or store near heat or open flame. Do not reuse empty container - dispose of properly.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

CAUTION: Important - this product is toxic to fish, shrimp, crab and other aquatic animals. Keep out of any body of water. Apply this product only as specified on this label.

