



GARDEN & HOME General Purpose Insecticidal Spray

Contains Methoxychlor 50% Wettable Powder

FOR CONTROL OF CERTAIN INSECT PESTS OF
VEGETABLES, FRUIT, ROSES, ORNAMENTALS & FLOWERS

Active Ingredients: Methoxychlor, technical* 50.0%
Inert Ingredients 50.0%
*Equivalent to 34% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloro-
ethane and 6% other isomers and related compounds.


**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Side Panels for Additional Precautionary Statements

EPA Est. No. 7478 QA-1

EPA Reg. No. 7472-4b

DISTRIBUTED BY
CNEM PAK CO.
Rt 1, Box 144, McRae, GA 31055



Directions for Use (continued)

VEGETABLE INSECTS - Asparagus beetles on asparagus, bean leaf beetles on snap or navy beans, flea beetles on tomatoes, Colorado potato beetles on potatoes, cucumber beetles on cucumbers, fall armyworms on beans, corn, tomatoes, flea beetles on cabbage, corn, potatoes, fruit worms on tomatoes, pea weevils on peas, Japanese beetles on beans, corn and tomatoes, leafhoppers on beans and carrots, Mexican bean beetle on snap or navy beans, rose chiggers on beans, corn, peppers, wire beetles on squash.

Asparagus beetles on asparagus, bean leaf beetles on snap or navy beans, flea beetles on tomatoes, Colorado potato beetles on potatoes, cucumber beetles on cucumbers, fall armyworms on beans, corn, tomatoes, flea beetles on cabbage, corn, potatoes, fruit worms on tomatoes, pea weevils on peas, Japanese beetles on beans, corn and tomatoes, leafhoppers on beans and carrots, Mexican bean beetle on snap or navy beans, rose chiggers on beans, corn, peppers, wire beetles on squash.

Do not apply to asparagus, summer squash, tomatoes, or peppers within 14 days of harvest. Do not apply to carrots within 7 days of harvest. Do not apply to beans, corn, and peas within 14 days of harvest.

FRUIT INSECTS - Plum curculio on peaches, pears, plums and aprins. Apply 2 TBS per gallon of water as thorough cover spray. Make 2 applications at 7 day intervals. Do not use on peaches within 7 days of harvest. Do not use on apples, pears, plums within 7 days of harvest.

Japanese beetle, leafhoppers, leaf beetles, and other insects on cherry, plums, peaches, grapes, raspberries and strawberries, use 2 TBS per gallon of water as a thorough cover spray when insects first appear. Repeat application at 7 day intervals until infestation warrants control is complete. Do not spray within 14 days of harvest.

FLOWERS, ORNAMENTALS - Roses, Shrubs - Japanese Beetles, Sawflies, Cuckoo's Spite, Flea Beetles, Leaf Beetles, Rose Chiggers, Rose Suckers, Leafhoppers, Rose Chiggers, Rose Suckers, Leafhoppers. Apply 2 TBS per gallon of water as a thorough cover spray when insects first appear. Repeat application at 7 day intervals until infestation warrants control is complete.

Notice: User assumes all risk of use, storage or handling of this material and its strict liability. A full instruction given herewith.

NET WEIGHT 1 Pound (16 oz.)

NO
MALS
breathing dust. Wash
ENT
Take person with you
entire use. Insect
Do not apply when
contaminate water by
ornamental fish
Applied when bees are
when not in use. Store
put in trash collection
stent with its labeling
in a few hours. Keep
pounds to 100 gallons
in spray is most effect
ity and other environ
cover part of the day
in large areas and thir
ives. Under severe con