GARDEN & HOME

General Purpose Insecticidal Dust

Contains Methoxychlor 10%

FOR THE CONTROL OF CERTAIN INSECTS
ON LISTED
VEGETABLES – SMALL FRUITS – FLOWERS
ORNAMENTALS

Active Ingredients	By Weight
Methoxychlor Technical* . . .	10.0%
Inert Ingredients	90.0%

**Equivalent to 8.8% 2,2-bis (bis (p-methoxyphenyl)-1,1,1 trichloroethane and 1.2% other isomers and related compounds.*

Use on Tomatoes, Beans, Corn, Cucumbers, Strawberries

KEEP OUT OF REACH OF CHILDREN
CAUTION
See Additional Cautions on Other Panels

EPA Est. No. 7478-GA-1 EPA Reg. No. 7478-46

DISTRIBUTED BY
CHEM-PAK CO.
Rt 1, Box 144, McRae, GA 31055

NET WEIGHT 12 OUNCES

NOTICE: User assumes all risk of use, storage or handling of this material not in strict accordance with direction given herewith

PLEASE READ ENTIRE LABEL BEFORE USING
PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION

[illegible] Avoid breathing [illegible]

STATEMENT OF PRACTICAL TREATMENT

[illegible] amount of water to drink. Make person [illegible]
[illegible] wash eyes with fresh water for 15 minutes. If irritation contin[illegible]
[illegible] wash with [illegible]

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water. Do not apply when weather conditions favor drift from area being treated. Do not contaminate water by cleaning of spray equipment or disposal of waste. Do not contaminate ornamental fish ponds.

BEE CAUTION This product is toxic to bees and should not be applied when bees are actively visiting the area.

STORAGE and DISPOSAL DIRECTIONS

Avoid storage near feed and food products. Keep container closed when not in use. Store in a dry place.

Disposal Do not reuse empty container. Wrap in newspaper and put in trash collection.

DIRECTIONS FOR USE

Begin application at first sign of infestation (for fruits, begin at petal fall) and repeat at 7 to 14 day intervals or as needed. Do not apply to crops within the number of days specified before harvest or grazing as shown by PHL (re Harvest Limitation).

Gardens Apply dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. Repeat as necessary, generally at 7–14 day intervals. Apply with a good dust gun. Do not apply in excess. Apply when foliage is dry. Adjust amount applied for complete coverage.

Small Fruit Blueberry: Japanese Beetle, Leafhoppers, PHL 14 days. Blackberry, Raspberry, Strawberry: Rose Chafer, Strawberry Weevil, Flea Beetles, Spittlebugs, Japanese Beetles, PHL 14 days, 10–15 lbs per acre.

Vegetables Beans, Black-eyed Peas: Corn Earworm, Cucumber Beetles, Fall Armyworm, Alfalfa and Garden Webworms, Flea Beetles, Mexican Bean Beetle, Potato Leafhopper, Bean Leaf Beetle, Japanese Beetle, PHL 7 days. Apply within 3 days of harvest. If vines are not used for food or forage. As a general rule 15 to 20 pounds to an acre every 7–14 days is the usual application. Adjust according to size of plants.

Cucumber, Melons, Summer Squash: Squash Vine Borers, Fall Armyworm, Cucumber Beetle, Flea Beetle, apply 10–20 lbs per acre depending on size of vines. PHL 1 day.

Potato (Irish): Colorado Potato Beetle, Flea Beetles, Fall Armyworm, Leafhoppers, Blister Beetles, PHL (None) 10–20 pounds per acre.

Sweet Potato: Fall Armyworm, Flea Beetles, PHL None, 15–20 lbs per acre. Adjust for complete coverage.

Tomato: Colorado Potato Beetle, Flea Beetles, Fall Armyworm, Leafhoppers, Blister Beetle, PHL 1 day, 15–20 lbs per acre. Adjust for complete coverage.

Flowers & Ornamentals such as roses, chrysanthemums, dahlias, shrubs, shade trees: Blister Beetles, Cankerworms, Cucumber Beetles, Flea Beetles, Fleahoppers, Flower Thrips, Japanese Beetle, Leafhoppers, Rose Chafer, Rose Slugs, Sawflies, Tent Caterpillars. Begin applications when insects [illegible] and repeat at 7 to 14 day intervals or as needed for [illegible] coverage.